



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/31, C07K 14/35, A61K 39/04, 48/00, 49/00, C12N 15/62, C07K 19/00, G01N 33/50, 33/60, 33/569, C12N 1/19, 1/20, 1/21, 5/10	A2	(11) International Publication Number: WO 98/16646 (43) International Publication Date: 23 April 1998 (23.04.98)
(21) International Application Number: PCT/US97/18293 (22) International Filing Date: 7 October 1997 (07.10.97) (30) Priority Data: 08/730,510 11 October 1996 (11.10.96) US 08/818,112 13 March 1997 (13.03.97) US (71) Applicant: CORIXA CORPORATION [US/US]; 1124 Columbia Street, Seattle, WA 98104 (US). (72) Inventors: REED, Steven, G.; 2843 - 122nd Place N.E., Bellevue, WA 98005 (US). SKEIKY, Yasir, A., W.; 8327 - 25th Avenue N.W., Seattle, WA 98107 (US). DILLON, Davin, C.; 21607 N.E. 24th Street, Redmond, WA 98053 (US). CAMPOS-NETO, Antonio; 9308 Midship Court N.E., Bainbridge Island, WA 98021 (US). HOUGHTON, Raymond; 2636 - 242nd Place S.E., Bothell, WA 98021 (US). VEDVICK, Thomas, S.; 124 South 300th Place, Federal Way, WA 98003 (US). TWARDZIK, Daniel, R.; 10195 South Beach Drive, Bainbridge Island, WA 98110 (US). LODES, Michael, J.; 9223 - 36th Avenue S.W., Seattle, WA 98126 (US).	(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US). (81) Designated States: AL, AM, AT, AU, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: COMPOUNDS AND METHODS FOR IMMUNOTHERAPY AND DIAGNOSIS OF TUBERCULOSIS		
(57) Abstract <p>Compounds and methods for inducing protective immunity against tuberculosis are disclosed. The compounds provided include polypeptides that contain at least one immunogenic portion of one or more <i>M. tuberculosis</i> proteins and DNA molecules encoding such polypeptides. Such compounds may be formulated into vaccines and/or pharmaceutical compositions for immunization against <i>M. tuberculosis</i> infection, or may be used for the diagnosis of tuberculosis.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

COMPOUNDS AND METHODS FOR IMMUNOTHERAPY AND DIAGNOSIS OF TUBERCULOSIS

5 TECHNICAL FIELD

The present invention relates generally to detecting, treating and preventing *Mycobacterium tuberculosis* infection. The invention is more particularly related to polypeptides comprising a *Mycobacterium tuberculosis* antigen, or a portion or other variant thereof, and the use of such polypeptides for diagnosing and vaccinating
10 against *Mycobacterium tuberculosis* infection.

BACKGROUND OF THE INVENTION

Tuberculosis is a chronic, infectious disease, that is generally caused by infection with *Mycobacterium tuberculosis*. It is a major disease in developing
15 countries, as well as an increasing problem in developed areas of the world, with about 8 million new cases and 3 million deaths each year. Although the infection may be asymptomatic for a considerable period of time, the disease is most commonly manifested as an acute inflammation of the lungs, resulting in fever and a nonproductive cough. If left untreated, serious complications and death typically result.

20 Although tuberculosis can generally be controlled using extended antibiotic therapy, such treatment is not sufficient to prevent the spread of the disease. Infected individuals may be asymptomatic, but contagious, for some time. In addition, although compliance with the treatment regimen is critical, patient behavior is difficult to monitor. Some patients do not complete the course of treatment, which can lead to
25 ineffective treatment and the development of drug resistance.

Inhibiting the spread of tuberculosis requires effective vaccination and accurate, early diagnosis of the disease. Currently, vaccination with live bacteria is the most efficient method for inducing protective immunity. The most common *Mycobacterium* employed for this purpose is *Bacillus Calmette-Guerin* (BCG), an
30 avirulent strain of *Mycobacterium bovis*. However, the safety and efficacy of BCG is a source of controversy and some countries, such as the United States, do not vaccinate

the general public. Diagnosis is commonly achieved using a skin test, which involves intradermal exposure to tuberculin PPD (protein-purified derivative). Antigen-specific T cell responses result in measurable induration at the injection site by 48-72 hours after injection, which indicates exposure to Mycobacterial antigens. Sensitivity and specificity have, however, been a problem with this test, and individuals vaccinated with BCG cannot be distinguished from infected individuals.

While macrophages have been shown to act as the principal effectors of *M. tuberculosis* immunity, T cells are the predominant inducers of such immunity. The essential role of T cells in protection against *M. tuberculosis* infection is illustrated by the frequent occurrence of *M. tuberculosis* in AIDS patients, due to the depletion of CD4 T cells associated with human immunodeficiency virus (HIV) infection. Mycobacterium-reactive CD4 T cells have been shown to be potent producers of gamma-interferon (IFN- γ), which, in turn, has been shown to trigger the anti-mycobacterial effects of macrophages in mice. While the role of IFN- γ in humans is less clear, studies have shown that 1,25-dihydroxy-vitamin D3, either alone or in combination with IFN- γ or tumor necrosis factor-alpha, activates human macrophages to inhibit *M. tuberculosis* infection. Furthermore, it is known that IFN- γ stimulates human macrophages to make 1,25-dihydroxy-vitamin D3. Similarly, IL-12 has been shown to play a role in stimulating resistance to *M. tuberculosis* infection. For a review of the immunology of *M. tuberculosis* infection see Chan and Kaufmann in *Tuberculosis: Pathogenesis, Protection and Control*, Bloom (ed.), ASM Press, Washington, DC, 1994.

Accordingly, there is a need in the art for improved vaccines and methods for preventing, treating and detecting tuberculosis. The present invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

Briefly stated, this invention provides compounds and methods for preventing and diagnosing tuberculosis. In one aspect, polypeptides are provided comprising an immunogenic portion of a soluble *M. tuberculosis* antigen, or a variant of such an antigen that differs only in conservative substitutions and/or modifications. In

one embodiment of this aspect, the soluble antigen has one of the following N-terminal sequences:

- (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Cys-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu; (SEQ ID No. 120)
- 5 (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser; (SEQ ID No. 121)
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg; (SEQ ID No. 122)
- (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro; (SEQ ID No. 123)
- 10 (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val; (SEQ ID No. 124)
- (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro; (SEQ ID No. 125)
- 15 (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Thr-Ala-Ala-Ser-Pro-Pro-Ser; (SEQ ID No. 126)
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly; (SEQ ID No. 127)
- (i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Leu-Thr-Ser-Leu-Leu-Asn-Ser-Leu-Ala-Asp-Pro-Asn-Val-Ser-Phe-Ala-Asn; (SEQ ID No. 128)
- 20 (j) Xaa-Asp-Ser-Glu-Lys-Ser-Ala-Thr-Ile-Lys-Val-Thr-Asp-Ala-Ser; (SEQ ID No. 134)
- (k) Ala-Gly-Asp-Thr-Xaa-Ile-Tyr-Ile-Val-Gly-Asn-Leu-Thr-Ala-Asp; (SEQ ID No. 135) or
- 25 (l) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly; (SEQ ID No. 136)

wherein Xaa may be any amino acid.

In a related aspect, polypeptides are provided comprising an immunogenic portion of an *M. tuberculosis* antigen, or a variant of such an antigen that

30

differs only in conservative substitutions and/or modifications, the antigen having one of the following N-terminal sequences:

- (m) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID No. 137) or
- 5 (n) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID No. 129)

wherein Xaa may be any amino acid.

In another embodiment, the soluble *M. tuberculosis* antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID Nos.: 1, 2, 4-10, 13-25, 52, 99 and 101, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID Nos.: 1, 2, 4-10, 13-25, 52, 99 and 101 or a complement thereof under moderately stringent conditions.

In a related aspect, the polypeptides comprise an immunogenic portion of a *M. tuberculosis* antigen, or a variant of such an antigen that differs only in conservative substitutions and/or modifications, wherein the antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID Nos.: 26-51, 138, 139, 163-183 and 201, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID Nos.: 26-51, 138, 139, 163-183 and 201 or a complement thereof under moderately stringent conditions.

In related aspects, DNA sequences encoding the above polypeptides, expression vectors comprising these DNA sequences and host cells transformed or transfected with such expression vectors are also provided.

25 In another aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, an inventive polypeptide and a known *M. tuberculosis* antigen.

Within other aspects, the present invention provides pharmaceutical compositions that comprise one or more of the above polypeptides, or a DNA molecule encoding such polypeptides, and a physiologically acceptable carrier. The invention

also provides vaccines comprising one or more of the polypeptides as described above and a non-specific immune response enhancer, together with vaccines comprising one or more DNA sequences encoding such polypeptides and a non-specific immune response enhancer.

5 In yet another aspect, methods are provided for inducing protective immunity in a patient, comprising administering to a patient an effective amount of one or more of the above polypeptides.

In further aspects of this invention, methods and diagnostic kits are provided for detecting tuberculosis in a patient. The methods comprise contacting
10 dermal cells of a patient with one or more of the above polypeptides and detecting an immune response on the patient's skin. The diagnostic kits comprise one or more of the above polypeptides in combination with an apparatus sufficient to contact the polypeptide with the dermal cells of a patient.

In yet other aspects, methods are provided for detecting tuberculosis in a
15 patient, such methods comprising contacting dermal cells of a patient with one or more polypeptides encoded by a DNA sequence selected from the group consisting of SEQ ID Nos.: 3, 11, 12, 140, 141, 156-160, 189-193, 199, 200 and 203, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID Nos.: 3, 11, 12, 140, 141, 156-160, 189-193, 199, 200 and 203; and detecting an
20 immune response on the patient's skin. Diagnostic kits for use in such methods are also provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All
25 references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCE IDENTIFIERS

Figure 1A and B illustrate the stimulation of proliferation and interferon-
30 γ production in T cells derived from a first and a second *M. tuberculosis*-immune donor, respectively, by the 14 Kd, 20 Kd and 26 Kd antigens described in Example 1.

Figure 2 illustrates the stimulation of proliferation and interferon- γ production in T cells derived from an *M. tuberculosis*-immune individual by the two representative polypeptides TbRa3 and TbRa9.

Figures 3A-D illustrate the reactivity of antisera raised against secretory
5 *M. tuberculosis* proteins, the known *M. tuberculosis* antigen 85b and the inventive antigens Tb38-1 and TbH-9, respectively, with *M. tuberculosis* lysate (lane 2), *M. tuberculosis* secretory proteins (lane 3), recombinant Tb38-1 (lane 4), recombinant TbH-9 (lane 5) and recombinant 85b (lane 5).

Figure 4A illustrates the stimulation of proliferation in a TbH-9-specific
10 T cell clone by secretory *M. tuberculosis* proteins, recombinant TbH-9 and a control antigen, TbRa11.

Figure 4B illustrates the stimulation of interferon- γ production in a TbH-9-specific T cell clone by secretory *M. tuberculosis* proteins, PPD and recombinant TbH-9.

15 Figures 5A and B illustrate the stimulation of proliferation and interferon- γ production in TbH9-specific T cells by the fusion protein TbH9-Tb38-1.

Figures 6A and B illustrate the stimulation of proliferation and interferon- γ production in Tb38-1-specific T cells by the fusion protein TbH9-Tb38-1.

Figures 7A and B illustrate the stimulation of proliferation and
20 interferon- γ production in T cells previously shown to respond to both TbH-9 and Tb38-1 by the fusion protein TbH9-Tb38-1.

Figures 8A and B illustrate the stimulation of proliferation and interferon- γ production in T cells derived from a first *M. tuberculosis*-immune individual by the representative polypeptides XP-1, RDIF6, RDIF8, RDIF10 and
25 RDIF11.

Figures 9A and B illustrate the stimulation of proliferation and interferon- γ production in T cells derived from a second *M. tuberculosis*-immune individual by the representative polypeptides XP-1, RDIF6, RDIF8, RDIF10 and RDIF11.

SEQ. ID NO. 1 is the DNA sequence of TbRa1.
SEQ. ID NO. 2 is the DNA sequence of TbRa10.
SEQ. ID NO. 3 is the DNA sequence of TbRa11.
SEQ. ID NO. 4 is the DNA sequence of TbRa12.
5 SEQ. ID NO. 5 is the DNA sequence of TbRa13.
SEQ. ID NO. 6 is the DNA sequence of TbRa16.
SEQ. ID NO. 7 is the DNA sequence of TbRa17.
SEQ. ID NO. 8 is the DNA sequence of TbRa18.
SEQ. ID NO. 9 is the DNA sequence of TbRa19.
10 SEQ. ID NO. 10 is the DNA sequence of TbRa24.
SEQ. ID NO. 11 is the DNA sequence of TbRa26.
SEQ. ID NO. 12 is the DNA sequence of TbRa28.
SEQ. ID NO. 13 is the DNA sequence of TbRa29.
SEQ. ID NO. 14 is the DNA sequence of TbRa2A.
15 SEQ. ID NO. 15 is the DNA sequence of TbRa3.
SEQ. ID NO. 16 is the DNA sequence of TbRa32.
SEQ. ID NO. 17 is the DNA sequence of TbRa35.
SEQ. ID NO. 18 is the DNA sequence of TbRa36.
SEQ. ID NO. 19 is the DNA sequence of TbRa4.
20 SEQ. ID NO. 20 is the DNA sequence of TbRa9.
SEQ. ID NO. 21 is the DNA sequence of TbRaB.
SEQ. ID NO. 22 is the DNA sequence of TbRaC.
SEQ. ID NO. 23 is the DNA sequence of TbRaD.
SEQ. ID NO. 24 is the DNA sequence of YYWCPG.
25 SEQ. ID NO. 25 is the DNA sequence of AAMK.
SEQ. ID NO. 26 is the DNA sequence of TbL-23.
SEQ. ID NO. 27 is the DNA sequence of TbL-24.
SEQ. ID NO. 28 is the DNA sequence of TbL-25.
SEQ. ID NO. 29 is the DNA sequence of TbL-28.
30 SEQ. ID NO. 30 is the DNA sequence of TbL-29.

- SEQ. ID NO. 31 is the DNA sequence of TbH-5.
- SEQ. ID NO. 32 is the DNA sequence of TbH-8.
- SEQ. ID NO. 33 is the DNA sequence of TbH-9.
- SEQ. ID NO. 34 is the DNA sequence of TbM-1.
- 5 SEQ. ID NO. 35 is the DNA sequence of TbM-3.
- SEQ. ID NO. 36 is the DNA sequence of TbM-6.
- SEQ. ID NO. 37 is the DNA sequence of TbM-7.
- SEQ. ID NO. 38 is the DNA sequence of TbM-9.
- SEQ. ID NO. 39 is the DNA sequence of TbM-12.
- 10 SEQ. ID NO. 40 is the DNA sequence of TbM-13.
- SEQ. ID NO. 41 is the DNA sequence of TbM-14.
- SEQ. ID NO. 42 is the DNA sequence of TbM-15.
- SEQ. ID NO. 43 is the DNA sequence of TbH-4.
- SEQ. ID NO. 44 is the DNA sequence of TbH-4-FWD.
- 15 SEQ. ID NO. 45 is the DNA sequence of TbH-12.
- SEQ. ID NO. 46 is the DNA sequence of Tb38-1.
- SEQ. ID NO. 47 is the DNA sequence of Tb38-4.
- SEQ. ID NO. 48 is the DNA sequence of TbL-17.
- SEQ. ID NO. 49 is the DNA sequence of TbL-20.
- 20 SEQ. ID NO. 50 is the DNA sequence of TbL-21.
- SEQ. ID NO. 51 is the DNA sequence of TbH-16.
- SEQ. ID NO. 52 is the DNA sequence of DPEP.
- SEQ. ID NO. 53 is the deduced amino acid sequence of DPEP.
- SEQ. ID NO. 54 is the protein sequence of DPV N-terminal Antigen.
- 25 SEQ. ID NO. 55 is the protein sequence of AVGS N-terminal Antigen.
- SEQ. ID NO. 56 is the protein sequence of AAMK N-terminal Antigen.
- SEQ. ID NO. 57 is the protein sequence of YYWC N-terminal Antigen.
- SEQ. ID NO. 58 is the protein sequence of DIGS N-terminal Antigen.
- SEQ. ID NO. 59 is the protein sequence of AEES N-terminal Antigen.
- 30 SEQ. ID NO. 60 is the protein sequence of DPEP N-terminal Antigen.

- SEQ. ID NO. 61 is the protein sequence of APKT N-terminal Antigen.
SEQ. ID NO. 62 is the protein sequence of DPAS N-terminal Antigen.
SEQ. ID NO. 63 is the deduced amino acid sequence of TbRa1.
SEQ. ID NO. 64 is the deduced amino acid sequence of TbRa10.
5 SEQ. ID NO. 65 is the deduced amino acid sequence of TbRa11.
SEQ. ID NO. 66 is the deduced amino acid sequence of TbRa12.
SEQ. ID NO. 67 is the deduced amino acid sequence of TbRa13.
SEQ. ID NO. 68 is the deduced amino acid sequence of TbRa16.
SEQ. ID NO. 69 is the deduced amino acid sequence of TbRa17.
10 SEQ. ID NO. 70 is the deduced amino acid sequence of TbRa18.
SEQ. ID NO. 71 is the deduced amino acid sequence of TbRa19.
SEQ. ID NO. 72 is the deduced amino acid sequence of TbRa24.
SEQ. ID NO. 73 is the deduced amino acid sequence of TbRa26.
SEQ. ID NO. 74 is the deduced amino acid sequence of TbRa28.
15 SEQ. ID NO. 75 is the deduced amino acid sequence of TbRa29.
SEQ. ID NO. 76 is the deduced amino acid sequence of TbRa2A.
SEQ. ID NO. 77 is the deduced amino acid sequence of TbRa3.
SEQ. ID NO. 78 is the deduced amino acid sequence of TbRa32.
SEQ. ID NO. 79 is the deduced amino acid sequence of TbRa35.
20 SEQ. ID NO. 80 is the deduced amino acid sequence of TbRa36.
SEQ. ID NO. 81 is the deduced amino acid sequence of TbRa4.
SEQ. ID NO. 82 is the deduced amino acid sequence of TbRa9.
SEQ. ID NO. 83 is the deduced amino acid sequence of TbRaB.
SEQ. ID NO. 84 is the deduced amino acid sequence of TbRaC.
25 SEQ. ID NO. 85 is the deduced amino acid sequence of TbRaD.
SEQ. ID NO. 86 is the deduced amino acid sequence of YYWCPG.
SEQ. ID NO. 87 is the deduced amino acid sequence of TbAAMK.
SEQ. ID NO. 88 is the deduced amino acid sequence of Tb38-1.
SEQ. ID NO. 89 is the deduced amino acid sequence of TbH-4.
30 SEQ. ID NO. 90 is the deduced amino acid sequence of TbH-8.

- SEQ. ID NO. 91 is the deduced amino acid sequence of TbH-9.
SEQ. ID NO. 92 is the deduced amino acid sequence of TbH-12.
SEQ. ID NO. 93 is the amino acid sequence of Tb38-1 Peptide 1.
SEQ. ID NO. 94 is the amino acid sequence of Tb38-1 Peptide 2.
5 SEQ. ID NO. 95 is the amino acid sequence of Tb38-1 Peptide 3.
SEQ. ID NO. 96 is the amino acid sequence of Tb38-1 Peptide 4.
SEQ. ID NO. 97 is the amino acid sequence of Tb38-1 Peptide 5.
SEQ. ID NO. 98 is the amino acid sequence of Tb38-1 Peptide 6.
SEQ. ID NO. 99 is the DNA sequence of DPAS.
10 SEQ. ID NO. 100 is the deduced amino acid sequence of DPAS.
SEQ. ID NO. 101 is the DNA sequence of DPV.
SEQ. ID NO. 102 is the deduced amino acid sequence of DPV.
SEQ. ID NO. 103 is the DNA sequence of ESAT-6.
SEQ. ID NO. 104 is the deduced amino acid sequence of ESAT-6.
15 SEQ. ID NO. 105 is the DNA sequence of TbH-8-2.
SEQ. ID NO. 106 is the DNA sequence of TbH-9FL.
SEQ. ID NO. 107 is the deduced amino acid sequence of TbH-9FL.
SEQ. ID NO. 108 is the DNA sequence of TbH-9-1.
SEQ. ID NO. 109 is the deduced amino acid sequence of TbH-9-1.
20 SEQ. ID NO. 110 is the DNA sequence of TbH-9-4.
SEQ. ID NO. 111 is the deduced amino acid sequence of TbH-9-4.
SEQ. ID NO. 112 is the DNA sequence of Tb38-1F2 IN.
SEQ. ID NO. 113 is the DNA sequence of Tb38-2F2 RP.
SEQ. ID NO. 114 is the deduced amino acid sequence of Tb37-FL.
25 SEQ. ID NO. 115 is the deduced amino acid sequence of Tb38-IN.
SEQ. ID NO. 116 is the DNA sequence of Tb38-1F3.
SEQ. ID NO. 117 is the deduced amino acid sequence of Tb38-1F3.
SEQ. ID NO. 118 is the DNA sequence of Tb38-1F5.
SEQ. ID NO. 119 is the DNA sequence of Tb38-1F6.
30 SEQ. ID NO. 120 is the deduced N-terminal amino acid sequence of DPV.

- SEQ. ID NO. 121 is the deduced N-terminal amino acid sequence of AVGS.
SEQ. ID NO. 122 is the deduced N-terminal amino acid sequence of AAMK.
SEQ. ID NO. 123 is the deduced N-terminal amino acid sequence of YYWC.
SEQ. ID NO. 124 is the deduced N-terminal amino acid sequence of DIGS.
5 SEQ. ID NO. 125 is the deduced N-terminal amino acid sequence of AEES.
SEQ. ID NO. 126 is the deduced N-terminal amino acid sequence of DPEP.
SEQ. ID NO. 127 is the deduced N-terminal amino acid sequence of APKT.
SEQ. ID NO. 128 is the deduced amino acid sequence of DPAS.
SEQ. ID NO. 129 is the protein sequence of DPPD N-terminal Antigen.
10 SEQ ID NO. 130-133 are the protein sequences of four DPPD cyanogen
bromide fragments.
SEQ ID NO. 134 is the N-terminal protein sequence of XDS antigen.
SEQ ID NO. 135 is the N-terminal protein sequence of AGD antigen.
SEQ ID NO. 136 is the N-terminal protein sequence of APE antigen.
15 SEQ ID NO. 137 is the N-terminal protein sequence of XYI antigen.
SEQ ID NO. 138 is the DNA sequence of TbH-29.
SEQ ID NO. 139 is the DNA sequence of TbH-30.
SEQ ID NO. 140 is the DNA sequence of TbH-32.
SEQ ID NO. 141 is the DNA sequence of TbH-33.
20 SEQ ID NO. 142 is the predicted amino acid sequence of TbH-29.
SEQ ID NO. 143 is the predicted amino acid sequence of TbH-30.
SEQ ID NO. 144 is the predicted amino acid sequence of TbH-32.
SEQ ID NO. 145 is the predicted amino acid sequence of TbH-33.
SEQ ID NO: 146-151 are PCR primers used in the preparation of a fusion
25 protein containing TbRa3, 38 kD and Tb38-1.
SEQ ID NO: 152 is the DNA sequence of the fusion protein containing TbRa3,
38 kD and Tb38-1.
SEQ ID NO: 153 is the amino acid sequence of the fusion protein containing
TbRa3, 38 kD and Tb38-1.
30 SEQ ID NO: 154 is the DNA sequence of the *M. tuberculosis* antigen 38 kD.

SEQ ID NO: 155 is the amino acid sequence of the *M. tuberculosis* antigen 38 kD.

SEQ ID NO: 156 is the DNA sequence of XP14.

SEQ ID NO: 157 is the DNA sequence of XP24.

5 SEQ ID NO: 158 is the DNA sequence of XP31.

SEQ ID NO: 159 is the 5' DNA sequence of XP32.

SEQ ID NO: 160 is the 3' DNA sequence of XP32.

SEQ ID NO: 161 is the predicted amino acid sequence of XP14.

10 SEQ ID NO: 162 is the predicted amino acid sequence encoded by the reverse complement of XP14.

SEQ ID NO: 163 is the DNA sequence of XP27.

SEQ ID NO: 164 is the DNA sequence of XP36.

SEQ ID NO: 165 is the 5' DNA sequence of XP4.

SEQ ID NO: 166 is the 5' DNA sequence of XP5.

15 SEQ ID NO: 167 is the 5' DNA sequence of XP17.

SEQ ID NO: 168 is the 5' DNA sequence of XP30.

SEQ ID NO: 169 is the 5' DNA sequence of XP2.

SEQ ID NO: 170 is the 3' DNA sequence of XP2.

SEQ ID NO: 171 is the 5' DNA sequence of XP3.

20 SEQ ID NO: 172 is the 3' DNA sequence of XP3.

SEQ ID NO: 173 is the 5' DNA sequence of XP6.

SEQ ID NO: 174 is the 3' DNA sequence of XP6.

SEQ ID NO: 175 is the 5' DNA sequence of XP18.

SEQ ID NO: 176 is the 3' DNA sequence of XP18.

25 SEQ ID NO: 177 is the 5' DNA sequence of XP19.

SEQ ID NO: 178 is the 3' DNA sequence of XP19.

SEQ ID NO: 179 is the 5' DNA sequence of XP22.

SEQ ID NO: 180 is the 3' DNA sequence of XP22.

SEQ ID NO: 181 is the 5' DNA sequence of XP25.

30 SEQ ID NO: 182 is the 3' DNA sequence of XP25.

SEQ ID NO: 183 is the full-length DNA sequence of TbH4-XP1.

SEQ ID NO: 184 is the predicted amino acid sequence of TbH4-XP1.

SEQ ID NO: 185 is the predicted amino acid sequence encoded by the reverse complement of TbH4-XP1.

5 SEQ ID NO: 186 is a first predicted amino acid sequence encoded by XP36.

SEQ ID NO: 187 is a second predicted amino acid sequence encoded by XP36.

SEQ ID NO: 188 is the predicted amino acid sequence encoded by the reverse complement of XP36.

SEQ ID NO: 189 is the DNA sequence of RDIF2.

10 SEQ ID NO: 190 is the DNA sequence of RDIF5.

SEQ ID NO: 191 is the DNA sequence of RDIF8.

SEQ ID NO: 192 is the DNA sequence of RDIF10.

SEQ ID NO: 193 is the DNA sequence of RDIF11.

SEQ ID NO: 194 is the predicted amino acid sequence of RDIF2.

15 SEQ ID NO: 195 is the predicted amino acid sequence of RDIF5.

SEQ ID NO: 196 is the predicted amino acid sequence of RDIF8.

SEQ ID NO: 197 is the predicted amino acid sequence of RDIF10.

SEQ ID NO: 198 is the predicted amino acid sequence of RDIF11.

SEQ ID NO: 199 is the 5' DNA sequence of RDIF12.

20 SEQ ID NO: 200 is the 3' DNA sequence of RDIF12.

SEQ ID NO: 201 is the DNA sequence of RDIF7.

SEQ ID NO: 202 is the predicted amino acid sequence of RDIF7.

SEQ ID NO: 203 is the DNA sequence of DIF2-1.

SEQ ID NO: 204 is the predicted amino acid sequence of DIF2-1.

25 SEQ ID NO: 205-212 are PCR primers used in the preparation of a fusion protein containing TbRa3, 38 kD, Tb38-1 and DPEP (hereinafter referred to as TbF-2).

SEQ ID NO: 213 is the DNA sequence of the fusion protein TbF-2.

SEQ ID NO: 214 is the amino acid sequence of the fusion protein TbF-2.

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for preventing, treating and diagnosing tuberculosis. The compositions of the subject invention include polypeptides that comprise at least one immunogenic portion of a *M. tuberculosis* antigen, or a variant of such an antigen that differs only in conservative substitutions and/or modifications. Polypeptides within the scope of the present invention include, but are not limited to, immunogenic soluble *M. tuberculosis* antigens. A "soluble *M. tuberculosis* antigen" is a protein of *M. tuberculosis* origin that is present in *M. tuberculosis* culture filtrate. As used herein, the term "polypeptide" encompasses amino acid chains of any length, including full length proteins (*i.e.*, antigens), wherein the amino acid residues are linked by covalent peptide bonds. Thus, a polypeptide comprising an immunogenic portion of one of the above antigens may consist entirely of the immunogenic portion, or may contain additional sequences. The additional sequences may be derived from the native *M. tuberculosis* antigen or may be heterologous, and such sequences may (but need not) be immunogenic.

"Immunogenic," as used herein, refers to the ability to elicit an immune response (*e.g.*, cellular) in a patient, such as a human, and/or in a biological sample. In particular, antigens that are immunogenic (and immunogenic portions or other variants of such antigens) are capable of stimulating cell proliferation, interleukin-12 production and/or interferon- γ production in biological samples comprising one or more cells selected from the group of T cells, NK cells, B cells and macrophages, where the cells are derived from an *M. tuberculosis*-immune individual. Polypeptides comprising at least an immunogenic portion of one or more *M. tuberculosis* antigens may generally be used to detect tuberculosis or to induce protective immunity against tuberculosis in a patient.

The compositions and methods of this invention also encompass variants of the above polypeptides. A "variant," as used herein, is a polypeptide that differs from the native antigen only in conservative substitutions and/or modifications, such that the ability of the polypeptide to induce an immune response is retained. Such variants may generally be identified by modifying one of the above polypeptide

sequences, and evaluating the immunogenic properties of the modified polypeptide using, for example, the representative procedures described herein.

A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his.

10 Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenic properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

20 In a related aspect, combination polypeptides are disclosed. A "combination polypeptide" is a polypeptide comprising at least one of the above immunogenic portions and one or more additional immunogenic *M. tuberculosis* sequences, which are joined via a peptide linkage into a single amino acid chain. The sequences may be joined directly (i.e., with no intervening amino acids) or may be joined by way of a linker sequence (e.g., Gly-Cys-Gly) that does not significantly diminish the immunogenic properties of the component polypeptides.

25 In general, *M. tuberculosis* antigens, and DNA sequences encoding such antigens, may be prepared using any of a variety of procedures. For example, soluble antigens may be isolated from *M. tuberculosis* culture filtrate by procedures known to those of ordinary skill in the art, including anion-exchange and reverse phase chromatography. Purified antigens are then evaluated for their ability to elicit an

30

appropriate immune response (*e.g.*, cellular) using, for example, the representative methods described herein. Immunogenic antigens may then be partially sequenced using techniques such as traditional Edman chemistry. *See* Edman and Berg, *Eur. J. Biochem.* 80:116-132, 1967.

5 Immunogenic antigens may also be produced recombinantly using a DNA sequence that encodes the antigen, which has been inserted into an expression vector and expressed in an appropriate host. DNA molecules encoding soluble antigens may be isolated by screening an appropriate *M. tuberculosis* expression library with anti-sera (*e.g.*, rabbit) raised specifically against soluble *M. tuberculosis* antigens. DNA
10 sequences encoding antigens that may or may not be soluble may be identified by screening an appropriate *M. tuberculosis* genomic or cDNA expression library with sera obtained from patients infected with *M. tuberculosis*. Such screens may generally be performed using techniques well known to those of ordinary skill in the art, such as those described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold
15 Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989.

DNA sequences encoding soluble antigens may also be obtained by screening an appropriate *M. tuberculosis* cDNA or genomic DNA library for DNA sequences that hybridize to degenerate oligonucleotides derived from partial amino acid sequences of isolated soluble antigens. Degenerate oligonucleotide sequences for use in
20 such a screen may be designed and synthesized, and the screen may be performed, as described (for example) in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989 (and references cited therein). Polymerase chain reaction (PCR) may also be employed, using the above oligonucleotides in methods well known in the art, to isolate a nucleic acid probe from a
25 cDNA or genomic library. The library screen may then be performed using the isolated probe.

Alternatively, genomic or cDNA libraries derived from *M. tuberculosis* may be screened directly using peripheral blood mononuclear cells (PBMCs) or T cell lines or clones derived from one or more *M. tuberculosis*-immune individuals. In
30 general, PBMCs and/or T cells for use in such screens may be prepared as described

below. Direct library screens may generally be performed by assaying pools of expressed recombinant proteins for the ability to induce proliferation and/or interferon- γ production in T cells derived from an *M. tuberculosis*-immune individual. Alternatively, potential T cell antigens may be first selected based on antibody
5 reactivity, as described above.

Regardless of the method of preparation, the antigens (and immunogenic portions thereof) described herein (which may or may not be soluble) have the ability to induce an immunogenic response. More specifically, the antigens have the ability to induce proliferation and/or cytokine production (*i.e.*, interferon- γ and/or interleukin-12
10 production) in T cells, NK cells, B cells and/or macrophages derived from an *M. tuberculosis*-immune individual. The selection of cell type for use in evaluating an immunogenic response to a antigen will, of course, depend on the desired response. For example, interleukin-12 production is most readily evaluated using preparations containing B cells and/or macrophages. An *M. tuberculosis*-immune individual is one
15 who is considered to be resistant to the development of tuberculosis by virtue of having mounted an effective T cell response to *M. tuberculosis* (*i.e.*, substantially free of disease symptoms). Such individuals may be identified based on a strongly positive (*i.e.*, greater than about 10 mm diameter induration) intradermal skin test response to tuberculosis proteins (PPD) and an absence of any signs or symptoms of tuberculosis
20 disease. T cells, NK cells, B cells and macrophages derived from *M. tuberculosis*-immune individuals may be prepared using methods known to those of ordinary skill in the art. For example, a preparation of PBMCs (*i.e.*, peripheral blood mononuclear cells) may be employed without further separation of component cells. PBMCs may generally be prepared, for example, using density centrifugation through Ficoll™
25 (Winthrop Laboratories, NY). T cells for use in the assays described herein may also be purified directly from PBMCs. Alternatively, an enriched T cell line reactive against mycobacterial proteins, or T cell clones reactive to individual mycobacterial proteins, may be employed. Such T cell clones may be generated by, for example, culturing PBMCs from *M. tuberculosis*-immune individuals with mycobacterial proteins for a
30 period of 2-4 weeks. This allows expansion of only the mycobacterial protein-specific

T cells, resulting in a line composed solely of such cells. These cells may then be cloned and tested with individual proteins, using methods known to those of ordinary skill in the art, to more accurately define individual T cell specificity. In general, antigens that test positive in assays for proliferation and/or cytokine production (*i.e.*,
5 interferon- γ and/or interleukin-12 production) performed using T cells, NK cells, B cells and/or macrophages derived from an *M. tuberculosis*-immune individual are considered immunogenic. Such assays may be performed, for example, using the representative procedures described below. Immunogenic portions of such antigens may be identified using similar assays, and may be present within the polypeptides described herein.

10 The ability of a polypeptide (*e.g.*, an immunogenic antigen, or a portion or other variant thereof) to induce cell proliferation is evaluated by contacting the cells (*e.g.*, T cells and/or NK cells) with the polypeptide and measuring the proliferation of the cells. In general, the amount of polypeptide that is sufficient for evaluation of about 10^5 cells ranges from about 10 ng/mL to about 100 μ g/mL and preferably is about
15 10 μ g/mL. The incubation of polypeptide with cells is typically performed at 37°C for about six days. Following incubation with polypeptide, the cells are assayed for a proliferative response, which may be evaluated by methods known to those of ordinary skill in the art, such as exposing cells to a pulse of radiolabeled thymidine and measuring the incorporation of label into cellular DNA. In general, a polypeptide that
20 results in at least a three fold increase in proliferation above background (*i.e.*, the proliferation observed for cells cultured without polypeptide) is considered to be able to induce proliferation.

The ability of a polypeptide to stimulate the production of interferon- γ and/or interleukin-12 in cells may be evaluated by contacting the cells with the
25 polypeptide and measuring the level of interferon- γ or interleukin-12 produced by the cells. In general, the amount of polypeptide that is sufficient for the evaluation of about 10^5 cells ranges from about 10 ng/mL to about 100 μ g/mL and preferably is about 10 μ g/mL. The polypeptide may, but need not, be immobilized on a solid support, such as a bead or a biodegradable microsphere, such as those described in U.S. Patent
30 Nos. 4,897,268 and 5,075,109. The incubation of polypeptide with the cells is typically

performed at 37°C for about six days. Following incubation with polypeptide, the cells are assayed for interferon- γ and/or interleukin-12 (or one or more subunits thereof), which may be evaluated by methods known to those of ordinary skill in the art, such as an enzyme-linked immunosorbent assay (ELISA) or, in the case of IL-12 P70 subunit, a
5 bioassay such as an assay measuring proliferation of T cells. In general, a polypeptide that results in the production of at least 50 pg of interferon- γ per mL of cultured supernatant (containing 10^4 - 10^5 T cells per mL) is considered able to stimulate the production of interferon- γ . A polypeptide that stimulates the production of at least 10 pg/mL of IL-12 P70 subunit, and/or at least 100 pg/mL of IL-12 P40 subunit, per 10^5
10 macrophages or B cells (or per 3×10^5 PBMC) is considered able to stimulate the production of IL-12.

In general, immunogenic antigens are those antigens that stimulate proliferation and/or cytokine production (*i.e.*, interferon- γ and/or interleukin-12 production) in T cells, NK cells, B cells and/or macrophages derived from at least about
15 25% of *M. tuberculosis*-immune individuals. Among these immunogenic antigens, polypeptides having superior therapeutic properties may be distinguished based on the magnitude of the responses in the above assays and based on the percentage of individuals for which a response is observed. In addition, antigens having superior therapeutic properties will not stimulate proliferation and/or cytokine production *in*
20 *vitro* in cells derived from more than about 25% of individuals that are not *M. tuberculosis*-immune, thereby eliminating responses that are not specifically due to *M. tuberculosis*-responsive cells. Those antigens that induce a response in a high percentage of T cell, NK cell, B cell and/or macrophage preparations from *M. tuberculosis*-immune individuals (with a low incidence of responses in cell
25 preparations from other individuals) have superior therapeutic properties.

Antigens with superior therapeutic properties may also be identified based on their ability to diminish the severity of *M. tuberculosis* infection in experimental animals, when administered as a vaccine. Suitable vaccine preparations for use on experimental animals are described in detail below. Efficacy may be
30 determined based on the ability of the antigen to provide at least about a 50% reduction

in bacterial numbers and/or at least about a 40% decrease in mortality following experimental infection. Suitable experimental animals include mice, guinea pigs and primates.

Antigens having superior diagnostic properties may generally be identified based on the ability to elicit a response in an intradermal skin test performed on an individual with active tuberculosis, but not in a test performed on an individual who is not infected with *M. tuberculosis*. Skin tests may generally be performed as described below, with a response of at least 5 mm induration considered positive.

Immunogenic portions of the antigens described herein may be prepared and identified using well known techniques, such as those summarized in Paul, *Fundamental Immunology*, 3d ed., Raven Press, 1993, pp. 243-247 and references cited therein. Such techniques include screening polypeptide portions of the native antigen for immunogenic properties. The representative proliferation and cytokine production assays described herein may generally be employed in these screens. An immunogenic portion of a polypeptide is a portion that, within such representative assays, generates an immune response (e.g., proliferation, interferon- γ production and/or interleukin-12 production) that is substantially similar to that generated by the full length antigen. In other words, an immunogenic portion of an antigen may generate at least about 20%, and preferably about 100%, of the proliferation induced by the full length antigen in the model proliferation assay described herein. An immunogenic portion may also, or alternatively, stimulate the production of at least about 20%, and preferably about 100%, of the interferon- γ and/or interleukin-12 induced by the full length antigen in the model assay described herein.

Portions and other variants of *M. tuberculosis* antigens may be generated by synthetic or recombinant means. Synthetic polypeptides having fewer than about 100 amino acids, and generally fewer than about 50 amino acids, may be generated using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.*

85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Applied BioSystems, Inc., Foster City, CA, and may be operated according to the manufacturer's instructions. Variants of a native antigen may generally be prepared using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis. Sections of the DNA sequence may also be removed using standard techniques to permit preparation of truncated polypeptides.

Recombinant polypeptides containing portions and/or variants of a native antigen may be readily prepared from a DNA sequence encoding the polypeptide using a variety of techniques well known to those of ordinary skill in the art. For example, supernatants from suitable host/vector systems which secrete recombinant protein into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix or an ion exchange resin. Finally, one or more reverse phase HPLC steps can be employed to further purify a recombinant protein.

Any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant polypeptides of this invention. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line such as COS or CHO. The DNA sequences expressed in this manner may encode naturally occurring antigens, portions of naturally occurring antigens, or other variants thereof.

In general, regardless of the method of preparation, the polypeptides disclosed herein are prepared in substantially pure form. Preferably, the polypeptides are at least about 80% pure, more preferably at least about 90% pure and most preferably at least about 99% pure. In certain preferred embodiments, described in

detail below, the substantially pure polypeptides are incorporated into pharmaceutical compositions or vaccines for use in one or more of the methods disclosed herein.

In certain specific embodiments, the subject invention discloses polypeptides comprising at least an immunogenic portion of a soluble *M. tuberculosis* antigen having one of the following N-terminal sequences, or a variant thereof that differs only in conservative substitutions and/or modifications:

- (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Cys-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu; (SEQ ID No. 120)
- (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser; (SEQ ID No. 121)
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg; (SEQ ID No. 122)
- (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro; (SEQ ID No. 123)
- (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val; (SEQ ID No. 124)
- (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro; (SEQ ID No. 125)
- (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Ala-Ala-Ala-Ser-Pro-Pro-Ser; (SEQ ID No. 126)
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly; (SEQ ID No. 127)
- (i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Leu-Thr-Ser-Leu-Leu-Asn-Ser-Leu-Ala-Asp-Pro-Asn-Val-Ser-Phe-Ala-Asn; (SEQ ID No. 128)
- (j) Xaa-Asp-Ser-Glu-Lys-Ser-Ala-Thr-Ile-Lys-Val-Thr-Asp-Ala-Ser; (SEQ ID No. 134)
- (k) Ala-Gly-Asp-Thr-Xaa-Ile-Tyr-Ile-Val-Gly-Asn-Leu-Thr-Ala-Asp; (SEQ ID No. 135) or

- (l) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly; (SEQ ID No. 136)

wherein Xaa may be any amino acid, preferably a cysteine residue. A DNA sequence encoding the antigen identified as (g) above is provided in SEQ ID No. 52, and the polypeptide encoded by SEQ ID No. 52 is provided in SEQ ID No. 53. A DNA sequence encoding the antigen defined as (a) above is provided in SEQ ID No. 101; its deduced amino acid sequence is provided in SEQ ID No. 102. A DNA sequence corresponding to antigen (d) above is provided in SEQ ID No. 24 a DNA sequence corresponding to antigen (c) is provided in SEQ ID No. 25 and a DNA sequence corresponding to antigen (i) is provided in SEQ ID No. 99; its deduced amino acid sequence is provided in SEQ ID No. 100.

In a further specific embodiment, the subject invention discloses polypeptides comprising at least an immunogenic portion of an *M. tuberculosis* antigen having one of the following N-terminal sequences, or a variant thereof that differs only in conservative substitutions and/or modifications:

- (m) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID No 137) or
- (n) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID No. 129)

wherein Xaa may be any amino acid, preferably a cysteine residue.

In other specific embodiments, the subject invention discloses polypeptides comprising at least an immunogenic portion of a soluble *M. tuberculosis* antigen (or a variant of such an antigen) that comprises one or more of the amino acid sequences encoded by (a) the DNA sequences of SEQ ID Nos.: 1, 2, 4-10, 13-25 and 52; (b) the complements of such DNA sequences, or (c) DNA sequences substantially homologous to a sequence in (a) or (b).

In further specific embodiments, the subject invention discloses polypeptides comprising at least an immunogenic portion of a *M. tuberculosis* antigen (or a variant of such an antigen), which may or may not be soluble, that comprises one or more of the amino acid sequences encoded by (a) the DNA sequences of SEQ ID

Nos.: 26-51, 138, 139, 163-183 and 201, (b) the complements of such DNA sequences or (c) DNA sequences substantially homologous to a sequence in (a) or (b).

In the specific embodiments discussed above, the *M. tuberculosis* antigens include variants that are encoded by DNA sequences which are substantially homologous to one or more of DNA sequences specifically recited herein. "Substantial homology," as used herein, refers to DNA sequences that are capable of hybridizing under moderately stringent conditions. Suitable moderately stringent conditions include prewashing in a solution of 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5X SSC, overnight or, in the case of cross-species homology at 45°C, 0.5X SSC; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS). Such hybridizing DNA sequences are also within the scope of this invention, as are nucleotide sequences that, due to code degeneracy, encode an immunogenic polypeptide that is encoded by a hybridizing DNA sequence.

In a related aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, a polypeptide of the present invention and a known *M. tuberculosis* antigen, such as the 38 kD antigen described in Andersen and Hansen, *Infect. Immun.* 57:2481-2488, 1989, (Genbank Accession No. M30046) or ESAT-6 (SEQ ID Nos. 103 and 104), together with variants of such fusion proteins. The fusion proteins of the present invention may also include a linker peptide between the first and second polypeptides.

A DNA sequence encoding a fusion protein of the present invention is constructed using known recombinant DNA techniques to assemble separate DNA sequences encoding the first and second polypeptides into an appropriate expression vector. The 3' end of a DNA sequence encoding the first polypeptide is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide so that the reading frames of the sequences are in phase to permit mRNA translation of the two DNA sequences into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptides by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons require to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

In another aspect, the present invention provides methods for using one or more of the above polypeptides or fusion proteins (or DNA molecules encoding such polypeptides) to induce protective immunity against tuberculosis in a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with a disease, or may be free of detectable disease and/or infection. In other words, protective immunity may be induced to prevent or treat tuberculosis.

In this aspect, the polypeptide, fusion protein or DNA molecule is generally present within a pharmaceutical composition and/or a vaccine. Pharmaceutical compositions may comprise one or more polypeptides, each of which may contain one or more of the above sequences (or variants thereof), and a physiologically acceptable carrier. Vaccines may comprise one or more of the above polypeptides and a non-specific immune response enhancer, such as an adjuvant or a liposome (into which the polypeptide is incorporated). Such pharmaceutical compositions and vaccines may also contain other *M. tuberculosis* antigens, either incorporated into a combination polypeptide or present within a separate polypeptide.

Alternatively, a vaccine may contain DNA encoding one or more polypeptides as described above, such that the polypeptide is generated *in situ*. In such vaccines, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacterial and viral expression systems. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface. In a preferred embodiment, the DNA may be introduced using a viral expression system (*e.g.*, vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

In a related aspect, a DNA vaccine as described above may be administered simultaneously with or sequentially to either a polypeptide of the present invention or a known *M. tuberculosis* antigen, such as the 38 kD antigen described above. For example, administration of DNA encoding a polypeptide of the present

invention, either "naked" or in a delivery system as described above, may be followed by administration of an antigen in order to enhance the protective immune effect of the vaccine.

Routes and frequency of administration, as well as dosage, will vary from individual to individual and may parallel those currently being used in immunization using BCG. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Between 1 and 3 doses may be administered for a 1-36 week period. Preferably, 3 doses are administered, at intervals of 3-4 months, and booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of polypeptide or DNA that, when administered as described above, is capable of raising an immune response in an immunized patient sufficient to protect the patient from *M. tuberculosis* infection for at least 1-2 years. In general, the amount of polypeptide present in a dose (or produced *in situ* by the DNA in a dose) ranges from about 1 pg to about 100 mg per kg of host, typically from about 10 pg to about 1 mg, and preferably from about 100 pg to about 1 μ g. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactic galactide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Any of a variety of adjuvants may be employed in the vaccines of this invention to nonspecifically enhance the immune response. Most adjuvants contain a

substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a nonspecific stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis*. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Freund's
5 Complete Adjuvant (Difco Laboratories) and Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ). Other suitable adjuvants include alum, biodegradable microspheres, monophosphoryl lipid A and quil A.

In another aspect, this invention provides methods for using one or more of the polypeptides described above to diagnose tuberculosis using a skin test. As used
10 herein, a "skin test" is any assay performed directly on a patient in which a delayed-type hypersensitivity (DTH) reaction (such as swelling, reddening or dermatitis) is measured following intradermal injection of one or more polypeptides as described above. Such injection may be achieved using any suitable device sufficient to contact the polypeptide or polypeptides with dermal cells of the patient, such as a tuberculin
15 syringe or 1 mL syringe. Preferably, the reaction is measured at least 48 hours after injection, more preferably 48-72 hours.

The DTH reaction is a cell-mediated immune response, which is greater in patients that have been exposed previously to the test antigen (*i.e.*, the immunogenic portion of the polypeptide employed, or a variant thereof). The response may be
20 measured visually, using a ruler. In general, a response that is greater than about 0.5 cm in diameter, preferably greater than about 1.0 cm in diameter, is a positive response, indicative of tuberculosis infection, which may or may not be manifested as an active disease.

The polypeptides of this invention are preferably formulated, for use in a
25 skin test, as pharmaceutical compositions containing a polypeptide and a physiologically acceptable carrier, as described above. Such compositions typically contain one or more of the above polypeptides in an amount ranging from about 1 μ g to about 100 μ g, preferably from about 10 μ g to about 50 μ g in a volume of 0.1 mL. Preferably, the carrier employed in such pharmaceutical compositions is a saline
30 solution with appropriate preservatives, such as phenol and/or Tween 80™.

In a preferred embodiment, a polypeptide employed in a skin test is of sufficient size such that it remains at the site of injection for the duration of the reaction period. In general, a polypeptide that is at least 9 amino acids in length is sufficient. The polypeptide is also preferably broken down by macrophages within hours of
5 injection to allow presentation to T-cells. Such polypeptides may contain repeats of one or more of the above sequences and/or other immunogenic or nonimmunogenic sequences.

The following Examples are offered by way of illustration and not by
10 way of limitation.

EXAMPLES

EXAMPLE 1

PURIFICATION AND CHARACTERIZATION OF POLYPEPTIDES FROM *M. TUBERCULOSIS* CULTURE FILTRATE

This example illustrates the preparation of *M. tuberculosis* soluble polypeptides from culture filtrate. Unless otherwise noted, all percentages in the
20 following example are weight per volume.

M. tuberculosis (either H37Ra, ATCC No. 25177, or H37Rv, ATCC No. 25618) was cultured in sterile GAS media at 37°C for fourteen days. The media was then vacuum filtered (leaving the bulk of the cells) through a 0.45 μ filter into a sterile 2.5 L bottle. The media was next filtered through a 0.2 μ filter into a sterile 4 L
25 bottle and NaN_3 was added to the culture filtrate to a concentration of 0.04%. The bottles were then placed in a 4°C cold room.

The culture filtrate was concentrated by placing the filtrate in a 12 L reservoir that had been autoclaved and feeding the filtrate into a 400 ml Amicon stir cell which had been rinsed with ethanol and contained a 10,000 kDa MWCO membrane.

The pressure was maintained at 60 psi using nitrogen gas. This procedure reduced the 12 L volume to approximately 50 ml.

The culture filtrate was dialyzed into 0.1% ammonium bicarbonate using a 8,000 kDa MWCO cellulose ester membrane, with two changes of ammonium bicarbonate solution. Protein concentration was then determined by a commercially available BCA assay (Pierce, Rockford, IL).

The dialyzed culture filtrate was then lyophilized, and the polypeptides resuspended in distilled water. The polypeptides were dialyzed against 0.01 mM 1,3 bis[tris(hydroxymethyl)-methylamino]propane, pH 7.5 (Bis-Tris propane buffer), the initial conditions for anion exchange chromatography. Fractionation was performed using gel perfusion chromatography on a POROS 146 II Q/M anion exchange column 4.6 mm x 100 mm (Perseptive BioSystems, Framingham, MA) equilibrated in 0.01 mM Bis-Tris propane buffer pH 7.5. Polypeptides were eluted with a linear 0-0.5 M NaCl gradient in the above buffer system. The column eluent was monitored at a wavelength of 220 nm.

The pools of polypeptides eluting from the ion exchange column were dialyzed against distilled water and lyophilized. The resulting material was dissolved in 0.1% trifluoroacetic acid (TFA) pH 1.9 in water, and the polypeptides were purified on a Delta-Pak C18 column (Waters, Milford, MA) 300 Angstrom pore size, 5 micron particle size (3.9 x 150 mm). The polypeptides were eluted from the column with a linear gradient from 0-60% dilution buffer (0.1% TFA in acetonitrile). The flow rate was 0.75 ml/minute and the HPLC eluent was monitored at 214 nm. Fractions containing the eluted polypeptides were collected to maximize the purity of the individual samples. Approximately 200 purified polypeptides were obtained.

The purified polypeptides were then screened for the ability to induce T-cell proliferation in PBMC preparations. The PBMCs from donors known to be PPD skin test positive and whose T-cells were shown to proliferate in response to PPD and crude soluble proteins from MTB were cultured in medium comprising RPMI 1640 supplemented with 10% pooled human serum and 50 µg/ml gentamicin. Purified polypeptides were added in duplicate at concentrations of 0.5 to 10 µg/mL. After six

days of culture in 96-well round-bottom plates in a volume of 200 μ l, 50 μ l of medium was removed from each well for determination of IFN- γ levels, as described below. The plates were then pulsed with 1 μ Ci/well of tritiated thymidine for a further 18 hours, harvested and tritium uptake determined using a gas scintillation counter.

- 5 Fractions that resulted in proliferation in both replicates three fold greater than the proliferation observed in cells cultured in medium alone were considered positive.

- IFN- γ was measured using an enzyme-linked immunosorbent assay (ELISA). ELISA plates were coated with a mouse monoclonal antibody directed to human IFN- γ (PharMingen, San Diego, CA) in PBS for four hours at room temperature.
- 10 Wells were then blocked with PBS containing 5% (W/V) non-fat dried milk for 1 hour at room temperature. The plates were then washed six times in PBS/0.2% TWEEN-20 and samples diluted 1:2 in culture medium in the ELISA plates were incubated overnight at room temperature. The plates were again washed and a polyclonal rabbit anti-human IFN- γ serum diluted 1:3000 in PBS/10% normal goat serum was added to
- 15 each well. The plates were then incubated for two hours at room temperature, washed and horseradish peroxidase-coupled anti-rabbit IgG (Sigma Chemical So., St. Louis, MO) was added at a 1:2000 dilution in PBS/5% non-fat dried milk. After a further two hour incubation at room temperature, the plates were washed and TMB substrate added. The reaction was stopped after 20 min with 1 N sulfuric acid. Optical density was
- 20 determined at 450 nm using 570 nm as a reference wavelength. Fractions that resulted in both replicates giving an OD two fold greater than the mean OD from cells cultured in medium alone, plus 3 standard deviations, were considered positive.

- For sequencing, the polypeptides were individually dried onto BiobreneTM (Perkin Elmer/Applied BioSystems Division, Foster City, CA)-treated-glass
- 25 fiber filters. The filters with polypeptide were loaded onto a Perkin Elmer/Applied BioSystems Division Procise 492 protein sequencer. The polypeptides were sequenced from the amino terminal and using traditional Edman chemistry. The amino acid sequence was determined for each polypeptide by comparing the retention time of the PTH amino acid derivative to the appropriate PTH derivative standards.

Using the procedure described above, antigens having the following N-terminal sequences were isolated:

- (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Xaa-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu; (SEQ ID No. 54)
- 5 (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser; (SEQ ID No. 55)
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg; (SEQ ID No. 56)
- (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro; (SEQ ID No. 57)
- 10 (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val; (SEQ ID No. 58)
- (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro; (SEQ ID No. 59)
- 15 (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Ala-Ala-Ala-Ala-Pro-Pro-Ala; (SEQ ID No. 60) and
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly; (SEQ ID No. 61)

wherein Xaa may be any amino acid.

20 An additional antigen was isolated employing a microbore HPLC purification step in addition to the procedure described above. Specifically, 20 μ l of a fraction comprising a mixture of antigens from the chromatographic purification step previously described, was purified on an Aquapore C18 column (Perkin Elmer/Applied Biosystems Division, Foster City, CA) with a 7 micron pore size, column size 1 mm x
25 100 mm, in a Perkin Elmer/Applied Biosystems Division Model 172 HPLC. Fractions were eluted from the column with a linear gradient of 1%/minute of acetonitrile (containing 0.05% TFA) in water (0.05% TFA) at a flow rate of 80 μ l/minute. The eluent was monitored at 250 nm. The original fraction was separated into 4 major peaks plus other smaller components and a polypeptide was obtained which was shown to

have a molecular weight of 12.054 Kd (by mass spectrometry) and the following N-terminal sequence:

- (i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Gln-
Thr-Ser-Leu-Leu-Asn-Asn-Leu-Ala-Asp-Pro-Asp-Val-Ser-Phe-
Ala-Asp (SEQ ID No. 62).

This polypeptide was shown to induce proliferation and IFN- γ production in PBMC preparations using the assays described above.

Additional soluble antigens were isolated from *M. tuberculosis* culture filtrate as follows. *M. tuberculosis* culture filtrate was prepared as described above. Following dialysis against Bis-Tris propane buffer, at pH 5.5, fractionation was performed using anion exchange chromatography on a Poros QE column 4.6 x 100 mm (Perseptive Biosystems) equilibrated in Bis-Tris propane buffer pH 5.5. Polypeptides were eluted with a linear 0-1.5 M NaCl gradient in the above buffer system at a flow rate of 10 ml/min. The column eluent was monitored at a wavelength of 214 nm.

The fractions eluting from the ion exchange column were pooled and subjected to reverse phase chromatography using a Poros R2 column 4.6 x 100 mm (Perseptive Biosystems). Polypeptides were eluted from the column with a linear gradient from 0-100% acetonitrile (0.1% TFA) at a flow rate of 5 ml/min. The eluent was monitored at 214 nm.

Fractions containing the eluted polypeptides were lyophilized and resuspended in 80 μ l of aqueous 0.1% TFA and further subjected to reverse phase chromatography on a Vydac C4 column 4.6 x 150 mm (Western Analytical, Temecula, CA) with a linear gradient of 0-100% acetonitrile (0.1% TFA) at a flow rate of 2 ml/min. Eluent was monitored at 214 nm.

The fraction with biological activity was separated into one major peak plus other smaller components. Western blot of this peak onto PVDF membrane revealed three major bands of molecular weights 14 Kd, 20 Kd and 26 Kd. These polypeptides were determined to have the following N-terminal sequences, respectively:

- (j) Xaa-Asp-Ser-Glu-Lys-Ser-Ala-Thr-Ile-Lys-Val-Thr-Asp-Ala-Ser; (SEQ ID No. 134)

(k) Ala-Gly-Asp-Thr-Xaa-Ile-Tyr-Ile-Val-Gly-Asn-Leu-Thr-Ala-Asp; (SEQ ID No. 135) and

(l) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly; (SEQ ID No. 136), wherein Xaa may be any amino acid.

5 Using the assays described above, these polypeptides were shown to induce proliferation and IFN- γ production in PBMC preparations. Figs. 1A and B show the results of such assays using PBMC preparations from a first and a second donor, respectively.

DNA sequences that encode the antigens designated as (a), (c), (d) and
10 (g) above were obtained by screening a genomic *M. tuberculosis* library using ³²P end labeled degenerate oligonucleotides corresponding to the N-terminal sequence and containing *M. tuberculosis* codon bias. The screen performed using a probe corresponding to antigen (a) above identified a clone having the sequence provided in SEQ ID No. 101. The polypeptide encoded by SEQ ID No. 101 is provided in SEQ ID
15 No. 102. The screen performed using a probe corresponding to antigen (g) above identified a clone having the sequence provided in SEQ ID No. 52. The polypeptide encoded by SEQ ID No. 52 is provided in SEQ ID No. 53. The screen performed using a probe corresponding to antigen (d) above identified a clone having the sequence provided in SEQ ID No. 24, and the screen performed with a probe corresponding to
20 antigen (c) identified a clone having the sequence provided in SEQ ID No. 25.

The above amino acid sequences were compared to known amino acid sequences in the gene bank using the DNA STAR system. The database searched contains some 173,000 proteins and is a combination of the Swiss, PIR databases along with translated protein sequences (Version 87). No significant homologies to the amino
25 acid sequences for antigens (a)-(h) and (l) were detected.

The amino acid sequence for antigen (i) was found to be homologous to a sequence from *M. leprae*. The full length *M. leprae* sequence was amplified from genomic DNA using the sequence obtained from GENBANK. This sequence was then used to screen the *M. tuberculosis* library described below in Example 2 and a full
30 length copy of the *M. tuberculosis* homologue was obtained (SEQ ID No. 99).

The amino acid sequence for antigen (j) was found to be homologous to a known *M. tuberculosis* protein translated from a DNA sequence. To the best of the inventors' knowledge, this protein has not been previously shown to possess T-cell stimulatory activity. The amino acid sequence for antigen (k) was found to be related to a sequence from *M. leprae*.

In the proliferation and IFN- γ assays described above, using three PPD positive donors, the results for representative antigens provided above are presented in Table 1:

10

TABLE 1RESULTS OF PBMC PROLIFERATION AND IFN- γ ASSAYS

Sequence	Proliferation	IFN- γ
(a)	+	-
(c)	+++	+++
(d)	++	++
(g)	+++	+++
(h)	+++	+++

In Table 1, responses that gave a stimulation index (SI) of between 2 and 4 (compared to cells cultured in medium alone) were scored as +, an SI of 4-8 or 2-4 at a concentration of 1 μ g or less was scored as ++ and an SI of greater than 8 was scored as +++. The antigen of sequence (i) was found to have a high SI (+++) for one donor and lower SI (++ and +) for the two other donors in both proliferation and IFN- γ assays. These results indicate that these antigens are capable of inducing proliferation and/or interferon- γ production.

20

EXAMPLE 2

USE OF PATIENT SERA TO ISOLATE *M. TUBERCULOSIS* ANTIGENS

This example illustrates the isolation of antigens from *M. tuberculosis* lysate by screening with serum from *M. tuberculosis*-infected individuals.

Dessicated *M. tuberculosis* H37Ra (Difco Laboratories) was added to a 2% NP40 solution, and alternately homogenized and sonicated three times. The resulting suspension was centrifuged at 13,000 rpm in microfuge tubes and the supernatant put through a 0.2 micron syringe filter. The filtrate was bound to Macro Prep DEAE beads (BioRad, Hercules, CA). The beads were extensively washed with 20 mM Tris pH 7.5 and bound proteins eluted with 1M NaCl. The 1M NaCl elute was dialyzed overnight against 10 mM Tris, pH 7.5. Dialyzed solution was treated with DNase and RNase at 0.05 mg/ml for 30 min. at room temperature and then with α -D-mannosidase, 0.5 U/mg at pH 4.5 for 3-4 hours at room temperature. After returning to pH 7.5, the material was fractionated via FPLC over a Bio Scale-Q-20 column (BioRad). Fractions were combined into nine pools, concentrated in a Centriprep 10 (Amicon, Beverley, MA) and then screened by Western blot for serological activity using a serum pool from *M. tuberculosis*-infected patients which was not immunoreactive with other antigens of the present invention.

The most reactive fraction was run in SDS-PAGE and transferred to PVDF. A band at approximately 85 Kd was cut out yielding the sequence:

(m) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID No. 137), wherein Xaa may be any amino acid.

Comparison of this sequence with those in the gene bank as described above, revealed no significant homologies to known sequences.

A DNA sequence that encodes the antigen designated as (m) above was obtained by screening a genomic *M. tuberculosis* Erdman strain library using labeled degenerate oligonucleotides corresponding to the N-terminal sequence of SEQ ID NO:137. A clone was identified having the DNA sequence provided in SEQ ID NO:

203. This sequence was found to encode the amino acid sequence provided in SEQ ID NO: 204. Comparison of these sequences with those in the genebank revealed some similarity to sequences previously identified in *M. tuberculosis* and *M. bovis*.

5

EXAMPLE 3

PREPARATION OF DNA SEQUENCES ENCODING *M. TUBERCULOSIS* ANTIGENS

This example illustrates the preparation of DNA sequences encoding *M. tuberculosis* antigens by screening a *M. tuberculosis* expression library with sera
10 obtained from patients infected with *M. tuberculosis*, or with anti-sera raised against soluble *M. tuberculosis* antigens.

A. PREPARATION OF *M. TUBERCULOSIS* SOLUBLE ANTIGENS USING RABBIT ANTI-SERA RAISED AGAINST *M. TUBERCULOSIS* SUPERNATANT

15 Genomic DNA was isolated from the *M. tuberculosis* strain H37Ra. The DNA was randomly sheared and used to construct an expression library using the Lambda ZAP expression system (Stratagene, La Jolla, CA). Rabbit anti-sera was generated against secretory proteins of the *M. tuberculosis* strains H37Ra, H37Rv and Erdman by immunizing a rabbit with concentrated supernatant of the *M. tuberculosis*
20 cultures. Specifically, the rabbit was first immunized subcutaneously with 200 µg of protein antigen in a total volume of 2 ml containing 10 µg muramyl dipeptide (Calbiochem, La Jolla, CA) and 1 ml of incomplete Freund's adjuvant. Four weeks later the rabbit was boosted subcutaneously with 100 µg antigen in incomplete Freund's adjuvant. Finally, the rabbit was immunized intravenously four weeks later with 50 µg
25 protein antigen. The anti-sera were used to screen the expression library as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989. Bacteriophage plaques expressing immunoreactive antigens were purified. Phagemid from the plaques was rescued and the nucleotide sequences of the *M. tuberculosis* clones deduced.

30

Thirty two clones were purified. Of these, 25 represent sequences that have not been previously identified in human *M. tuberculosis*. Recombinant antigens

were expressed and purified antigens used in the immunological analysis described in Example 1. Proteins were induced by IPTG and purified by gel elution, as described in Skeiky et al., *J. Exp. Med.* 181:1527-1537, 1995. Representative sequences of DNA molecules identified in this screen are provided in SEQ ID Nos.: 1-25. The
5 corresponding predicted amino acid sequences are shown in SEQ ID Nos. 63-87.

On comparison of these sequences with known sequences in the gene bank using the databases described above, it was found that the clones referred to hereinafter as TbRA2A, TbRA16, TbRA18, and TbRA29 (SEQ ID Nos. 76, 68, 70, 75) show some homology to sequences previously identified in *Mycobacterium leprae* but
10 not in *M. tuberculosis*. TbRA11, TbRA26, TbRA28 and TbDPEP (SEQ ID Nos.: 65, 73, 74, 53) have been previously identified in *M. tuberculosis*. No significant homologies were found to TbRA1, TbRA3, TbRA4, TbRA9, TbRA10, TbRA13, TbRA17, TbRa19, TbRA29, TbRA32, TbRA36 and the overlapping clones TbRA35 and TbRA12 (SEQ ID Nos. 63, 77, 81, 82, 64, 67, 69, 71, 75, 78, 80, 79, 66). The
15 clone TbRa24 is overlapping with clone TbRa29.

The results of PBMC proliferation and interferon- γ assays performed on representative recombinant antigens, and using T-cell preparations from several different *M. tuberculosis*-immune patients, are presented in Tables 2 and 3, respectively.

TABLE 2
RESULTS OF PBMC PROLIFERATION TO REPRESENTATIVE SOLUBLE ANTIGENS

Antigen	Patient												
	1	2	3	4	5	6	7	8	9	10	11	12	13
TbRa1	-	-	±	++	-	-	±	±	-	-	+	±	-
TbRa3	-	±	++	-	±	-	-	++	±	-	-	-	-
TbRa9	-	-	nt	nt	++	++	nt	nt	nt	nt	nt	nt	nt
TbRa10	-	-	±	±	±	+	nt	±	-	+	±	±	-
TbRa11	±	±	+	++	++	+	nt	-	++	++	++	±	nt
TbRa12	-	-	+	+	±	++	+	±	±	-	+	-	-
TbRa16	nt	nt	nt	nt	-	+	nt	nt	nt	nt	nt	nt	nt
TbRa24	nt	nt	nt	nt	-	-	nt	nt	nt	nt	nt	nt	nt
TbRa26	-	+	nt	nt	-	-	nt	nt	nt	nt	nt	nt	nt
TbRa29	nt	nt	nt	nt	-	-	nt	nt	nt	nt	nt	nt	nt
TbRa35	++	nt	++	++	++	++	nt	++	++	++	++	++	nt
TbRaB	nt	nt	nt	nt	-	-	nt	nt	nt	nt	nt	nt	nt
TbRaC	nt	nt	nt	nt	-	-	nt	nt	nt	nt	nt	nt	nt
TbRaD	nt	nt	nt	nt	-	-	nt	nt	nt	nt	nt	nt	nt
AAMK	-	-	±	-	-	-	nt	-	-	-	nt	±	nt
YY	-	-	-	-	-	-	nt	-	-	-	nt	+	nt
DPEP	-	+	-	++	-	-	nt	++	±	+	±	±	nt
Control	-	-	-	-	-	-	-	-	-	-	-	-	-

nt = not tested

100

100

In Tables 2 and 3, responses that gave a stimulation index (SI) of between 1.2 and 2 (compared to cells cultured in medium alone) were scored as \pm , a SI of 2-4 was scored as +, as SI of 4-8 or 2-4 at a concentration of 1 μ g or less was scored as ++ and an SI of greater than 8 was scored as +++. In addition, the effect of
5 concentration on proliferation and interferon- γ production is shown for two of the above antigens in the attached Figure. For both proliferation and interferon- γ production, TbRa3 was scored as ++ and TbRa9 as +.

These results indicate that these soluble antigens can induce proliferation and/or interferon- γ production in T-cells derived from an *M. tuberculosis*-immune
10 individual.

B. USE OF SERA FROM PATIENTS HAVING PULMONARY OR PLEURAL TUBERCULOSIS
TO IDENTIFY DNA SEQUENCES ENCODING *M. TUBERCULOSIS* ANTIGENS

The genomic DNA library described above, and an additional H37Rv
15 library, were screened using pools of sera obtained from patients with active tuberculosis. To prepare the H37Rv library, *M. tuberculosis* strain H37Rv genomic DNA was isolated, subjected to partial *Sau*3A digestion and used to construct an expression library using the Lambda Zap expression system (Stratagene, La Jolla, Ca). Three different pools of sera, each containing sera obtained from three individuals with
20 active pulmonary or pleural disease, were used in the expression screening. The pools were designated TbL, TbM and TbH, referring to relative reactivity with H37Ra lysate (*i.e.*, TbL = low reactivity, TbM = medium reactivity and TbH = high reactivity) in both ELISA and immunoblot format. A fourth pool of sera from seven patients with active pulmonary tuberculosis was also employed. All of the sera lacked increased reactivity
25 with the recombinant 38 kD *M. tuberculosis* H37Ra phosphate-binding protein.

All pools were pre-adsorbed with *E. coli* lysate and used to screen the H37Ra and H37Rv expression libraries, as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989. Bacteriophage plaques expressing immunoreactive antigens were purified.
30 Phagemid from the plaques was rescued and the nucleotide sequences of the *M. tuberculosis* clones deduced.

Thirty two clones were purified. Of these, 31 represented sequences that had not been previously identified in human *M. tuberculosis*. Representative sequences of the DNA molecules identified are provided in SEQ ID Nos.: 26-51 and 105. Of these, TbH-8-2 (SEQ. ID NO. 105) is a partial clone of TbH-8. and TbH-4 (SEQ. ID NO. 43) and TbH-4-FWD (SEQ. ID NO. 44) are non-contiguous sequences from the same clone. Amino acid sequences for the antigens hereinafter identified as Tb38-1, TbH-4, TbH-8, TbH-9, and TbH-12 are shown in SEQ ID Nos.: 88-92. Comparison of these sequences with known sequences in the gene bank using the databases identified above revealed no significant homologies to TbH-4, TbH-8, TbH-9 and TbM-3, although weak homologies were found to TbH-9. TbH-12 was found to be homologous to a 34 kD antigenic protein previously identified in *M. paratuberculosis* (Acc. No. S28515). Tb38-1 was found to be located 34 base pairs upstream of the open reading frame for the antigen ESAT-6 previously identified in *M. bovis* (Acc. No. U34848) and in *M. tuberculosis* (Sorensen et al., *Infect. Immun.* 63:1710-1717, 1995).

Probes derived from Tb38-1 and TbH-9, both isolated from an H37Ra library, were used to identify clones in an H37Rv library. Tb38-1 hybridized to Tb38-1F2, Tb38-1F3, Tb38-1F5 and Tb38-1F6 (SEQ. ID NOS. 112, 113, 116, 118, and 119). (SEQ ID NOS. 112 and 113 are non-contiguous sequences from clone Tb38-1F2.) Two open reading frames were deduced in Tb38-1F2; one corresponds to Tb37FL (SEQ. ID. NO. 114), the second, a partial sequence, may be the homologue of Tb38-1 and is called Tb38-IN (SEQ. ID NO. 115). The deduced amino acid sequence of Tb38-1F3 is presented in SEQ. ID. NO. 117. A TbH-9 probe identified three clones in the H37Rv library: TbH-9-FL (SEQ. ID NO. 106), which may be the homologue of TbH-9 (R37Ra), TbH-9-1 (SEQ. ID NO. 108), and TbH-9-4 (SEQ. ID NO. 110), all of which are highly related sequences to TbH-9. The deduced amino acid sequences for these three clones are presented in SEQ ID NOS. 107, 109 and 111.

Further screening of the *M. tuberculosis* genomic DNA library, as described above, resulted in the recovery of ten additional reactive clones, representing seven different genes. One of these genes was identified as the 38 Kd antigen discussed

above, one was determined to be identical to the 14Kd alpha crystallin heat shock protein previously shown to be present in *M. tuberculosis*, and a third was determined to be identical to the antigen TbH-8 described above. The determined DNA sequences for the remaining five clones (hereinafter referred to as TbH-29, TbH-30, TbH-32 and
5 TbH-33) are provided in SEQ ID NO: 138-141, respectively, with the corresponding predicted amino acid sequences being provided in SEQ ID NO: 142-145, respectively. The DNA and amino acid sequences for these antigens were compared with those in the gene bank as described above. No homologies were found to the 5' end of TbH-29 (which contains the reactive open reading frame), although the 3' end of TbH-29
10 found to be identical to the *M. tuberculosis* cosmid Y227. TbH-32 and TbH-33 were found to be identical to the previously identified *M. tuberculosis* insertion element IS6110 and to the *M. tuberculosis* cosmid Y50, respectively. No significant homologies to TbH-30 were found.

Positive phagemid from this additional screening were used to infect *E.*
15 *coli* XL-1 Blue MRF', as described in Sambrook et al., *supra*. Induction of recombinant protein was accomplished by the addition of IPTG. Induced and uninduced lysates were run in duplicate on SDS-PAGE and transferred to nitrocellulose filters. Filters were reacted with human *M. tuberculosis* sera (1:200 dilution) reactive with TbH and a rabbit sera (1:200 or 1:250 dilution) reactive with the N-terminal 4 Kd portion of lacZ.
20 Sera incubations were performed for 2 hours at room temperature. Bound antibody was detected by addition of ¹²⁵I-labeled Protein A and subsequent exposure to film for variable times ranging from 16 hours to 11 days. The results of the immunoblots are summarized in Table 4.

TABLE 4

<u>Antigen</u>	Human M. tb <u>Sera</u>	Anti-lacZ <u>Sera</u>
TbH-29	45 Kd	45 Kd
TbH-30	No reactivity	29 Kd
TbH-32	12 Kd	12 Kd
TbH-33	16 Kd	16 Kd

Positive reaction of the recombinant human *M. tuberculosis* antigens with both the human *M. tuberculosis* sera and anti-lacZ sera indicate that reactivity of the human *M. tuberculosis* sera is directed towards the fusion protein. Antigens reactive with the anti-lacZ sera but not with the human *M. tuberculosis* sera may be the result of the human *M. tuberculosis* sera recognizing conformational epitopes, or the antigen-antibody binding kinetics may be such that the 2 hour sera exposure in the immunoblot is not sufficient.

The results of T-cell assays performed on Tb38-1, ESAT-6 and other representative recombinant antigens are presented in Tables 5A, B and 6, respectively, below:

TABLE 5A
RESULTS OF PBMC PROLIFERATION TO REPRESENTATIVE ANTIGENS

Antigen	Donor										
	1	2	3	4	5	6	7	8	9	10	11
Tb38.1	+++	+	-	-	-	++	-	+	-	++	+++
ESAT-6	+++	+	+	+	-	+	-	+	+	++	+++
TbH-9	++	++	-	++	±	±	++	++	++	++	++

TABLE 5B
RESULTS OF PBMC INTERFERON- γ PRODUCTION TO REPRESENTATIVE ANTIGENS

Antigen	Donor										
	1	2	3	4	5	6	7	8	9	10	11
Tb38.1	+++	+	-	+	+	+++	-	++	-	+++	+++
ESAT-6	+++	+	+	+	+-	+	-	+	+	+++	+++
TbH-9	++	++	-	+++	\pm	\pm	+++	+++	++	+++	++

5

TABLE 6
SUMMARY OF T-CELL RESPONSES TO REPRESENTATIVE ANTIGENS

Antigen	Proliferation			Interferon- γ			total
	patient 4	patient 5	patient 6	patient 4	patient 5	patient 6	
TbH9	++	++	++	+++	++	++	13
TbM7	-	+	-	++	+	-	4
TbH5	-	+	+	++	++	++	8
TbL23	-	+	\pm	++	++	+	7.5
TbH4	-	++	\pm	++	++	\pm	7
- control	-	-	-	-	-	-	0

10

These results indicate that both the inventive *M. tuberculosis* antigens and ESAT-6 can induce proliferation and/or interferon- γ production in T-cells derived from an *M. tuberculosis*-immune individual. To the best of the inventors' knowledge, ESAT-6 has not been previously shown to stimulate human immune responses

15

A set of six overlapping peptides covering the amino acid sequence of the antigen Tb38-1 was constructed using the method described in Example 6. The sequences of these peptides, hereinafter referred to as pep1-6, are provided in SEQ ID Nos. 93-98, respectively. The results of T-cell assays using these peptides are shown in Tables 7 and 8. These results confirm the existence, and help to localize T-cell epitopes within Tb38-1 capable of inducing proliferation and interferon- γ production in T-cells derived from an *M. tuberculosis* immune individual.

20

TABLE 7
RESULTS OF PBMC PROLIFERATION TO TB38-1 PEPTIDES

[illegible]

Studies were undertaken to determine whether the antigens TbH-9 and Tb38-1 represent cellular proteins or are secreted into *M. tuberculosis* culture media. In the first study, rabbit sera were raised against A) secretory proteins of *M. tuberculosis*, B) the known secretory recombinant *M. tuberculosis* antigen 85b, C) recombinant Tb38-1 and D) recombinant TbH-9, using protocols substantially the same as that as described in Example 3A. Total *M. tuberculosis* lysate, concentrated supernatant of *M. tuberculosis* cultures and the recombinant antigens 85b, TbH-9 and Tb38-1 were resolved on denaturing gels, immobilized on nitrocellulose membranes and duplicate blots were probed using the rabbit sera described above.

The results of this analysis using control sera (panel I) and antisera (panel II) against secretory proteins, recombinant 85b, recombinant Tb38-1 and recombinant TbH-9 are shown in Figures 3A-D, respectively, wherein the lane designations are as follows: 1) molecular weight protein standards; 2) 5 μ g of *M. tuberculosis* lysate; 3) 5 μ g secretory proteins; 4) 50 ng recombinant Tb38-1; 5) 50 ng recombinant TbH-9; and 6) 50 ng recombinant 85b. The recombinant antigens were engineered with six terminal histidine residues and would therefore be expected to migrate with a mobility approximately 1 kD larger than the native protein. In Figure 3D, recombinant TbH-9 is lacking approximately 10 kD of the full-length 42 kD antigen, hence the significant difference in the size of the immunoreactive native TbH-9 antigen in the lysate lane (indicated by an arrow). These results demonstrate that Tb38-1 and TbH-9 are intracellular antigens and are not actively secreted by *M. tuberculosis*.

The finding that TbH-9 is an intracellular antigen was confirmed by determining the reactivity of TbH-9-specific human T cell clones to recombinant TbH-9, secretory *M. tuberculosis* proteins and PPD. A TbH-9-specific T cell clone (designated 131TbH-9) was generated from PBMC of a healthy PPD-positive donor. The proliferative response of 131TbH-9 to secretory proteins, recombinant TbH-9 and a control *M. tuberculosis* antigen, TbRa11, was determined by measuring uptake of tritiated thymidine, as described in Example 1. As shown in Figure 4A, the clone 131TbH-9 responds specifically to TbH-9, showing that TbH-9 is not a significant component of *M. tuberculosis* secretory proteins. Figure 4B shows the production of IFN- γ by a second TbH-9-specific T cell clone

(designated PPD 800-10) prepared from PBMC from a healthy PPD-positive donor, following stimulation of the T cell clone with secretory proteins, PPD or recombinant TbH-9. These results further confirm that TbH-9 is not secreted by *M. tuberculosis*.

5 C. USE OF SERA FROM PATIENTS HAVING EXTRAPULMONARY TUBERCULOSIS TO IDENTIFY
DNA SEQUENCES ENCODING *M. TUBERCULOSIS* ANTIGENS

Genomic DNA was isolated from *M. tuberculosis* Erdman strain, randomly sheared and used to construct an expression library employing the Lambda ZAP expression
10 system (Stratagene, La Jolla, CA). The resulting library was screened using pools of sera obtained from individuals with extrapulmonary tuberculosis, as described above in Example 3B, with the secondary antibody being goat anti-human IgG + A + M (H+L) conjugated with alkaline phosphatase.

Eighteen clones were purified. Of these, 4 clones (hereinafter referred to as
15 XP14, XP24, XP31 and XP32) were found to bear some similarity to known sequences. The determined DNA sequences for XP14, XP24 and XP31 are provided in SEQ ID Nos.: 156-158, respectively, with the 5' and 3' DNA sequences for XP32 being provided in SEQ ID Nos.: 159 and 160, respectively. The predicted amino acid sequence for XP14 is provided in SEQ ID No: 161. The reverse complement of XP14 was found to encode the amino acid
20 sequence provided in SEQ ID No.: 162.

Comparison of the sequences for the remaining 14 clones (hereinafter referred to as XP1-XP6, XP17-XP19, XP22, XP25, XP27, XP30 and XP36) with those in the genebank as described above, revealed no homologies with the exception of the 3' ends of XP2 and XP6 which were found to bear some homology to known *M. tuberculosis* cosmids.
25 The DNA sequences for XP27 and XP36 are shown in SEQ ID Nos.: 163 and 164, respectively, with the 5' sequences for XP4, XP5, XP17 and XP30 being shown in SEQ ID Nos: 165-168, respectively, and the 5' and 3' sequences for XP2, XP3, XP6, XP18, XP19, XP22 and XP25 being shown in SEQ ID Nos: 169 and 170; 171 and 172; 173 and 174; 175 and 176; 177 and 178; 179 and 180; and 181 and 182, respectively. XP1 was found to
30 overlap with the DNA sequences for TbH4, disclosed above. The full-length DNA sequence for TbH4-XP1 is provided in SEQ ID No.: 183. This DNA sequence was found to contain an

open reading frame encoding the amino acid sequence shown in SEQ ID No: 184. The reverse complement of TbH4-XP1 was found to contain an open reading frame encoding the amino acid sequence shown in SEQ ID No.: 185. The DNA sequence for XP36 was found to contain two open reading frames encoding the amino acid sequence shown in SEQ ID Nos.: 186 and 187, with the reverse complement containing an open reading frame encoding the amino acid sequence shown in SEQ ID No.: 188.

Recombinant XP1 protein was prepared as described above in Example 3B, with a metal ion affinity chromatography column being employed for purification. As illustrated in Figures 8A-B and 9A-B, using the assays described herein, recombinant XP1 was found to stimulate cell proliferation and IFN- γ production in T cells isolated from an *M. tuberculosis*-immune donors.

D. PREPARATION OF *M. TUBERCULOSIS* SOLUBLE ANTIGENS USING RABBIT ANTI-SERA RAISED AGAINST *M. TUBERCULOSIS* FRACTIONATED PROTEINS

M. tuberculosis lysate was prepared as described above in Example 2. The resulting material was fractionated by HPLC and the fractions screened by Western blot for serological activity with a serum pool from *M. tuberculosis*-infected patients which showed little or no immunoreactivity with other antigens of the present invention. Rabbit anti-sera was generated against the most reactive fraction using the method described in Example 3A. The anti-sera was used to screen an *M. tuberculosis* Erdman strain genomic DNA expression library prepared as described above. Bacteriophage plaques expressing immunoreactive antigens were purified. Phagemid from the plaques was rescued and the nucleotide sequences of the *M. tuberculosis* clones determined.

Ten different clones were purified. Of these, one was found to be TbRa35, described above, and one was found to be the previously identified *M. tuberculosis* antigen, HSP60. Of the remaining eight clones, seven (hereinafter referred to as RDIF2, RDIF5, RDIF8, RDIF10, RDIF11 and RDIF 12) were found to bear some similarity to previously identified *M. tuberculosis* sequences. The determined DNA sequences for RDIF2, RDIF5, RDIF8, RDIF10 and RDIF11 are provided in SEQ ID Nos.: 189-193, respectively, with the corresponding predicted amino acid sequences being provided in SEQ ID Nos: 194-198, respectively. The 5' and 3' DNA sequences for RDIF12 are provided in SEQ ID Nos.: 199

and 200, respectively. No significant homologies were found to the antigen RDIF-7. The determined DNA and predicted amino acid sequences for RDIF7 are provided in SEQ ID Nos.: 201 and 202, respectively. One additional clone, referred to as RDIF6 was isolated, however, this was found to be identical to RDIF5.

5 Recombinant RDIF6, RDIF8, RDIF10 and RDIF11 were prepared as described above. As shown in Figures 8A-B and 9A-B, these antigens were found to stimulate cell proliferation and IFN- γ production in T cells isolated from *M. tuberculosis*-immune donors.

10

EXAMPLE 4

PURIFICATION AND CHARACTERIZATION OF A POLYPEPTIDE FROM TUBERCULIN PURIFIED PROTEIN DERIVATIVE

15 An *M. tuberculosis* polypeptide was isolated from tuberculin purified protein derivative (PPD) as follows.

PPD was prepared as published with some modification (Seibert, F. et al., Tuberculin purified protein derivative. Preparation and analyses of a large quantity for standard. The American Review of Tuberculosis 44:9-25, 1941).

20 *M. tuberculosis* Rv strain was grown for 6 weeks in synthetic medium in roller bottles at 37°C. Bottles containing the bacterial growth were then heated to 100° C in water vapor for 3 hours. Cultures were sterile filtered using a 0.22 μ filter and the liquid phase was concentrated 20 times using a 3 kD cut-off membrane. Proteins were precipitated once with 50% ammonium sulfate solution and eight times with 25% ammonium sulfate solution. The resulting proteins (PPD) were fractionated by reverse phase liquid chromatography (RP-
25 HPLC) using a C18 column (7.8 x 300 mM; Waters, Milford, MA) in a Biocad HPLC system (Perseptive Biosystems, Framingham, MA). Fractions were eluted from the column with a linear gradient from 0-100% buffer (0.1% TFA in acetonitrile). The flow rate was 10 ml/minute and eluent was monitored at 214 nm and 280 nm.

30 Six fractions were collected, dried, suspended in PBS and tested individually in *M. tuberculosis*-infected guinea pigs for induction of delayed type hypersensitivity (DTH)

reaction. One fraction was found to induce a strong DTH reaction and was subsequently fractionated further by RP-HPLC on a microbore Vydac C18 column (Cat. No. 218TP5115) in a Perkin Elmer/Applied Biosystems Division Model 172 HPLC. Fractions were eluted with a linear gradient from 5-100% buffer (0.05% TFA in acetonitrile) with a flow rate of 80 μ l/minute. Eluent was monitored at 215 nm. Eight fractions were collected and tested for induction of DTH in *M. tuberculosis*-infected guinea pigs. One fraction was found to induce strong DTH of about 16 mm induration. The other fractions did not induce detectable DTH. The positive fraction was submitted to SDS-PAGE gel electrophoresis and found to contain a single protein band of approximately 12 kD molecular weight.

This polypeptide, herein after referred to as DPPD, was sequenced from the amino terminal using a Perkin Elmer/Applied Biosystems Division Procise 492 protein sequencer as described above and found to have the N-terminal sequence shown in SEQ ID No.: 129. Comparison of this sequence with known sequences in the gene bank as described above revealed no known homologies. Four cyanogen bromide fragments of DPPD were isolated and found to have the sequences shown in SEQ ID Nos.: 130-133.

The ability of the antigen DPPD to stimulate human PBMC to proliferate and to produce IFN- γ was assayed as described in Example 1. As shown in Table 9, DPPD was found to stimulate proliferation and elicit production of large quantities of IFN- γ ; more than that elicited by commercial PPD.

TABLE 9RESULTS OF PROLIFERATION AND INTERFERON- γ ASSAYS TO DPPD

PBMC Donor	Stimulator	Proliferation (CPM)	IFN- γ (OD ₄₅₀)
A	Medium	1,089	0.17
	PPD (commercial)	8,394	1.29
	DPPD	13,451	2.21
B	Medium	450	0.09
	PPD (commercial)	3,929	1.26
	DPPD	6,184	1.49
C	Medium	541	0.11
	PPD (commercial)	8,907	0.76
	DPPD	23,024	>2.70

5

EXAMPLE 5USE OF REPRESENTATIVE ANTIGENS FOR DIAGNOSIS OF TUBERCULOSIS

This example illustrates the effectiveness of several representative polypeptides in skin tests for the diagnosis of *M. tuberculosis* infection.

Individuals were injected intradermally with 100 μ l of either PBS or PBS plus Tween 20TM containing either 0.1 μ g of protein (for TbH-9 and TbRa35) or 1.0 μ g of protein (for TbRa38-1). Induration was measured between 5-7 days after injection, with a response of 5 mm or greater being considered positive. Of the 20 individuals tested, 2 were PPD negative and 18 were PPD positive. Of the PPD positive individuals, 3 had active tuberculosis, 3 had been previously infected with tuberculosis and 9 were healthy. In a second study, 13 PPD positive individuals were tested with 0.1 μ g TbRa11 in either PBS or PBS plus Tween 20TM as described above. The results of both studies are shown in Table 10.

TABLE 10
RESULTS OF DTH TESTING WITH REPRESENTATIVE ANTIGENS

	TbH-9 Pos/Total	Tb38-1 Pos/Total	TbRa35 Pos/Total	Cumulative Pos/Total	TbRa11 Pos/Total
PPD negative	0/2	0/2	0/2	0/2	
PPD positive					
healthy	5/9	4/9	4/9	6/9	1/4
prior TB	3/5	2/5	2/5	4/5	3/5
active	3/4	3/4	0/4	4/4	1/4
TOTAL	11/18	9/18	6/18	14/18	5/13

5

EXAMPLE 6

SYNTHESIS OF SYNTHETIC POLYPEPTIDES

10 Polypeptides may be synthesized on a Millipore 9050 peptide synthesizer using Fmoc chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using the following

15 cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0%-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to elute the

20 peptides. Following lyophilization of the pure fractions, the peptides may be characterized using electrospray mass spectrometry and by amino acid analysis.

EXAMPLE 7PREPARATION AND CHARACTERIZATION OF *M. TUBERCULOSIS* FUSION PROTEINS

A fusion protein containing TbRa3, the 38 kD antigen and Tb38-1 was
5 prepared as follows.

Each of the DNA constructs TbRa3, 38 kD and Tb38-1 were modified by PCR
in order to facilitate their fusion and the subsequent expression of the fusion protein TbRa3-
38 kD-Tb38-1. TbRa3, 38 kD and Tb38-1 DNA was used to perform PCR using the primers
PDM-64 and PDM-65 (SEQ ID NO: 146 and 147), PDM-57 and PDM-58 (SEQ ID NO: 148
10 and 149), and PDM-69 and PDM-60 (SEQ ID NO: 150 and 151), respectively. In each case,
the DNA amplification was performed using 10 µl 10X Pfu buffer, 2 µl 10 mM dNTPs, 2 µl
each of the PCR primers at 10 µM concentration, 81.5 µl water, 1.5 µl Pfu DNA polymerase
(Stratagene, La Jolla, CA) and 1 µl DNA at either 70 ng/µl (for TbRa3) or 50 ng/µl (for 38
kD and Tb38-1). For TbRa3, denaturation at 94°C was performed for 2 min, followed by 40
15 cycles of 96°C for 15 sec and 72°C for 1 min, and lastly by 72°C for 4 min. For 38 kD,
denaturation at 96°C was performed for 2 min, followed by 40 cycles of 96°C for 30 sec,
68°C for 15 sec and 72°C for 3 min, and finally by 72°C for 4 min. For Tb38-1 denaturation
at 94°C for 2 min was followed by 10 cycles of 96°C for 15 sec, 68°C for 15 sec and 72°C for
1.5 min, 30 cycles of 96°C for 15 sec, 64°C for 15 sec and 72°C for 1.5, and finally by 72°C
20 for 4 min.

The TbRa3 PCR fragment was digested with NdeI and EcoRI and cloned
directly into pT7⁺L2 IL 1 vector using NdeI and EcoRI sites. The 38 kD PCR fragment was
digested with Sse8387I, treated with T4 DNA polymerase to make blunt ends and then
digested with EcoRI for direct cloning into the pT7⁺L2Ra3-1 vector which was digested with
25 StuI and EcoRI. The 38-1 PCR fragment was digested with Eco47III and EcoRI and directly
subcloned into pT7⁺L2Ra3/38kD-17 digested with the same enzymes. The whole fusion was
then transferred to pET28b – using NdeI and EcoRI sites. The fusion construct was
confirmed by DNA sequencing.

The expression construct was transformed into BLR pLys S *E. coli* (Novagen,
30 Madison, WI) and grown overnight in LB broth with kanamycin (30 µg/ml) and
chloramphenicol (34 µg/ml). This culture (12 ml) was used to inoculate 500 ml 2XYT with

the same antibiotics and the culture was induced with IPTG at an OD₅₆₀ of 0.44 to a final concentration of 1.2 mM. Four hours post-induction, the bacteria were harvested and sonicated in 20 mM Tris (8.0), 100 mM NaCl, 0.1% DOC, 20 µg/ml Leupeptin, 20 mM PMSF followed by centrifugation at 26,000 X g. The resulting pellet was resuspended in 8 M urea, 20 mM Tris (8.0), 100 mM NaCl and bound to Pro-bond nickel resin (Invitrogen, Carlsbad, CA). The column was washed several times with the above buffer then eluted with an imidazole gradient (50 mM, 100 mM, 500 mM imidazole was added to 8 M urea, 20 mM Tris (8.0), 100 mM NaCl). The eluates containing the protein of interest were then dialyzed against 10 mM Tris (8.0).

The DNA and amino acid sequences for the resulting fusion protein (hereinafter referred to as TbRa3-38 kD-Tb38-1) are provided in SEQ ID NO: 152 and 153, respectively.

A fusion protein containing the two antigens TbH-9 and Tb38-1 (hereinafter referred to as TbH9-Tb38-1) without a hinge sequence, was prepared using a similar procedure to that described above. The DNA sequence for the TbH9-Tb38-1 fusion protein is provided in SEQ ID NO: 156.

The ability of the fusion protein TbH9-Tb38-1 to induce T cell proliferation and IFN-γ production in PBMC preparations was examined using the protocol described above in Example 1. PBMC from three donors were employed: one who had been previously shown to respond to TbH9 but not Tb38-1 (donor 131); one who had been shown to respond to Tb38-1 but not TbH9 (donor 184); and one who had been shown to respond to both antigens (donor 201). The results of these studies (Figs. 5-7, respectively) demonstrate the functional activity of both the antigens in the fusion protein.

A fusion protein containing TbRa3, the antigen 38kD, Tb38-1 and DPEP was prepared as follows.

Each of the DNA constructs TbRa3, 38 kD and Tb38-1 were modified by PCR and cloned into vectors essentially as described above, with the primers PDM-69 (SEQ ID NO:150 and PDM-83 (SEQ ID NO: 205) being used for amplification of the Tb38-1A fragment. Tb38-1A differs from Tb38-1 by a *DraI* site at the 3' end of the coding region that

keeps the final amino acid intact while creating a blunt restriction site that is in frame. The TbRa3/38kD/Tb38-1A fusion was then transferred to pET28b using NdeI and EcoRI sites.

DPEP DNA was used to perform PCR using the primers PDM-84 and PDM-85 (SEQ ID NO: 206 and 207, respectively) and 1 μ l DNA at 50 ng/ μ l. Denaturation at 94 °C was performed for 2 min, followed by 10 cycles of 96 °C for 15 sec, 68 °C for 15 sec and 72 °C for 1.5 min; 30 cycles of 96 °C for 15 sec, 64 °C for 15 sec and 72 °C for 1.5 min; and finally by 72 °C for 4 min. The DPEP PCR fragment was digested with EcoRI and Eco72I and clones directly into the pET28Ra3/38kD/38-1A construct which was digested with DraI and EcoRI. The fusion construct was confirmed to be correct by DNA sequencing.

10 Recombinant protein was prepared as described above. The DNA and amino acid sequences for the resulting fusion protein (hereinafter referred to as TbF-2) are provided in SEQ ID NO: 208 and 209, respectively.

The reactivity of the fusion protein TbF-2 with sera from *M. tuberculosis*-infected patients was examined by ELISA using the protocol described above. The results of these studies (Table 11) demonstrate that all four antigens function independently in the fusion protein.

15

TABLE 11
 REACTIVITY OF TbF-2 FUSION RECOMBINANT WITH TB AND NORMAL SERA

Serum ID	Status	TbF OD450	Status	TbF-2 OD450	Status	ELISA Reactivity			
						38 kD	TbRa3	Tb38-1	DPEP
B931-40	TB	0.57	+	0.321	+	-	+	-	+
B931-41	TB	0.601	+	0.396	+	+	+	+	-
B931-109	TB	0.494	+	0.404	+	+	+	±	-
B931-132	TB	1.502	+	1.292	+	+	+	+	±
5004	TB	1.806	+	1.666	+	±	±	+	-
15004	TB	2.862	+	2.468	+	+	+	+	-
39004	TB	2.443	+	1.722	+	+	+	+	-
68004	TB	2.871	+	2.575	+	+	+	+	-
99004	TB	0.691	+	0.971	+	-	±	+	-
107004	TB	0.875	+	0.732	+	-	±	+	-
92004	TB	1.632	+	1.394	+	+	±	±	-
97004	TB	1.491	+	1.979	+	+	±	-	+
118004	TB	3.182	+	3.045	+	+	±	-	-
173004	TB	3.644	+	3.578	+	+	+	+	-
175004	TB	3.332	+	2.916	+	+	+	-	-
274004	TB	3.696	+	3.716	+	-	+	-	+
276004	TB	3.243	+	2.56	+	-	-	+	-
282004	TB	1.249	+	1.234	+	+	-	-	-
289004	TB	1.373	+	1.17	+	-	+	-	-
308004	TB	3.708	+	3.355	+	-	-	+	-
314004	TB	1.663	+	1.399	+	-	-	+	-
317004	TB	1.163	+	0.92	+	+	-	-	-
312004	TB	1.709	+	1.453	+	-	+	-	-
380004	TB	0.238	-	0.461	+	-	±	-	+
451004	TB	0.18	-	0.2	-	-	-	-	±
478004	TB	0.188	-	0.469	+	-	-	-	±
410004	TB	0.384	+	2.392	+	±	-	-	+
411004	TB	0.306	+	0.874	+	-	+	-	+
421004	TB	0.357	+	1.456	+	-	+	-	+
528004	TB	0.047	-	0.196	-	-	-	-	+
A6-87	Normal	0.094	-	0.063	-	-	-	-	-
A6-88	Normal	0.214	-	0.19	-	-	-	-	-
A6-89	Normal	0.248	-	0.125	-	-	-	-	-
A6-90	Normal	0.179	-	0.206	-	-	-	-	-
A6-91	Normal	0.135	-	0.151	-	-	-	-	-
A6-92	Normal	0.064	-	0.097	-	-	-	-	-
A6-93	Normal	0.072	-	0.098	-	-	-	-	-
A6-94	Normal	0.072	-	0.064	-	-	-	-	-
A6-95	Normal	0.125	-	0.159	-	-	-	-	-
A6-96	Normal	0.121	-	0.12	-	-	-	-	-
Cut-off		0.284		0.266					

One of skill in the art will appreciate that the order of the individual antigens within the fusion protein may be changed and that comparable activity would be expected provided each of the epitopes is still functionally available. In addition, truncated forms of the proteins containing active epitopes may be used in the construction of fusion proteins.

5

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purpose of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANTS: Reed, Steven G.
Skeiky, Yasir A.W.
Dillon, Davin C.
Campos-Neto, Antonio
Houghton, Raymond
Vedvick, Thomas S.
Twardzik, Daniel R.
Lodes, Michael J.
- (ii) TITLE OF INVENTION: COMPOUNDS AND METHODS FOR IMMUNOTHERAPY
AND DIAGNOSIS OF TUBERCULOSIS
- (iii) NUMBER OF SEQUENCES: 214
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: SEED and BERRY LLP
 - (B) STREET: 6300 Columbia Center, 701 Fifth Avenue
 - (C) CITY: Seattle
 - (D) STATE: Washington
 - (E) COUNTRY: USA
 - (F) ZIP: 98104-7092
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 01-OCT-1997
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Maki, David J.
 - (B) REGISTRATION NUMBER: 31,392
 - (C) REFERENCE/DOCKET NUMBER: 210121.411C7
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (206) 622-4900
 - (B) TELEFAX: (206) 682-6031

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 766 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGAGGCACCG GTAGTTTGAA CCAAACGCAC AATCGACGGG CAAACGAACG GAAGAACACA	60
ACCATGAAGA TGGTGAAATC GATCGCCGCA GGTCTGACCG CCGCGGCTGC AATCGGCGCC	120
GCTGCGGCCG GTGTGACTTC GATCATGGCT GGCGGCCCGG TCGTATACCA GATGCAGCCG	180
GTCGTCTTCG GCGCGCCACT GCCGTTGGAC CCGGCATCCG CCCCTGACGT CCCGACCGCC	240
GCCCAGTTGA CCAGCCTGCT CAACAGCCTC GCCGATCCCA ACGTGTCGTT TGCGAACAAG	300
GGCAGTCTGG TCGAGGGCGG CATCGGGGGC ACCGAGGCGC GCATCGCCGA CCACAAGCTG	360
AAGAAGGCCG CCGAGCACGG GGATCTGCCG CTGTCTGTTCA GCGTGACGAA CATCCAGCCG	420
GCGGCCGCCG GTTCGGCCAC CGCCGACGTT TCCGTCTCGG GTCCGAAGCT CTCGTGCGCG	480
GTCACGCAGA ACGTCACGTT CGTGAATCAA GGCGGCTGGA TGCTGTCACG CGCATCGGCG	540
ATGGAGTTGC TGCAGGCCGC AGGGNAACTG ATTGGCGGGC CGGNTTCAGC CCGCTGTTCA	600
GCTACGCCGC CCGCCTGGTG ACGCGTCCAT GTCGAACACT CGCGCGTGTA GCACGGTGCG	660
GTNTGCGCAG GGNCGCACGC ACCGCCCGGT GCAAGCCGTC CTCGAGATAG GTGGTGNCTC	720
GNCACCAGNG ANCACCCCN NNTCGNCNNT TCTCGNTGNT GNATGA	766

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 752 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGCATCACC ATCACCATCA CGATGAAGTC ACGGTAGAGA CGACCTCCGT CTTCCGCGCA	60
GACTTCCTCA GCGAGCTGGA CGCTCCTGCG CAAGCGGGTA CGGAGAGCGC GGTCTCCGGG	120
GTGGAAGGGC TCCCGCCGGG CTCGGCGTTG CTGGTAGTCA AACGAGGCCC CAACGCCGGG	180
TCCCGGTTCC TACTCGACCA AGCCATCACG TCGGCTGGTC GGCATCCCGA CAGCGACATA	240
TTTETCGACG ACGTGACCGT GAGCCGTCGC CATGCTGAAT TCCGGTTGGA AAACAACGAA	300
TTCAATGTCG TCGATGTCGG GAGTCTCAAC GGCACCTACG TCAACCGCGA GCCCGTGGAT	360
TCGGCGGTGC TGGCGAACGG CGACGAGGTC CAGATCGGCA AGCTCCGGTT GGTGTTCTTG	420
ACCGGACCCA AGCAAGGCGA GGATGACGGG AGTACCGGGG GCCCGTGAGC GCACCCGATA	480
GCCCCGCGCT GGCCGGGATG TCGATCGGGG CGGTCCTCCG ACCTGCTACG ACCGGATTTT	540
CCCTGATGTC CACCATCTCC AAGATTCGAT TCTTGGGAGG CTTGAGGGTC NGGGTGACCC	600
CCCCGCGGGC CTCATTCTGG GGTNTCGGCN GGTTTCACCC CNTACCNACT GCCNCCCGGN	660

TTGCNAATTC NTTCTTCNCT GCCCNNAAG GGACCNTTAN CTTGCCGCTN GAAANGGTNA 720
TCCNGGGCCC NTCCTNGAAN CCCCNTCCCC CT 752

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 813 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CATATGCATC ACCATCACCA TCACACTTCT AACCGCCCAG CGCGTCGGGG GCGTCGAGCA 60
CCACGCGACA CCGGGCCCGA TCGATCTGCT AGCTTGAGTC TGGTCAGGCA TCGTCGTCAG 120
CAGCGCGATG CCCTATGTTT GTCGTCGACT CAGATATCGC GGCAATCCAA TCTCCCGCCT 180
GCGGCCGGCG GTGCTGCAAA CTA CTACTCCCGG AGGAATTTTCG ACGTGCGCAT CAAGATCTTC 240
ATGCTGGTCA CGGCTGTCGT TTTGCTCTGT TGTTCCGGGTG TGGCCACGGC CGCGCCCAAG 300
ACCTACTGCG AGGAGTTGAA AGGCACCGAT ACCGGCCAGG CGTGCCAGAT TCAAATGTCC 360
GACCCGGCCT ACAACATCAA CATCAGCCTG CCCAGTTACT ACCCCGACCA GAAGTCGCTG 420
GAAAATTACA TCGCCAGAC GCGCGACAAG TTCCTCAGCG CGGCCACATC GTCCACTCCA 480
CGCGAAGCCC CCTACGAATT GAATATCACC TCGGCCACAT ACCAGTCCGC GATACCGCCG 540
CGTGGTACGC AGGCCGTGGT GCTCAMGGTC TACCACAACG CCGGCGGCAC GCACCCAACG 600
ACCACGTACA AGGCCTTCGA TTGGGACCAG GCCTATCGCA AGCCAATCAC CTATGACACG 660
CTGTGGCAGG CTGACACCGA TCCGCTGCCA GTCGTCTTCC CCATTGTTGC AAGGTGAACT 720
GAGCAACGCA GACCGGGACA ACWGGTATCG ATAGCCGCCN AATGCCGGCT TGGAAACCCNG 780
TGAAATTATC ACAACTTCGC AGTCACNAAA NAA 813

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 447 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CGGTATGAAC ACGGCCGCGT CCGATAACTT CCAGCTGTCC CAGGGTGGGC AGGGATTTCG 60

CATTCCGATC GGGCAGGCGA TGGCGATCGC GGGCCAGATC CGATCGGGTG GGGGGTCACC	120
CACCGTTCAT ATCGGGCCTA CCGCCTTCCT CGGCTTGGGT GTTGTGACACA ACAACGGCAA	180
CGGCGCACGA GTCCAACGCG TGGTCGGGAG CGCTCCGGCG GCAAGTCTCG GCATCTCCAC	240
CGGCGACGTG ATCACC GCGG TCGACGGCGC TCCGATCAAC TCGGCCACCG CGATGGCGGA	300
CGCGCTTAAC GGGCATCATC CCGGTGACGT CATCTCGGTG AACTGGCAAA CCAAGTCGGG	360
CGGCACGCGT ACAGGGAACG TGACATTGGC CGAGGGACCC CCGGCCTGAT TTCGTCGYGG	420
ATACCACCCG CCGGCCGGCC AATTGGA	447

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 604 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GTCCCACTGC GGTGCGCCGAG TATGTCGCCC AGCAAATGTC TGGCAGCCGC CCAACGGAAT	60
CCGGTGATCC GACGTGCGAG GTTGTGCAAC CCGCCGCCGC GGAAGTATCG GTCCATGCCT	120
AGCCCGGCGA CGGCGAGCGC CGGAATGGCG CGAGTGAGGA GGCGGGCAAT TTGGCGGGGC	180
CCGGCGACGG NGAGCGCCGG AATGGCGCGA GTGAGGAGGT GGNCAATCAT GCCCAGNGTG	240
ATCCAATCAA CCTGNATTCG GNCTGNNGGN CCATTTGACA ATCGAGGTAG TGAGCGCAAA	300
TGAATGATGG AAAACGGGNG GNGACGTCCG NTGTTCTGGT GGTGNTAGGT GNCTGNCTGG	360
NGTNGNGGNT ATCAGGATGT TCTTCGNCGA AANCTGATGN CGAGGAACAG GGTGTNCCCG	420
NNANNCCNAN GGNGTCCNAN CCCNNNNNTCC TCGNCGANAT CANANAGNCG NTTGATGNGA	480
NAAAAGGGTG GANCAGNNNN AANTNGNGGN CCNAANAANC NNNANNGNNG NNAGNTNGNT	540
NNNTNTTNNC ANNNNNNNNTG NNGNNGNNCN NNNCAANCNN NTNNNNGNAA NNGGNTTNTT	600
NAAT	604

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 633 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TTGCANGTCG AACCACCTCA CTAAAGGGAA CAAAAGCTNG AGCTCCACCG CGGTGGCGGC	60
CGCTCTAGAA CTAGTGKATM YYYCKGGCTG CAGSAATYCG GYACGAGCAT TAGGACAGTC	120
TAACGGTCCT GTTACGGTGA TCGAATGACC GACGACATCC TGCTGATCGA CACCGACGAA	180
CGGGTGCGAA CCCTCACCTT CAACCGGCCG CAGTCCCGYA ACGCGCTCTC GGCGGCGCTA	240
CGGGATCGGT TTTTCGCGGY GTTGGYCGAC GCCGAGGYCG ACGACGACAT CGACGTCGTC	300
ATCCTCACCG GYGCCGATCC GGTGTTCTGC GCCGACTGG ACCTCAAGGT AGCTGGCCGG	360
GCAGACCGCG CTGCCGGACA TCTCACCGCG GTGGGCGGCC ATGACCAAGC CGGTGATCGG	420
CGCGATCAAC GGCGCCGCGG TCACCGGCGG GCTCGAACTG GCGCTGTACT GCGACATCCT	480
GATCGCCTCC GAGCAGCCC GCTTCGNCGA CACCCACGCC CGGGTGGGGC TGCTGCCCCA	540
CTGGGGACTC AGTGTGTGCT TGCCGCAAAA GGTCGGCATC GGNCTGGGCC GGTGGATGAG	600
CCTGACCGGC GACTACCTGT CCGTGACCGA CGC	633

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1362 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGACGACGAC GGCGCCGGAG AGCGGGCGCG AACGGCGATC GACGCGGCC TGGCCAGAGT	60
CGGCAACCACC CAGGAGGGAG TCGAATCATG AAATTTGTCA ACCATATTGA GCGCGTCGCG	120
CCCCGCCGAG CCGGCGGCGC GGTCGCCGAG GTCTATGCCG AGGCCCGCCG CGAGTTCGGC	180
CGGCTGCCCC AGCCGCTCGC CATGCTGTCC CCGGACGAGG GACTGCTCAC CGCCGGCTGG	240
GCGACGTTGC GCGAGAACT GCTGGTGGGC CAGGTGCCGC GTGGCCGCAA GGAAGCCGTC	300
GCCGCCGCCG TCGCGGCCAG CCTGCGCTGC CCCTGGTGCG TCGACGCACA CACCACCATG	360
CTGTACGCGG CAGGCCAAAC CGACACCGCC GCGGCGATCT TGGCCGGCAC AGCACCTGCC	420
GCCGGTGACC CGAACGCGCC GATGTGGCG TGGGCGGCAG GAACCGGAC ACCGGCGGGA	480
CCGCCGGCAC CGTTCGGCCC GGATGTCGCC GCCGAATACC TGGGCACCGC GGTGCAATTC	540
CACTTCATCG CACGCCTGGT CCTGGTGCTG CTGGACGAAA CCTTCCTGCC GGGGGGCCCC	600
CGCGCCCAAC AGCTCATGCG CCGCGCCGGT GGACTGGTGT TCGCCCGCAA GGTGCGCGCG	660
GAGCATCGGC CGGGCCGCTC CACCCGCCGG CTCGAGCCGC GAACGCTGCC CGACGATCTG	720

GCATGGGCAA CACCGTCCGA GCCCATAGCA ACCGCGTTCG CCGCGCTCAG CCACCACCTG	780
GACACCGCGC CGCACCTGCC GCCACCGACT CGTCAGGTGG TCAGGCGGGT CGTGGGGTCG	840
TGGCACGGCG AGCCAATGCC GATGAGCAGT CGCTGGACGA ACGAGCACAC CGCCGAGCTG	900
CCCGCCGACC TGCACGCGCC CACCCGTCTT GCCCTGCTGA CCGGCCTGGC CCCGCATCAG	960
GTGACCGACG ACGACGTGCG CGCGGCCCCGA TCCCTGCTCG ACACCGATGC GGCCTGGTT	1020
GGCGCCCTGG CCTGGGCGCG CTTCACCGCC GCGCGGCGCA TCGGCACCTG GATCGGCGCC	1080
GCCGCCGAGG GCCAGGTGTC GCGGCAAAAC CCGACTGGGT GAGTGTGCGC GCCCTGTCGG	1140
TAGGGTGTCA TCGCTGGCCC GAGGGATCTC GCGGCGGCGA ACGGAGGTGG CGACACAGGT	1200
GGAAGCTGCG CCCACTGGCT TGCGCCCAA CGCCGTCGTG GGCGTTCGGT TGGCCGCACT	1260
GGCCGATCAG GTCGGCGCCG GCCCTTGGCC GAAGGTCCAG CTCAACGTGC CGTCACCGAA	1320
GGACCGGACG GTCACCGGGG GTCACCCTGC GCGCCAAGG AA	1362

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1458 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCGACGACCC CGATATGCCG GGCACCGTAG CGAAAGCCGT CGCCGACGCA CTCGGGCGCG	60
GTATCGCTCC CGTTGAGGAC ATTCAGGACT GCGTGAGGC CCGGCTGGGG GAAGCCGGTC	120
TGGATGACGT GGCCCGTGTT TACATCATCT ACCGGCAGCG GCGCGCCGAG CTGCGGACGG	180
CTAAGGCCTT GCTCGGCGTG CGGGACGAGT TAAAGCTGAG CTTGGCGGCC GTGACGGTAC	240
TGCGCGAGCG CTATCTGCTG CACGACGAGC AGGGCCGGCC GGCCGAGTCG ACCGGCGAGC	300
TGATGGACCG ATCGGCGCGC TGTGTCGCGG CGGCCGAGGA CCAGTATGAG CCGGGCTCGT	360
CGAGGCGGTG GGCCGAGCGG TTCGCCACGC TATTACGCAA CCTGGAATTC CTGCCGAATT	420
CGCCACGTT GATGAACTCT GGCACCGACC TGGGACTGCT CGCCGGCTGT TTTGTTCTGC	480
CGATTGAGGA TTCGCTGCAA TCGATCTTTG CGACGCTGGG ACAGGCCGCC GAGCTGCAGC	540
GGGCTGGAGG CGGCACCGGA TATGCGTTCA GCCACCTGCG ACCCGCCGGG GATCGGGTGG	600
CCTCCACGGG CGGCACGGCC AGCGGACCGG TGTCGTTTCT ACGGCTGTAT GACAGTGCCG	660
CGGGTGTGGT CTCCATGGGC GGTCGCCGGC GTGGCGCCTG TATGGCTGTG CTTGATGTGT	720
CGCACCCGGA TATCTGTGAT TTCGTCACCG CCAAGGCCGA ATCCCCAGC GAGCTCCCGC	780

ATTTCAACCT ATCGGTTGGT GTGACCGACG CGTTCCTGCG GGCCGTCGAA CGCAACGGCC	840
TACACCGGCT GGTCAATCCG CGAACCGGCA AGATCGTCGC GCGGATGCCC GCCGCCGAGC	900
TGTTCGACGC CATCTGCAAA GCCGCGCACG CCGGTGGCGA TCCCGGGCTG GTGTTTCTCG	960
ACACGATCAA TAGGGCAAAC CCGGTGCCGG GGAGAGGCCG CATCGAGGCG ACCAACCCGT	1020
GCGGGGAGGT CCCACTGCTG CCTTACGAGT CATGTAATCT CGGCTCGATC AACCTCGCCC	1080
GGATGCTCGC CGACGGTCGC GTCGACTGGG ACCGGCTCGA GGAGGTCGCC GGTGTGGCGG	1140
TGCGGTTTCT TGATGACGTC ATCGATGTCA GCCGCTACCC CTTCCCCGAA CTGGGTGAGG	1200
CGGCCCCGCG CACCCGCAAG ATCGGGCTGG GAGTCATGGG TTTGGCGGAA CTGCTTGCCG	1260
CACTGGGTAT TCCGTACGAC AGTGAAGAAG CCGTGCGGTT AGCCACCCGG CTCATGCGTC	1320
GCATACAGCA GGCGGCGCAC ACGGCATCGC GGAGGCTGGC CGAAGAGCGG GGCGCATTC	1380
CGGCGTTCAC CGATAGCCGG TTCGCGCGGT CGGGCCCGAG GCGCAACGCA CAGGTCACCT	1440
CCGTCGCTCC GACGGGCA	1458

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 862 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ACGGTGTAAT CGTGCTGGAT CTGGAACCGC GTGGCCCGCT ACCTACCGAG ATCTACTGGC	60
GGCGCAGGGG GCTGGCCCTG GGCATCGCGG TCGTCGTAGT CGGGATCGCG GTGGCCATCG	120
TCATCGCCTT CGTCGACAGC AGCGCCGGTG CCAAACCGGT CAGCGCCGAC AAGCCGGCCT	180
CCGCCCAGAG CCATCCGGGC TCGCCGGCAC CCAAGCACC CCAGCCGGCC GGGCAAACCG	240
AAGGTAACGC CGCCGCGGCC CCGCCGAGG GCCAAAACCC CGAGACACCC ACGCCACCG	300
CCGCGGTGCA GCCGCCGCCG GTGCTCAAGG AAGGGGACGA TTGCCCCGAT TCGACGCTGG	360
CCGTCAAAGG TTTGACCAAC GCGCCGAGT ACTACGTCGG CGACCAGCCG AAGTTCACCA	420
TGGTGGTCAC CAACATCGGC CTGGTGTCTT GTAAACGCGA CGTTGGGGCC GCGGTGTTGG	480
CCGCCTACGT TTA CTGCTG GACAACAAGC GGTGTGGTCAACCTGGAC TGC GCGCCCT	540
CGAATGAGAC GCTGGTCAAG ACGTTTTCCC CCGGTGAGCA GGTAACGACC GCGGTGACCT	600
GGACCGGGAT GGGATCGGCG CCGCGCTGCC CATTGCCGCG GCCGGCGATC GGGCCGGGCA	660
CCTACAATCT CGTGGTACAA CTGGGCAATC TGC GCTCGCT GCCGGTTCCG TTCATCCTGA	720

ATCAGCCGCC GCGCCGCCC GGGCCGGTAC CCGCTCCGGG TCCAGCGCAG GCGCCTCCGC	780
CGGAGTCTCC CGCGCAAGGC GGATAATTAT TGATCGCTGA TGGTCGATTC CGCCAGCTGT	840
GACAACCCCT CGCCTCGTGC CG	862

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 622 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

TTGATCAGCA CCGGCAAGGC GTCACATGCC TCCCTGGGTG TGCAGGTGAC CAATGACAAA	60
GACACCCCGG GCGCCAAGAT CGTCGAAGTA GTGGCCGGTG GTGCTGCCGC GAACGCTGGA	120
GTGCCGAAGG GCGTCGTTGT CACCAAGGTC GACGACCGCC CGATCAACAG CGCGGACGCG	180
TTGGTTGCCG CCGTGCGGTC CAAAGCGCCG GCGGCCACGG TGGCGCTAAC CTTTCAGGAT	240
CCCTCGGGCG GTAGCCGCAC AGTGCAAGTC ACCCTCGGCA AGGCGGAGCA GTGATGAAGG	300
TCGCCGCGCA GTGTTCAAAG CTCGGATATA CGGTGGCACC CATGGAACAG CGTGCGGAGT	360
TGGTGTTGG CCGGGCACTT GTCGTCGTCG TTGACGATCG CACGGCGCAC GGCGATGAAG	420
ACCACAGCGG GCCGCTTGTC ACCGAGCTGC TCACCGAGGC CGGGTTTGTT GTCGACGGCG	480
TGGTGGCGGT GTCGGCCGAC GAGGTCGAGA TCCGAAATGC GCTGAACACA GCGGTGATCG	540
GCGGGGTGGA CCTGGTGGTG TCGGTCGGCG GGACCGGNGT GACGNCTCGC GATGTCACCC	600
CGGAAGCCAC CCGNGACATT CT	622

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1200 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GGCGCAGCGG TAAGCCTGTT GGCCGCCGGC AACTGGTGT TGACAGCATG CGGCGGTGGC	60
ACCAACAGCT CGTCGTCAGG CGCAGGCGGA ACGTCTGGGT CGGTGCACTG CGGCGGCAAG	120
AAGGAGCTCC ACTCCAGCGG CTCGACCGCA CAAGAAAATG CCATGGAGCA GTTCGTCTAT	180

GCCTACGTGC	GATCGTGCCC	GGGCTACACG	TTGGACTACA	ACGCCAACGG	GTCCGGTGCC	240
GGGGTGACCC	AGTTTCTCAA	CAACGAAACC	GATTTCGCCG	GCTCGGATGT	CCCGTTGAAT	300
CCGTGACCG	GTCAACCTGA	CCGGTCGGCG	GAGCGGTGCG	GTTCCCCGGC	ATGGGACCTG	360
CCGACGGTGT	TCGGCCCCGAT	CGCGATCACC	TACAATATCA	AGGGCGTGAG	CACGCTGAAT	420
CTTGACGGAC	CCACTACCGC	CAAGATTTTC	AACGGCACCA	TCACCGTGTG	GAATGATCCA	480
CAGATCCAAG	CCCTCAACTC	CGGCACCGAC	CTGCCGCCAA	CACCGATTAG	CGTTATCTTC	540
CGCAGCGACA	AGTCCGGTAC	GTCGGACAAC	TTCCAGAAAT	ACCTCGACGG	TGTATCCAAC	600
GGGGCGTGGG	GCAAAGGCGC	CAGCGAAACG	TTCAGCGGGG	GCGTCGGCGT	CGGCGCCAGC	660
GGGAACAACG	GAACGTCGGC	CCTACTGCAG	ACGACCGACG	GGTCGATCAC	CTACAACGAG	720
TGGTCGTTTG	CGGTGGGTAA	GCAGTTGAAC	ATGGCCCAGA	TCATCACGTC	GGCGGGTCCG	780
GATCCAGTGG	CGATCACCAC	CGAGTCGGTC	GGTAAGACAA	TCGCCGGGGC	CAAGATCATG	840
GGACAAGGCA	ACGACCTGGT	ATTGGACACG	TCGTCGTTCT	ACAGACCCAC	CCAGCCTGGC	900
TCTTACCCGA	TCGTGCTGGC	GACCTATGAG	ATCGTCTGCT	CGAAATACCC	GGATGCGACG	960
ACCGGTACTG	CGGTAAGGGC	GTTTATGCAA	GCCGCGATTG	GTCCAGGCCA	AGAAGGCCTG	1020
GACCAATACG	GCTCCATTCC	GTTGCCCAAA	TCGTTCCAAG	CAAAATTGGC	GGCCGCGGTG	1080
AATGCTATTT	CTTGACCTAG	TGAAGGGAAT	TCGACGGTGA	GCGATGCCGT	TCCGCAGGTA	1140
GGGTCGCAAT	TTGGGCCGTA	TCAGCTATTG	CGGCTGCTGG	GCCGAGGCGG	GATGGGCGAG	1200

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1155 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GCAAGCAGCT	GCAGGTCGTG	CTGTTGACG	AACTGGGCAT	GCCGAAGACC	AAACGCACCA	60
AGACCGGCTA	CACCACGGAT	GCCGACGCGC	TGCAGTCGTT	GTTCGACAAG	ACCGGGCATC	120
CGTTTCTGCA	ACATCTGCTC	GCCCACCGCG	ACGTCACCCG	GCTCAAGGTC	ACCGTCGACG	180
GTTTGCTCCA	AGCGGTGGCC	GCCGACGGCC	GCATCCACAC	CACGTTCAAC	CAGACGATCG	240
CCGCGACCGG	CCGGCTCTCC	TCGACCGAAC	CCAACCTGCA	GAACATCCCG	ATCCGCACCG	300
ACGCGGGCCG	GCGGATCCGG	GACGCGTTG	TGGTCGGGGA	CGGTTACGCC	GAGTTGATGA	360
CGGCCGACTA	CAGCCAGATC	GAGATGCGGA	TCATGGGGCA	CCTGTCCGGG	GACGAGGGCC	420

TCATCGAGGC GTTCAACACC GGGGAGGACC TGTATTTCGTT CGTCGCGTCC CGGGTGTTCG	480
GTGTGCCCCAT CGACGAGGTC ACCGGCGAGT TGCGGCGCCG GGTCAAGGCG ATGTCCTACG	540
GGCTGGTTTA CGGGTTGAGC GCCTACGGCC TGTCGCAGCA GTTGAAAATC TCCACCGAGG	600
AAGCCAACGA GCAGATGGAC GCGTATTTTCG CCCGATTTCGG CGGGGTGCGC GACTACCTGC	660
GCGCCGTAAGT CGAGCGGGCC CGCAAGGACG GCTACACCTC GACGGTGCTG GGCCGTCGCC	720
GCTACCTGCC CGAGCTGGAC AGCAGCAACC GTCAAGTGCG GGAGGCCGCC GAGCGGGCGG	780
CGCTGAACGC GCCGATCCAG GGCAGCGCGG CCGACATCAT CAAGGTGGCC ATGATCCAGG	840
TCGACAAGGC GCTCAACGAG GCACAGCTGG CGTCGCGCAT GCTGCTGCAG GTCCACGACG	900
AGCTGCTGTT CGAAATCGCC CCCGGTGAAC GCGAGCGGGT CGAGGCCCTG GTGCGCGACA	960
AGATGGGCGG CGCTTACCCG CTCGACGTCC CGCTGGAGGT GTCGGTGGGC TACGGCCGCA	1020
GCTGGGACGC GGC GGCGGCAC TGAGTGCCGA GCGTGATCT GGGGCGGGAA TTCGGCGATT	1080
TTTCCGCCCT GAGTTCACGC TCGGCGCAAT CGGGACCGAG TTTGTCCAGC GTGTACCCGT	1140
CGAGTAGCCT CGTCA	1155

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1771 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GAGCGCCGTC TGGTGTTTGA ACGGTTTTAC CGGTCGGCAT CGGCACGGGC GTTGCCGGGT	60
TCGGGCCTCG GGTGCGGAT CGTCAACAG GTGGTGCTCA ACCACGGCGG ATTGCTGCGC	120
ATCGAAGACA CCGACCCAGG CGGCCAGCCC CCTGGAACGT CGATTTACGT GCTGCTCCCC	180
GGCCGTCGGA TGCCGATTCC GCAGCTTCCC GGTGCGACGG CTGGCGCTCG GAGCACGGAC	240
ATCGAGAACT CTCGGGGTTC GGCGAACGTT ATCTCAGTGG AATCTCAGTC CACGCGCGCA	300
ACCTAGTTGT GCAGTTACTG TTGAAAGCCA CACCCATGCC AGTCCACGCA TGGCCAAGTT	360
GGCCCGAGTA GTGGGCCTAG TACAGGAAGA GCAACCTAGC GACATGACGA ATCACCACG	420
GATTTCGCCA CCGCCGCAGC AGCCGGGAAC CCCAGGTTAT GCTCAGGGGC AGCAGCAAAC	480
GTACAGCCAG CAGTTCGACT GGC GTTACCC ACCGTCCCCG CCCCCGCAGC CAACCCAGTA	540
CCGTCAACCC TACGAGGCGT TGGGTGGTAC CCGGCCGGGT CTGATACCTG GCGTGATTCC	600
GACCATGACG CCCCCTCCTG GGATGGTTCG CCAACGCCCT CGTGCAGGCA TGTTGGCCAT	660

CGGCGCGGTG	ACGATAGCGG	TGGTGTCCGC	CGGCATCGGC	GGCGCGGCCG	CATCCCTGGT	720
CGGGTTCAAC	CGGGCACCCG	CCGGCCCCAG	CGGCGGCCCA	GTGGCTGCCA	GCGCGGCGCC	780
AAGCATCCCC	GCAGCAAACA	TGCCGCCGGG	GTCGGTCGAA	CAGGTGGCGG	CCAAGGTGGT	840
GCCCAGTGTC	GTCATGTTGG	AAACCGATCT	GGGCCGCCAG	TCGGAGGAGG	GCTCCGGCAT	900
CATTCTGTCT	GCCGAGGGGC	TGATCTTGAC	CAACAACCAC	GTGATCGCGG	CGGCCGCCAA	960
GCCTCCCCTG	GGCAGTCCGC	CGCCGAAAAC	GACGGTAACC	TTCTCTGACG	GGCGGACCGC	1020
ACCCTTCACG	GTGGTGCGGG	CTGACCCAC	CAGTGATATC	GCCGTCGTCC	GTGTTCAGGG	1080
CGTCTCCGGG	CTCACCCCGA	TCTCCCTGGG	TTCCTCCTCG	GACCTGAGGG	TCGGTCAGCC	1140
GGTGCTGGCG	ATCGGGTCGC	CGCTCGGTTT	GGAGGGCACC	GTGACCACGG	GGATCGTCAG	1200
CGCTCTCAAC	CGTCCAGTGT	CGACGACCGG	CGAGGCCGGC	AACCAGAACA	CCGTGCTGGA	1260
CGCCATTAG	ACCGACGCCG	CGATCAACCC	CGGTAACTCC	GGGGGCGCGC	TGGTGAACAT	1320
GAACGCTCAA	CTCGTCGGAG	TCAACTCGGC	CATTGCCACG	CTGGGCGCGG	ACTCAGCCGA	1380
TGCGCAGAGC	GGCTCGATCG	GTCTCGGTTT	TGCGATTCCA	GTGACCAGG	CCAAGCGCAT	1440
CGCCGACGAG	TTGATCAGCA	CCGGCAAGGC	GTCACATGCC	TCCCTGGGTG	TGCAGGTGAC	1500
CAATGACAAA	GACACCCCGG	GCGCCAAGAT	CGTCGAAGTA	GTGGCCGGTG	GTGCTGCCGC	1560
GAACGCTGGA	GTGCCGAAGG	GCGTCGTTGT	CACCAAGGTC	GACGACCGCC	CGATCAACAG	1620
CGCGGACGCG	TTGGTTGCCG	CCGTGCGGTC	CAAAGCGCCG	GGCGCCACGG	TGGCGCTAAC	1680
CTTTCAGGAT	CCCTCGGGCG	GTAGCCGCAC	AGTGCAAGTC	ACCCTCGGCA	AGGCGGAGCA	1740
GTGATGAAGG	TCGCCGCGCA	GTGTTCAAAG	C			1771

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1058 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CTCCACCGCG	GTGGCGGCCG	CTCTAGAACT	AGTGGATCCC	CCGGGCTGCA	GGAATTCGGC	60
ACGAGGATCC	GACGTCGCAG	GTTGTGGAAC	CCGCCGCCGC	GGAAGTATCG	GTCCATGCCT	120
AGCCCGGCGA	CGGCGAGCGC	CGGAATGGCG	CGAGTGAGGA	GGCGGGCAAT	TTGGCGGGGC	180
CCGGCGACGG	CGAGCGCCGG	AATGGCGCGA	GTGAGGAGGC	GGGCAGTCAT	GCCCAGCGTG	240
ATCCAATCAA	CCTGCATTCT	GCCTGCGGGC	CCATTTGACA	ATCGAGGTAG	TGAGCGCAAA	300

TGAATGATGG AAAACGGGCG GTGACGTCCG CTGTTCTGGT GGTGCTAGGT GCCTGCCTGG	360
CGTTGTGGCT ATCAGGATGT TCTTCGCCGA AACCTGATGC CGAGGAACAG GGTGTTCCCG	420
TGAGCCCCGAC GGC GTCCGAC CCCGCGCTCC TCGCCGAGAT CAGGCAGTCG CTTGATGCGA	480
CAAAAGGGTT GACCAGCGTG CACGTAGCGG TCCGAACAAC CGGGAAAGTC GACAGCTTGC	540
TGGGTATTAC CAGTGCCGAT GTCGACGTCC GGGCCAATCC GCTCGCGGCA AAGGGCGTAT	600
GCACCTACAA CGACGAGCAG GGTGTCCCGT TTCGGGTACA AGGCGACAAC ATCTCGGTGA	660
AACTGTTCTGA CGACTGGAGC AATCTCGGCT CGATTTCTGA ACTGTCAACT TCACGCGTGC	720
TCGATCCTGC CGCTGGGGTG ACGCAGCTGC TGTCCGGTGT CACGAACCTC CAAGCGCAAG	780
GTACCGAAGT GATAGACGGA ATTTTCGACCA CCAAATCAC CGGGACCATC CCCGCGAGCT	840
CTGTCAAGAT GCTTGATCCT GGCGCCAAGA GTGCAAGGCC GGCGACCGTG TGGATTGCCC	900
AGGACGGCTC GCACCACCTC GTCCGAGCGA GCATCGACCT CGGATCCGGG TCGATTACAGC	960
TCACGCAGTC GAAATGGAAC GAACCCGTCA ACGTCGACTA GGCCGAAGTT GCGTCGACGC	1020
GTTGNTCGAA ACGCCCTTGT GAACGGTGTC AACGGNAC	1058

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 542 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GAATTCGGCA CGAGAGGTGA TCGACATCAT CGGGACCAGC CCCACATCCT GGGAACAGGC	60
GGCGGCGGAG GCGGTCCAGC GGGCGCGGGA TAGCGTCGAT GACATCCGCG TCGCTCGGGT	120
CATTGAGCAG GACATGGCCG TGGACAGCGC CGGCAAGATC ACCTACCGCA TCAAGCTCGA	180
AGTGTCGTTT AAGATGAGGC CGGCGCAACC GCGCTAGCAC GGGCCGGCGA GCAAGACGCA	240
AAATCGCACG GTTTGCGGTT GATTCGTGCG ATTTTGTGTC TGCTCGCCGA GGCCTACCAG	300
GCGCGGCCCA GGTCCGCGTG CTGCCGTATC CAGGCGTGCA TCGCGATTCC GGCGGCCACG	360
CCGGAGTTAA TGCTTCGCGT CGACCCGAAC TGGGCGATCC GCCGNGAGC TGATCGATGA	420
CCGTGGCCAG CCCGTCGATG CCCGAGTTGC CCGAGGAAAC GTGCTGCCAG GCCGGTAGGA	480
AGCGTCCGTA GGCGGCGGTG CTGACCGGCT CTGCCTGCGC CCTCAGTGCG GCCAGCGAGC	540
GG	542

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 913 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGGTGCCGCC CGCGCCTCCG TTGCCCCCAT TGCCGCCGTC GCCGATCAGC TGCGCATCGC	60
CACCATCACC GCCTTTGCCG CCGGCACCGC CGGTGGCGCC GGGGCCGCCG ATGCCACCGC	120
TTGACCCTGG CCGCCGGCGC CGCCATTGCC ATACAGCACC CCGCCGGGGG CACCGTTACC	180
GCCGTCGCCA CCGTCGCCGC CGCTGCCGTT TCAGGCCGGG GAGGCCGAAT GAACCGCCGC	240
CAAGCCCGCC GCCGGCACCG TTGCCGCCTT TTCCGCCCGC CCCGCCGGCG CCGCCAATTG	300
CCGAACAGCC AMGCACCGTT GCCGCCAGCC CCGCCGCCGT TAACGGCGCT GCCGGGCGCC	360
GCCGCCGGAC CCGCCATTAC CGCCGTTCCC GTTCGGTGCC CCGCCGTTAC CGGCGCCGCC	420
GTTTGCCGCC AATATTCGGC GGGCACCGCC AGACCCGCCG GGGCCACCAT TGCCGCCGGG	480
CACCGAAACA ACAGCCCAAC GGTGCCGCCG GCCCCGCCGT TTGCCGCCAT CACCGGCCAT	540
TCACCGCCAG CACCGCCGTT AATGTTTATG AACCCGGTAC CGCCAGCGCG GCCCCTATTG	600
CCGGGCGCCG GAGNGCGTGC CCGCCGGCGC CGCCAACGCC CAAAAGCCCG GGGTTGCCAC	660
CGGCCCCGCC GGACCCACCG GTCCCGCCGA TCCCCCGTT GCCGCCGGTG CCGCCGCCAT	720
TGGTGCTGCT GAAGCCGTTA GCGCCGGTTC CGCSGGTTCC GCGGGTGGCG CCNTGGCCGC	780
CGGCCCCGCC GTTGCCGTAC AGCCACCCCC CGGTGGCGCC GTTGCCGCCA TTGCCGCCAT	840
TGCCGCCGTT GCCGCCATTG CCGCCGTTCC CGCCGCCACC GCCGNTTGG CCGCCGGCGC	900
CGCCGGCGGC CGC	913

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1872 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GACTACGTTG GTGTAGAAAA ATCCTGCCGC CCGGACCCCT AAGGCTGGGA CAATTTCTGA	60
TAGTACCCC GACACAGGAG GTTACGGGAT GAGCAATTCT CGCCGCCGCT CACTCAGGTG	120

GTCATGGTTG CTGAGCGTGC TGGCTGCCGT CGGGCTGGGC CTGGCCACGG CGCCGGCCCA	180
GGCGGCCCCG CCGGCCTTGT CGCAGGACCG GTTCGCCGAC TTCCCCGCGC TGCCCCCTCGA	240
CCCGTCCGCG ATGGTCGCCC AAGTGGCGCC ACAGGTGGTC AACATCAACA CCAAACCTGGG	300
CTACAACAAC GCCGTGGGCG CCGGGACCGG CATCGTCATC GATCCCAACG GTGTCGTGCT	360
GACCAACAAC CACGTGATCG CGGGCGCCAC CGACATCAAT GCGTTCAGCG TCGGCTCCGG	420
CCAAACCTAC GCGTCGATG TGGTCGGGTA TGACCGCACC CAGGATGTCTG CGGTGCTGCA	480
GCTGCGCGGT GCCGGTGGCC TGCCGTCCGC GGCATCGGT GGCGGCGTCG CGGTTGGTGA	540
GCCCCGTCGTC GCGATGGGCA ACAGCGGTGG GCAGGGCGGA ACGCCCGTG CGGTGCCTGG	600
CAGGGTGGTC GCGCTCGGCC AAACCGTGCA GCGTCGGAT TCGCTGACCG GTGCCGAAGA	660
GACATTGAAC GGGTTGATCC AGTTCGATGC CGCAATCCAG CCCGGTGATT CGGGCGGGCC	720
CGTCGTCAAC GGCCTAGGAC AGGTGGTCGG TATGAACACG GCCGCGTCCG ATAACCTCCA	780
GCTGTCCCAG GGTGGGCAGG GATTCGCCAT TCCGATCGGG CAGGCGATGG CGATCGCGGG	840
CCAAATCCGA TCGGGTGGGG GGTCACCCAC CGTTCATATC GGGCCTACCG CCTTCCTCGG	900
CTTGGGTGTT GTCGACAACA ACGGCAACGG CGCACGAGTC CAACGCGTGG TCGGAAGCGC	960
TCCGGCGGCA AGTCTCGGCA TCTCCACCGG CGACGTGATC ACCGCGGTCTG ACGGCGCTCC	1020
GATCAACTCG GCCACCGCGA TGGCGGACGC GCTTAACGGG CATCATCCCG GTGACGTCAT	1080
CTCGGTGAAC TGGCAAACCA AGTCGGGCGG CACGCGTACA GGAACGTGA CATTGGCCGA	1140
GGGACCCCCG GCCTGATTTG TCGCGGATAC CACCCGCCGG CCGGCCAATT GGATTGGCGC	1200
CAGCCGTGAT TGCCGCGTGA GCCCCGAGT TCCGTCTCCC GTGCGCGTGG CATTGTGGAA	1260
GCAATGAACG AGGCAGAACA CAGCGTTGAG CACCCTCCCG TGCAGGGCAG TTACGTGCAA	1320
GGCGGTGTGG TCGAGCATCC GGATGCCAAG GACTTCGGCA GCGCCGCCGC CCTGCCCCGC	1380
GATCCGACCT GGTTTAAGCA CGCCGTCTTC TACGAGGTGC TGGTCCGGGC GTTCTTCGAC	1440
GCCAGCGCGG ACGGTTCCGN CGATCTGCGT GGA CTATCG ATCGCCTCGA CTACCTGCAG	1500
TGGCTTGGA TCGACTGCAT CTGTTGCCGC CGTTCCTACG ACTCACCCTG GCGCGACGGC	1560
GGTTACGACA TTCGCGACTT CTACAAGGTG CTGCCCCAAT TCGGCACCGT CGACGATTTT	1620
GTCGCCCTGG TCGACACCGC TCACCGGCGA GGTATCCGCA TCATCACC GA CCTGGTGATG	1680
AATCACACCT CGGAGTCGCA CCCCTGGTTT CAGGAGTCCC GCCGCGACCC AGACGGACCG	1740
TACGGTGACT ATTACGTGTG GAGCGACACC AGCGAGCGCT ACACCGACGC CCGGATCATC	1800
TTCGTGACCA CCGAAGAGTC GAACTGGTCA TTCGATCCTG TCCGCCGACA GTTNCTACTG	1860
GCACCGATTC TT	1872

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1482 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CTTCGCCGAA ACCTGATGCC GAGGAACAGG GTGTTCCCGT GAGCCCGACG GCGTCCGACC	60
CCGCGCTCCT CGCCGAGATC AGGCAGTCGC TTGATGCGAC AAAAGGGTTG ACCAGCGTGC	120
ACGTAGCGGT CCGAACAACC GGGAAAGTCG ACAGCTTGCT GGGTATTACC AGTGCCGATG	180
TCGACGTCCG GGCCAATCCG CTCGCGGCAA AGGGCGTATG CACCTACAAC GACGAGCAGG	240
GTGTCCCGTT TCGGGTACAA GCGACAACA TCTCGGTGAA ACTGTTCGAC GACTGGAGCA	300
ATCTCGGCTC GATTTCTGAA CTGTCAACTT CACGCGTGCT CGATCCTGCC GCTGGGGTGA	360
CGCAGCTGCT GTCCGGTGTC ACGAACCTCC AAGCGCAAGG TACCGAAGTG ATAGACGGAA	420
TTTCGACCAC CAAAATCACC GGGACCATCC CCGCGAGCTC TGTCAAGATG CTTGATCCTG	480
GCGCCAAGAG TGCAAGGCCG GCGACCGTGT GGATTGCCCA GGACGGCTCG CACCACCTCG	540
TCCGAGCGAG CATCGACCTC GGATCCGGGT CGATTGAGCT CACGCAGTCG AAATGGAACG	600
AACCCGTCAA CGTCGACTAG GCCGAAGTTG CGTCGACGCG TTGCTCGAAA CGCCCTTGTTG	660
AACGGTGTCA ACGGCACCCG AAAACTGACC CCCTGACGGC ATCTGAAAAT TGACCCCCTA	720
GACCGGGCGG TTGGTGGTTA TTCTTCGGTG GTTCCGGCTG GTGGGACGCG GCCGAGGTCG	780
CGGTCTTTGA GCCGGTAGCT GTCGCCTTTG AGGGCGACGA CTTGAGCATG GTGGACGAGG	840
CGGTCGATCA TGGCGGCAGC AACGACGTCG TCGCCGCCGA AAACCTCGCC CCACCGGCCG	900
AAGGCCTTAT TGGACGTGAC GATCAAGCTG GCCCGCTCAT ACCGGGAGGA CACCAGCTGG	960
AAGAAGAGGT TGGCGGCCTC GGGCTCAAAC GGAATGTAAAC CGACTTCGTC AACCACCAGG	1020
AGCGGATAGC GGCCAAACCG GGTGAGTTG GCGTAGATGC GCCCGGCGTG GTGAGCCTCG	1080
GCGAACCGTG CTACCCATTC GCGGGCGGTG GCGAACAGCA CCCGATGACC GGCCTGACAC	1140
GCGCGTATCG CCAGGCCGAC CGCAAGATGA GTCTTCCCGG TGCCAGGCGG GGCCCAAAAA	1200
CACGACGTTA TCGCGGGCGG TGATGAAATC CAGGGTGCCC AGATGTGCGA TGGTGTGCGG	1260
TTTGAGGCCA CGAGCATGCT CAAAGTCGAA CTCTTCCAAC GACTTCCGAA CCGGGAAGCG	1320
GGCGGCGCGG ATGCGGCCCT CACCACCATG GGAATCCCGG GCTGACACTT CCCGCTGCAG	1380
GCAGGCGGCC AGGTATTCTT CGTGGCTCCA GTTCTCGGCG CGGGCGCGAT CGGCCAGCCG	1440
GGAACTGAC TCACGCAGGG TGGGAGCTTT CAATGCTCTT GT	1482

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 876 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

```
GAATTCGGCA CGAGCCGGCG ATAGCTTCTG GGCCGCGGCC GACCAGATGG CTCGAGGGTT      60
CGTGCTCGGG GCCACCGCCG GGCGCACCAC CCTGACCGGT GAGGGCCTGC AACACGCCGA      120
CGGTCACTCG TTGCTGCTGG ACGCCACCAA CCCGGCGGTG GTTGCTACG ACCCGGCCTT      180
CGCCTACGAA ATCGGCTACA TCGNGGAAAG CGGACTGGCC AGGATGTGCG GGGAGAACCC      240
GGAGAACATC TTCTTCTACA TCACCGTCTA CAACGAGCCG TACGTGCAGC CGCCGGAGCC      300
GGAGAACTTC GATCCCGAGG GCGTGCTGGG GGGTATCTAC CGNTATCACG CGGCCACCGA      360
GCAACGCACC AACAAGGNGC AGATCCTGGC CTCCGGGGTA GCGATGCCCC CGGCGCTGCG      420
GGCAGCACAG ATGCTGGCCG CCGAGTGGGA TGTCGCCGCC GACGTGTGGT CGGTGACCAG      480
TTGGGGCGAG CTAAACCGCG ACGGGGTGGT CATCGAGACC GAGAAGCTCC GCCACCCCGA      540
TCGGCCGGCG GCGTGCCCT ACGTGACGAG AGCGCTGGAG AATGCTCGGG GCCCGGTGAT      600
CGCGGTGTCG GACTGGATGC GCGCGGTCCC CGAGCAGATC CGACCGTGGG TGCCGGGCAC      660
ATACCTCACG TTGGGCACCG ACGGGTTCGG TTTTCCGAC ACTCGGCCCG CCGGTCGTG      720
TTACTTCAAC ACCGACGCCG AATCCCAGGT TGGTCGCGGT TTTGGGAGGG GTTGGCCGGG      780
TCGACGGGTG AATATCGACC CATTCGGTGC CGGTCGTGGG CCGCCCCGCC AGTTACCCGG      840
ATTCGACGAA GGTGGGGGGT TGCGCCGAN TAAGTT      876
```

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1021 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

```
ATCCCCCGG GCTGCAGGAA TTCGGCACGA GAGACAAAAT TCCACGCGTT AATGCAGGAA      60
CAGATTCATA ACGAATTCAC AGCGGCACAA CAATATGTCG CGATCGCGGT TTATTTTCGAC      120
```

AGCGAAGACC TGCCGCAGTT GGCGAAGCAT TTTTACAGCC AAGCGGTCTGA GGAACGAAAC	180
CATGCAATGA TGCTCGTGCA ACACCTGCTC GACCGCGACC TTCGTGTCTGA AATTCCCGGC	240
GTAGACACGG TGCAGAACCA GTTCGACAGA CCCCCTGAGG CACTGGCGCT GGCCTCTGAT	300
CAGGAACGCA CAGTCACCGA CCAGGTCGGT CGGCTGACAG CGGTGGCCCG CGACGAGGGC	360
GATTTCTCTG GCGAGCAGTT CATGCAGTGG TTCTTGACAG AACAGATCTGA AGAGGTGGCC	420
TTGATGGCAA CCCTGGTGCG GGTTGCCGAT CGGGCCGGGG CCAACCTGTT CGAGCTAGAG	480
AACTTCGTCTG CACGTGAAGT GGATGTGGCG CCGGCCGCAT CAGGCGCCCC GCACGCTGCC	540
GGGGGCCGCC TCTAGATCCC TGGGGGGGAT CAGCGAGTGG TCCCGTTCGC CCGCCCGTCT	600
TCCAGCCAGG CCTTGGTGCG GCCGGGGTGG TGAGTACCAA TCCAGGCCAC CCCGACCTCC	660
CGGNAAAAGT CGATGTCCTC GTACTCATCG ACGTTCCAGG AGTACACCGC CCGGCCCTGA	720
GCTGCCGAGC GGTCAACGAG TTGCGGATAT TCCTTTAACG CAGGCAGTGA GGGTCCCACG	780
GCGGTTGGCC CGACCGCCGT GGCCGCACTG CTGGTCAGGT ATCGGGGGGT CTTGGCGAGC	840
AACAACGTCTG GCAGGAGGGG TGGAGCCCGC CGGATCCGCA GACCGGGGGG GCGAAAACGA	900
CATCAACACC GCACGGGATC GATCTGCGGA GGGGGGTGCG GGAATACCGA ACCGGTGTAG	960
GAGCGCCAGC AGTTGTTTTT CCACCAGCGA AGCGTTTTCTG GGTCATCGGN GGCNNTTAAG	1020
T	1021

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 321 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CGTGCCGACG AACGGAAGAA CACAACCATG AAGATGGTGA AATCGATCGC CGCAGGTCTG	60
ACCGCCGCGG CTGCAATCGG CGCCGCTGCG GCCGGTGTGA CTTGATCAT GGCTGGCGGN	120
CCGGTCGTAT ACCAGATGCA GCCGGTCGTC TTCGGCGCGC CACTGCCGTT GGACCCGGNA	180
TCCGCCCCCTG ANGTCCTGAC CGCCGCCAG TGGACCAGNC TGCTCAACAG NCTCGNCGAT	240
CCCAACGTGT CGTTTGNGAA CAAGGGNAGT CTGGTCGAGG GNNGNATCGG NGGNANCGAG	300
GGNGNGNATC GNCGANCACA A	321

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 373 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TCTTATCGGT TCCGGTTGGC GACGGGTTTT GGGNGCGGGT GGTAAACCCG CTCGGCCAGC	60
CGATCGACGG GCGCGGAGAC GTCGACTCCG ATACTCGGCG CGCGCTGGAG CTCCAGGCGC	120
CCTCGGTGGT GNACCGGCAA GCGTGAAGG AGCCGTTGNA GACCGGGATC AAGGCGATTG	180
ACGCGATGAC CCCGATCGGC CGCGGGCAGC GCCAGCTGAT CATCGGGGAC CGCAAGACCG	240
GCAAAAACCG CCGTCTGTGT CGGACACCAT CCTCAAACCA GCGGGAAGAA CTGGGAGTCC	300
GGTGGATCCC AAGAAGCAGG TGCGCTTGTG TATACGTTGG CCATCGGGCA AGAAGGGGAA	360
CTTACCATCG CCG	373

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 352 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GTGACGCCGT GATGGGATTC CTGGGCGGGG CCGGTCCGCT GGCGGTGGTG GATCAGCAAC	60
TGGTTACCCG GGTGCCGCAA GGCTGGTCGT TTGCTCAGGC AGCCGCTGTG CCGGTGGTGT	120
TCTTGACGGC CTGGTACGGG TTGGCCGATT TAGCCGAGAT CAAGGCGGGC GAATCGGTGC	180
TGATCCATGC CCGTACCGGC GGTGTGGGCA TGGCGGCTGT GCAGCTGGCT CGCCAGTGGG	240
GCGTGGAGGT TTTCGTCACC GCCAGCCGTG GNAAGTGGGA CACGCTGCGC GCCATNGNGT	300
TTGACGACGA NCCATATCGG NGATTCCNC ACATNCGAAG TTCCGANGGA GA	352

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 726 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GAAATCCGCG TTCATTCCGT TCGACCAGCG GCTGGCGATA ATCGACGAAG TGATCAAGCC	60
GCGGTTTCGCG GCGCTCATGG GTCACAGCGA GTAATCAGCA AGTTCTCTGG TATATCGCAC	120
CTAGCGTCCA GTTGCTTGCC AGATCGCTTT CGTACCGTCA TCGCATGTAC CGGTTCGCGT	180
GCCGCACGCT CATGCTGGCG GCGTGCATCC TGGCCACGGG TGTGGCGGGT CTCGGGGTCG	240
GCGCGCAGTC CGCAGCCCAA ACCGCGCCGG TGCCCGACTA CTACTGGTGC CCGGGGCAGC	300
CTTTCGACCC CGCATGGGGG CCCAACTGGG ATCCCTACAC CTGCCATGAC GACTTCCACC	360
GCGACAGCGA CGGCCCCGAC CACAGCCGCG ACTACCCCGG ACCCATCCTC GAAGGTCCCG	420
TGCTTGACGA TCCCGGTGCT GCGCCGCCGC CCCCGGCTGC CGGTGGCGGC GCATAGCGCT	480
CGTTGACCGG GCCGCATCAG CGAATACGCG TATAAACCCG GCGTGCCCC CGGCAAGCTA	540
CGACCCCGG CGGGGCAGAT TTACGCTCCC GTGCCGATGG ATCGCGCCGT CCGATGACAG	600
AAAATAGGCG ACGGTTTTTG CAACCGCTTG GAGGACGCTT GAAGGGAACC TGTCATGAAC	660
GGCGACAGCG CCTCCACCAT CGACATCGAC AAGGTTGTTA CCCGCACACC CGTTCGCCGG	720
ATCGTG	726

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 580 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CGCGACGACG ACGAACGTCG GGCCACCAC CGCCTATGCG TTGATGCAGG CGACCGGGAT	60
GGTCGCCGAC CATATCCAAG CATGCTGGGT GCCCACTGAG CGACCTTTTG ACCAGCCGGG	120
CTGCCCCGATG GCGGCCCGGT GAAGTCATTG CGCCGGGGCT TGTGCACCTG ATGAACCCGA	180
ATAGGGAACA ATAGGGGGGT GATTTGGCAG TTCAATGTCG GGTATGGCTG GAAATCCAAT	240
GGCGGGGCAT GCTCGGCGCC GACCAGGCTC GCGCAGGCGG GCCAGCCCGA ATCTGGAGGG	300
AGCACTCAAT GGCGGCGATG AAGCCCCGGA CCGGCGACGG TCCTTTGGAA GCAACTAAGG	360
AGGGGCGCGG CATTGTGATG CGAGTACCAC TTGAGGGTGG CGGTGCCTG GTCGTCGAGC	420
TGACACCCGA CGAAGCCGCC GCACTGGGTG ACGAACTCAA AGGCGTTACT AGCTAAGACC	480
AGCCCAACGG CGAATGGTCG GCGTTACGCG CACACCTTCC GGTAGATGTC CAGTGTCTGC	540
TCGGCGATGT ATGCCAGGA GAACTCTTGG ATACAGCGCT	580

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 160 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

```
AACGGAGGCG CCGGGGGTTT TGGCGGGGCC GGGGCGGTCG GCGGCAACGG CGGGGCCGGC      60
GGTACCGCCG GGTGTGTCGG TGTCGGCGGG GCCGGTGGGG CCGGAGGCAA CGGCATCGCC      120
GGTGTACAGG GTACGTCGGC CAGCACACCG GGTGGATCCG                               160
```

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 272 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

```
GACACCGATA CGATGGTGAT GTACGCCAAC GTTGTGACGA CGCTCGAGGC GTTCACGATC      60
CAGCGCACAC CCGACGGCGT GACCATCGGC GATGCGGCCC CGTTCGCGGA GGCGGCTGCC      120
AAGGCGATGG GAATCGACAA GCTGCGGGTA ATTCATACCG GAATGGACCC CGTCGTCGCT      180
GAACGCGAAC AGTGGGACGA CGGCAACAAC ACGTTGGCGT TGGCGCCCGG TGTCGTTGTC      240
GCCTACGAGC GCAACGTACA GACCAACGCC CG                               272
```

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 317 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

```
GCAGCCGGTG GTTCTCGGAC TATCTGCGCA CGGTGACGCA GCGCGACGTG CGCGAGCTGA      60
AGCGGATCGA GCAGACGGAT CGCCTGCCGC GGTTCATGCG CTACCTGGCC GCTATCACCG      120
```

CGCAGGAGCT GAACGTGGCC GAAGCGGCGC GGGTCATCGG GGTCGACGCG GGGACGATCC	180
GTTCGGATCT GGCCTGGTTC GAGACGGTCT ATCTGGTACA TCGCCTGCCC GCCTGGTCGC	240
GGAATCTGAC CGCGAAGATC AAGAAGCGGT CAAAGATCCA CGTCGTCGAC AGTGGCTTCG	300
CGGCCTGGTT GCGCGGG	317

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 182 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GATCGTGGAG CTGTCGATGA ACAGCGTTGC CGGACGCGCG GCGGCCAGCA CGTCGGTGTA	60
GCAGCGCCGG ACCACCTCGC CGGTGGGCAG CATGGTGATG ACCACGTCGG CCTCGGCCAC	120
CGCTTCGGGC GCGCTACGAA ACACCGCGAC ACCGTGCGCG GCGGCGCCGG ACGCCGCCGT	180
GG	182

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 308 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GATCGCGAAG TTTGGTGAGC AGGTGGTCGA CGCGAAAGTC TGGGCGCCTG CGAAGCGGGT	60
CGGCGTTTAC GAGGCGAAGA CACGCCTGTC CGAGCTGCTG CGGCTCGTCT ACGGCGGGCA	120
GAGGTTGAGA TTGCCC GCCG CGGCGAGCCG GTAGCAAAGC TTGTGCCGCT GCATCCTCAT	180
GAGACTCGGC GGTTAGGCAT TGACCATGGC GTGTACCGCG TGCCCGACGA TTTGGACGCT	240
CCGTTGTCAG ACGACGTGCT CGAACGCTTT CACCGGTGAA GCGCTACCTC ATCGACACCC	300
ACGTTTGG	308

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 267 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

CCGACGACGA	GCAACTCACG	TGGATGATGG	TCGGCAGCGG	CATTGAGGAC	GGAGAGAATC	60
CGGCCGAAGC	TGCCGCGCGG	CAAGTGCTCA	TAGTGACCGG	CCGTAGAGGG	CTCCCCCGAT	120
GGCACC GGAC	TATTCTGGTG	TGCCGCTGGC	CGGTAAGAGC	GGGTAAAAGA	ATGTGAGGGG	180
ACACGATGAG	CAATCACACC	TACCGAGTGA	TCGAGATCGT	CGGGACCTCG	CCCGACGGCG	240
TCGACGCGGC	AATCCAGGGC	GGTCTGG				267

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1539 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

CTCGTGCCGA	AAGAATGTGA	GGGGACACGA	TGAGCAATCA	CACCTACCGA	GTGATCGAGA	60
TCGTGCGGAC	CTCGCCCGAC	GGCGTCGACG	CGGCAATCCA	GGGCGGTCTG	GCCCCGAGCTG	120
CGCAGACCAT	GCGCGCGCTG	GACTGGTTCG	AAGTACAGTC	AATTCGAGGC	CACCTGGTCG	180
ACGGAGCGGT	CGCGCACTTC	CAGGTGACTA	TGAAAGTCGG	CTTCCGCTGG	AGGATTCCTG	240
AACCTTCAAG	CGCGGCCGAT	AACTGAGGTG	CATCATTAAG	CGACTTTTCC	AGAACATCCT	300
GACGCGCTCG	AAACGCGGTT	CAGCCGACGG	TGGCTCCGCC	GAGGCGCTGC	CTCCAAAATC	360
CCTGCGACAA	TTCGTGCGCG	GCGCCTACAA	GGAAGTCGGT	GCTGAATTCG	TCGGGTATCT	420
GGTCGACCTG	TGTGGGCTGC	AGCCGGACGA	AGCGGTGCTC	GACGTCGGCT	GCGGCTCGGG	480
GCGGATGGCG	TTGCCGCTCA	CCGGCTATCT	GAACAGCGAG	GGACGCTACG	CCGGCTTCGA	540
TATCTCGCAG	AAAGCCATCG	CGTGGTGCCA	GGAGCACATC	ACCTCGGCGC	ACCCCAACTT	600
CCAGTTCGAG	GTCTCCGACA	TCTACAATC	GCTGTACAAC	CCGAAAGGGA	AATACCAGTC	660
ACTAGACTTT	CGCTTTCCAT	ATCCGGATGC	GTCGTTTCGAT	GTGGTGTTTC	TTACCTCGGT	720
GTTCACCCAC	ATGTTTCCGC	CGGACGTGGA	GCACTATCTG	GACGAGATCT	CCCGCGTGCT	780
GAAGCCCGGC	GGACGATGCC	TGTGCACGTA	CTTCTTGCTC	AATGACGAGT	CGTTAGCCCA	840
CATCGCGGAA	GGAAAGAGTG	CGCACAACTT	CCAGCATGAG	GGACCGGGTT	ATCGGACAAT	900

CCACAAGAAG	CGGCCCCGAAG	AAGCAATCGG	CTTGCCGGAG	ACCTTCGTCA	GGGATGTCTA	960
TGGCAAGTTC	GGCCTCGCCG	TGCACGAACC	ATTGCACTAC	GGCTCATGGA	GTGGCCGGGA	1020
ACCACGCCTA	AGCTTCCAGG	ACATCGTCAT	CGCGACCAAA	ACCGCGAGCT	AGGTCCGGCAT	1080
CCGGGAAGCA	TCGCGACACC	GTGGCGCCGA	GCGCCGCTGC	CGGCAGGCCG	ATTAGGCGGG	1140
CAGATTAGCC	CGCCGCGGCT	CCCGGCTCCG	AGTACGGCGC	CCCGAATGGC	GTCACCGGCT	1200
GGTAACCACG	CTTGCGCGCC	TGGGCGGCGG	CCTGCCGGAT	CAGGTGGTAG	ATGCCGACAA	1260
AGCCTGCGTG	ATCGGTCATC	ACCAACGGTG	ACAGCAGCCG	GTTGTGCACC	AGCGCGAACG	1320
CCACCCCGGT	CTCCGGGTCT	GTCCAGCCGA	TCGAGCCGCC	CAAGCCCACA	TGACCAAACC	1380
CCGGCATCAC	GTTGCCGATC	GGCATAACGT	GATAGCCAAG	ATGAAAATTT	AAGGGCACCA	1440
ATAGATTTTCG	ATCCGGCAGA	ACTTGCCGTC	GGTTGCGGGT	CAGGCCCGTG	ACCAGCTCCC	1500
GCGACAAGAA	CCGTATGCCG	TCGATCTCGC	CTCGTGCCG			1539

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 851 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

CTGCAGGGTG	GCGTGGATGA	GCGTCACCGC	GGGGCAGGCC	GAGCTGACCG	CCGCCCAGGT	60
CCGGGTTGCT	GCGGCGGCCT	ACGAGACGGC	GTATGGGCTG	ACGGTGCCCC	CGCCGGTGAT	120
CGCCGAGAAC	CGTGCTGAAC	TGATGATTCT	GATAGCGACC	AACCTCTTGG	GGCAAAACAC	180
CCCGGCGATC	GCGGTCAACG	AGGCCGAATA	CGGCGAGATG	TGGGCCCAAG	ACGCCGCCGC	240
GATGTTTGGC	TACGCCGCGG	CGACGGCGAC	GGCGACGGCG	ACGTTGCTGC	CGTTTCGAGGA	300
GGCGCCGGAG	ATGACCAGCG	CGGGTGGGCT	CCTCGAGCAG	GCCGCCGCGG	TCGAGGAGGC	360
CTCCGACACC	GCCGCGGCGA	ACCAGTTGAT	GAACAATGTG	CCCCAGGCGC	TGAAACAGTT	420
GGCCCAGCCC	ACGCAGGGCA	CCACGCCTTC	TTCCAAGCTG	GGTGGCCTGT	GGAAGACGGT	480
CTCGCCGCAT	CGGTGCGCCG	TCAGCAACAT	GGTGTCGATG	GCCAACAACC	ACATGTGCAT	540
GACCAACTCG	GGTGTGTCGA	TGACCAACAC	CTTGAGCTCG	ATGTTGAAGG	GCTTTGCTCC	600
GGCGGCGGCC	GCCCAGGCCG	TGCAAACCGC	GGCGCAAAAC	GGGGTCCGGG	CGATGAGCTC	660
GCTGGGCAGC	TCGCTGGGTT	CTTCGGGTCT	GGGCGGTGGG	GTGGCCGCCA	ACTTGGGTCTG	720
GGCGGCCTCG	GTACGGTATG	GTCACCGGGA	TGGCGGAAAA	TATGCANAGT	CTGGTCGGCG	780

GAACGGTGGT CCGGCGTAAG GTTTACCCCC GTTTTCTGGA TGCGGTGAAC TTCGTCAACG 840
GAAACAGTTA C 851

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 254 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GATCGATCGG GCGGAAATTT GGACCAGATT CGCCTCCGGC GATAACCCAA TCAATCGAAC 60
 CTAGATTTAT TCCGTCCAGG GGCCCGAGTA ATGGCTCGCA GGAGAGGAAC CTTACTGCTG 120
 CGGGCACCTG TCGTAGGTCC TCGATACGGC GGAAGGCGTC GACATTTTCC ACCGACACCC 180
 CCATCCAAAC GTTCGAGGGC CACTCCAGCT TGTGAGCGAG GCGACGCAGT CGCAGGCTGC 240
 GCTTGGTCAA GATC 254

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1227 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

GATCCTGACC GAAGCGGCCG CCGCCAAGGC GAAGTCGCTG TTGGACCAGG AGGGACGGGA 60
 CGATCTGGCG CTGCGGATCG CGGTTAGCC GGGGGGGTGC GCTGGATTGC GCTATAACCT 120
 TTTCTTCGAC GACCGGACGC TGGATGGTGA CCAAACCGCG GAGTTCGGTG GTGTCAGGTT 180
 GATCGTGGAC CGGATGAGCG CGCCGTATGT GGAAGGCGCG TCGATCGATT TCGTCGACAC 240
 TATTGAGAAG CAAGGTTTAC CATCGACAAT CCCAACGCCA CCGGCTCCTG CGCGTGCGGG 300
 GATTCGTTCA ACTGATAAAA CGCTAGTACG ACCCGCGGGT GCGCAACACG TACGAGCACA 360
 CCAAGACCTG ACCGCGCTGG AAAAGCAACT GAGCGATGCC TTGCACCTGA CCGCGTGGCG 420
 GGCCGCCGGC GGCAGGTGTC ACCTGCATGG TGAACAGCAC CTGGGCCTGA TATTGCGACC 480
 AGTACACGAT TTTGTGATC GAGGTCACCT CGACCTGGGA GAACTGCTTG CGGAACGCGT 540
 CGCTGCTCAG CTTGGCCAAG GCCTGATCGG AGCGCTTGTC GCGCACGCCG TCGTGGATAC 600

CGCACAGCGC ATTGCGAACG ATGGTGTCCA CATCGCGGTT CTCCAGCGCG TTGAGGTATC	660
CCTGAATCGC GGTTTTGGCC GGTCCCTCCG AGAATGTGCC TGCCGTGTTG GCTCCGTTGG	720
TGCGGACCCC GTATATGATC GCCGCCGTCA TAGCCGACAC CAGCGCGAGG GCTACCACAA	780
TGCCGATCAG CAGCCGCTTG TGCCGTCGCT TCGGGTAGGA CACCTGCGGC GGCACGCCGG	840
GATATGCGGC GGGCGGCAGC GCCGCGTCGT CTGCCGGTCC CGGGGCGAAG GCCGGTTCGG	900
CGGCGCCGAG GTCGTGGGGG TAGTCCAGGG CTTGGGGTTC GTGGGATGAG GGCTCGGGGT	960
ACGGCGCCGG TCCGTTGGTG CCGACACCGG GGTTCGGCGA GTGGGGACCG GGCATTGTGG	1020
TTCTCCTAGG GTGGTGGACG GGACCAGCTG CTAGGGCGAC AACCGCCCGT CGCGTCAGCC	1080
GGCAGCATCG GCAATCAGGT GAGCTCCCTA GGCAGGCTAG CGCAACAGCT GCCGTCAGCT	1140
CTCAACGCGA CGGGGCGGGC CGCGGCGCCG ATAATGTTGA AAGACTAGGC AACCTTAGGA	1200
ACGAAGGACG GAGATTTTGT GACGATC	1227

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 181 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

GCGGTGTCGG CGGATCCGGC GGGTGGTTGA ACGGCAACGG CGGGGCCGGC GGGGCCGGCG	60
GGACCGGCGC TAACGGTGGT GCCGGCGGCA ACGCCTGGTT GTTCGGGGCC GGCGGGTCCG	120
GCGGNGCCGG CACCAATGGT GGNGTCGGCG GGTCCGGCGG ATTTGTCTAC GGCAACGGCG	180
G	181

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 290 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GCGGTGTCGG CGGATCCGGC GGGTGGTTGA ACGGCAACGG CGGTGTCGGC GGCCGGGGCG	60
GCGACGGCGT CTTTGCCGGT GCCGGCGGCC AGGGCGGCCT CGGTGGGCAG GGCGGCAATG	120

GCGGCGGCTC CACCGGCGGC AACGGCGGTC TTGGCGGCGC GGGCGGTGGC GGAGGCAACG 180
CCCCGGACGG CGGCTTCGGT GGCAACGGCG GTAAGGGTGG CCAGGGCGGN ATTGGCGGCG 240
GCACTCAGAG CGCGACCGGC CTCGGNGGTG ACGGCGGTGA CGGCGGTGAC 290

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

GATCCAGTGG CATGGNGGGT GTCAGTGGAA GCAT 34

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 155 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

GATCGCTGCT CGTCCCCCCC TTGCCGCCGA CGCCACCGGT CCCACCGTTA CCGAACAAGC 60
TGGCGTGGTC GCCAGCACCC CCGGCACCGC CGACGCCGGA GTCGAACAAT GGCACCGTCG 120
TATCCCCACC ATTGCCGCCG GNCCCACCGG CACCG 155

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 53 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

ATGGCGTTCA CGGGGCGCCG GGGACCGGGC AGCCCGGNGG GGCCGGGGGG TGG 53

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 132 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

GATCCACCGC GGGTGCAGAC GGTGCCCCGCG GCGCCACCCC GACCAGCGGC GGCAACGGCG	60
GCACCGGCGG CAACGGCGCG AACGCCACCG TCGTCGGNGG GGCCGGCGGG GCCGGCGGCA	120
AGGGCGGCAA CG	132

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 132 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GATCGGCGGC CGGNACGGNC GGGGACGGCG GCAAGGGCGG NAACGGGGGC GCCGNAGCCA	60
CCNGCCAAGA ATCCTCCGNG TCCNCCAATG GCGCGAATGG CGGACAGGGC GGCAACGGCG	120
GCANCGGCGG CA	132

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 702 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

CGGCACGAGG ATCGGTACCC CGCGGCATCG GCAGCTGCCG ATTCGCCGGG TTTCCCCACC	60
CGAGGAAAGC CGCTACCAGA TGGCGCTGCC GAAGTAGGGC GATCCGTTCG CGATGCCGGC	120
ATGAACGGGC GGCATCAAAT TAGTGCAGGA ACCTTTCAGT TTAGCGACGA TAATGGCTAT	180
AGCACTAAGG AGGATGATCC GATATGACGC AGTCGCAGAC CGTGACGGTG GATCAGCAAG	240
AGATTTTGAA CAGGGCCAAC GAGGTGGAGG CCCCAGTGGC GGACCCACCG ACTGATGTCC	300
CCATCACACC GTGCGAACTC ACGGNGGNTA AAAACGCCGC CCAACAGNTG GTNTTGTCCG	360

CCGACAACAT GCGGGAATAC CTGGCGGCCG GTGCCAAAGA GCGGCAGCGT CTGGCGACCT	420
CGCTGCGCAA CGCGGCCAAG GNGTATGGCG AGGTTGATGA GGAGGCTGCG ACCGCGCTGG	480
ACAACGACGG CGAAGGAACT GTGCAGGCAG AATCGGCCGG GGCCGTCGGA GGGGACAGTT	540
CGGCCGAACT AACCGATACG CCGAGGGTGG CCACGGCCGG TGAACCCAAC TTCATGGATC	600
TCAAAGAAGC GGCAAGGAAG CTCGAAACGG GCGACCAAGG CGCATCGCTC GCGCACTGNG	660
GGGATGGGTG GAACACTTNC ACCCTGACGC TGCAAGGCGA CG	702

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 298 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

GAAGCCGCAG CGCTGTCGGG CGACGTGGCG GTCAAAGCGG CATCGCTCGG TGGCGGTGGA	60
GGCGGCGGGG TGCCGTCGGC GCCGTTGGGA TCCGCGATCG GGGGCGCCGA ATCGGTGCGG	120
CCCGCTGGCG CTGGTGACAT TGCCGGCTTA GGCCAGGGAA GGGCCGGCGG CGGCGCCGCG	180
CTGGGCGGCG GTGGCATGGG AATGCCGATG GGTGCCGCGC ATCAGGGACA AGGGGGCGCC	240
AAGTCCAAGG GTTCTCAGCA GGAAGACGAG GCGCTCTACA CCGAGGATCC TCGTGCCG	298

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1058 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

CGGCACGAGG ATCGAATCGC GTCGCCGGGA GCACAGCGTC GCACTGCACC AGTGGAGGAG	60
CCATGACCTA CTCGCCGGGT AACCCCGGAT ACCCGCAAGC GCAGCCCGCA GGCTCCTACG	120
GAGGCGTCAC ACCCTCGTTC GCCCAGCCG ATGAGGGTGC GAGCAAGCTA CCGATGTACC	180
TGAACATCGC GGTGGCAGTG CTCGGTCTGG CTGCGTACTT CGCCAGCTTC GGCCCAATGT	240
TCACCCTCAG TACCGAACTC GGGGGGGGTG ATGGCGCAGT GTCCGGTGAC ACTGGGCTGC	300
CGGTGCGGGT GGCTCTGCTG GCTGCGCTGC TTGCCGGGGT GGTCTGGTG CCTAAGGCCA	360

AGAGCCATGT GACGGTAGTT GCGGTGCTCG GGGTACTCGG CGTATTTCTG ATGGTCTCGG	420
CGACGTTTAA CAAGCCCAGC GCCTATTCGA CCGGTTGGGC ATTGTGGGTT GTGTTGGCTT	480
TCATCGTGTT CCAGGCGGTT GCGGCAGTCC TGGCGCTCTT GGTGGAGACC GGCGCTATCA	540
CCGCGCCGGC GCCGCGGCCC AAGTTCGACC CGTATGGACA GTACGGGCGG TACGGGCAGT	600
ACGGGCAGTA CGGGGTGCAG CCGGGTGGGT ACTACGGTCA GCAGGGTGCT CAGCAGGCCG	660
CGGGACTGCA GTCGCCCCGC CCGCAGCAGT CTCCGCAGCC TCCCGGATAT GGGTCGCAGT	720
ACGGCGGCTA TTCGTCCAGT CCGAGCCAAT CGGGCAGTGG ATACACTGCT CAGCCCCCGG	780
CCCAGCCGCC GGCGCAGTCC GGGTCGCAAC AATCGCACCA GGGCCCATCC ACGCCACCTA	840
CCGGCTTTCC GAGCTTCAGC CCACCACCAC CGGTCAGTGC CGGGACGGGG TCGCAGGCTG	900
GTTCGGCTCC AGTCAACTAT TCAAACCCCA GCGGGGGCGA GCAGTCGTCG TCCCCCGGGG	960
GGGCGCCGGT CTAACCGGGC GTTCCCGCGT CCGGTCGCGC GTGTGCGCGA AGAGTGAACA	1020
GGGTGTCAGC AAGCGCGGAC GATCCTCGTG CCGAATTC	1058

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 327 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

CGGCACGAGA GACCGATGCC GCTACCCTCG CGCAGGAGGC AGGTAATTTT GAGCGGATCT	60
CCGGCGACCT GAAAACCCAG ATCGACCAGG TGGAGTCGAC GGCAGGTTCG TTGCAGGGCC	120
AGTGCGCGCG CGCGGCGGGG ACGGCCGCCC AGGCCGCGGT GGTGCGCTTC CAAGAAGCAG	180
CCAATAAGCA GAAGCAGGAA CTCGACGAGA TCTCGACGAA TATTCGTCAG GCCGGCGTCC	240
AATACTCGAG GGCCGACGAG GAGCAGCAGC AGGCGCTGTC CTCGCAAATG GGCTTCTGAC	300
CCGCTAATAC GAAAAGAAAC GGAGCAA	327

(2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 170 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

CGGTCGCGAT GATGGCGTTG TCGAACGTGA CCGATTCTGT ACCGCCGTCG TTGAGATCAA 60
CCAACAACGT GTTGGCGTCG GCAAATGTGC CGNACCCGTG GATCTCGGTG ATCTTGTTCT 120
TCTTCATCAG GAAGTGCACA CCGGCCACCC TGCCCTCGGN TACCTTTTCGG 170

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 127 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GATCCGGCGG CACGGGGGGT GCCGGCGGCA GCACCGCTGG CGCTGGCGGC AACGGCGGGG 60
CCGGGGGTGG CGGCGGAACC GGTGGGTTGC TCTTCGGCAA CGGCGGTGCC GGCGGGCACG 120
GGGCCGT 127

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 81 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

CGGCGGCAAG GGCGGCACCG CCGGCAACGG GAGCGGCGCG GCCGGCGGCA ACGGCGGCAA 60
CGGCGGCTCC GGCCTCAACG G 81

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 149 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

GATCAGGGCT GGCCGGCTCC GGCCAGAAGG GCGGTAACGG AGGAGCTGCC GGATTGTTTG 60

GCAACGGCGG GGCCGGNGGT GCCGGCGCGT CCAACCAAGC CGGTAACGGC GGNGCCGGCG 120
 GAAACGGTGG TGCCGGTGGG CTGATCTGG 149

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 355 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CGGCACGAGA TCACACCTAC CGAGTGATCG AGATCGTCGG GACCTCGCCC GACGGTGTCG 60
 ACGCGGNAAT CCAGGGCGGT CTGGCCCGAG CTGCGCAGAC CATGCGCGCG CTGGACTGGT 120
 TCGAAGTACA GTCAATTCGA GGCCACCTGG TCGACGGAGC GGTGCGGCAC TTCCAGGTGA 180
 CTATGAAAGT CGGCTTCCGC CTGGAGGATT CCTGAACCTT CAAGCGCGGC CGATAACTGA 240
 GGTGCATCAT TAAGCGACTT TTCCAGAACA TCCTGACGCG CTCGAAACGC GTTTCAGCCG 300
 ACGGTGGCTC CGCCGAGGCG CTGCCTCCAA AATCCCTGCG ACAATTCGTC GGCGG 355

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 999 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

ATGCATCACC ATCACCATCA CATGCATCAG GTGGACCCCA ACTTGACACG TCGCAAGGGA 60
 CGATTGGCGG CACTGGCTAT CGCGGCGATG GCCAGCGCCA GCCTGGTGAC CGTTGCGGTG 120
 CCCGCGACCG CCAACGCCGA TCCGGAGCCA GCGCCCCCGG TACCCACAAC GGCCGCCTCG 180
 CCGCCGTCGA CCGCTGCAGC GCCACCCGCA CCGGCGACAC CTGTTGCCCC CCCACCACCG 240
 GCCGCCGCCA ACACGCCGAA TGCCAGCCG GCGGATCCCA ACGCAGCACC TCCGCCGGCC 300
 GACCCGAACG CACCGCCGCC ACCTGTCATT GCCCAAACG CACCCCAACC TGTCCGGATC 360
 GACAACCCGG TTGGAGGATT CAGCTTCGCG CTGCCTGCTG GCTGGGTGGA GTCTGACGCC 420
 GCCCACTTCG ACTACGGTTC AGCACTCCTC AGCAAAACCA CCGGGGACCC GCCATTTCCC 480
 GGACAGCCGC CGCCGGTGGC CAATGACACC CGTATCGTGC TCGGCCGGCT AGACCAAAAAG 540

CTTTACGCCA GCGCCGAAGC CACCGACTCC AAGGCCGCGG CCCGGTTGGG CTCGGACATG 600
 GGTGAGTTCT ATATGCCCTA CCCGGGCACC CGGATCAACC AGGAAACCGT CTCGCTCGAC 660
 GCCAACGGGG TGTCTGGAAG CGCGTCGTAT TACGAAGTCA AGTTCAGCGA TCCGAGTAAG 720
 CCGAACGGCC AGATCTGGAC GGGCGTAATC GGCTCGCCCG CGGCGAACGC ACCGGACGCC 780
 GGGCCCCCTC AGCGCTGGTT TGTGGTATGG CTCGGGACCG CCAACAACCC GGTGGACAAG 840
 GCGCGGGCCA AGGCGCTGGC CGAATCGATC CGGCCTTTGG TCGCCCCGCC GCCGGCGCCG 900
 GCACCGGCTC CTGCAGAGCC CGCTCCGGCG CCGGCGCCCG CCGGGGAAGT CGCTCCTACC 960
 CCGACGACAC CGACACCGCA GCGGACCTTA CCGGCCTGA 999

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 332 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

Met His His His His His His Met His Gln Val Asp Pro Asn Leu Thr
 1 5 10 15
 Arg Arg Lys Gly Arg Leu Ala Ala Leu Ala Ile Ala Ala Met Ala Ser
 20 25 30
 Ala Ser Leu Val Thr Val Ala Val Pro Ala Thr Ala Asn Ala Asp Pro
 35 40 45
 Glu Pro Ala Pro Pro Val Pro Thr Thr Ala Ala Ser Pro Pro Ser Thr
 50 55 60
 Ala Ala Ala Pro Pro Ala Pro Ala Thr Pro Val Ala Pro Pro Pro Pro
 65 70 75 80
 Ala Ala Ala Asn Thr Pro Asn Ala Gln Pro Gly Asp Pro Asn Ala Ala
 85 90 95
 Pro Pro Pro Ala Asp Pro Asn Ala Pro Pro Pro Pro Val Ile Ala Pro
 100 105 110
 Asn Ala Pro Gln Pro Val Arg Ile Asp Asn Pro Val Gly Gly Phe Ser
 115 120 125
 Phe Ala Leu Pro Ala Gly Trp Val Glu Ser Asp Ala Ala His Phe Asp
 130 135 140
 Tyr Gly Ser Ala Leu Leu Ser Lys Thr Thr Gly Asp Pro Pro Phe Pro
 145 150 155 160
 Gly Gln Pro Pro Pro Val Ala Asn Asp Thr Arg Ile Val Leu Gly Arg
 165 170 175

Leu Asp Gln Lys Leu Tyr Ala Ser Ala Glu Ala Thr Asp Ser Lys Ala
 180 185 190
 Ala Ala Arg Leu Gly Ser Asp Met Gly Glu Phe Tyr Met Pro Tyr Pro
 195 200 205
 Gly Thr Arg Ile Asn Gln Glu Thr Val Ser Leu Asp Ala Asn Gly Val
 210 215 220
 Ser Gly Ser Ala Ser Tyr Tyr Glu Val Lys Phe Ser Asp Pro Ser Lys
 225 230 235 240
 Pro Asn Gly Gln Ile Trp Thr Gly Val Ile Gly Ser Pro Ala Ala Asn
 245 250 255
 Ala Pro Asp Ala Gly Pro Pro Gln Arg Trp Phe Val Val Trp Leu Gly
 260 265 270
 Thr Ala Asn Asn Pro Val Asp Lys Gly Ala Ala Lys Ala Leu Ala Glu
 275 280 285
 Ser Ile Arg Pro Leu Val Ala Pro Pro Pro Ala Pro Ala Pro Ala Pro
 290 295 300
 Ala Glu Pro Ala Pro Ala Pro Ala Pro Ala Gly Glu Val Ala Pro Thr
 305 310 315 320
 Pro Thr Thr Pro Thr Pro Gln Arg Thr Leu Pro Ala
 325 330

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

Asp Pro Val Asp Ala Val Ile Asn Thr Thr Xaa Asn Tyr Gly Gln Val
 1 5 10 15
 Val Ala Ala Leu
 20

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 15 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Ala Val Glu Ser Gly Met Leu Ala Leu Gly Thr Pro Ala Pro Ser
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

Ala Ala Met Lys Pro Arg Thr Gly Asp Gly Pro Leu Glu Ala Ala Lys
1 5 10 15

Glu Gly Arg

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Tyr Tyr Trp Cys Pro Gly Gln Pro Phe Asp Pro Ala Trp Gly Pro
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Asp Ile Gly Ser Glu Ser Thr Glu Asp Gln Gln Xaa Ala Val
1 5 10

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

Ala Glu Glu Ser Ile Ser Thr Xaa Glu Xaa Ile Val Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:60:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

Asp Pro Glu Pro Ala Pro Pro Val Pro Thr Ala Ala Ala Ala Pro Pro
1 5 10 15
Ala

(2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Ala Pro Lys Thr Tyr Xaa Glu Glu Leu Lys Gly Thr Asp Thr Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:62:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

Asp Pro Ala Ser Ala Pro Asp Val Pro Thr Ala Ala Gln Gln Thr Ser
 1 5 10 15
 Leu Leu Asn Asn Leu Ala Asp Pro Asp Val Ser Phe Ala Asp
 20 25 30

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 187 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

Thr Gly Ser Leu Asn Gln Thr His Asn Arg Arg Ala Asn Glu Arg Lys
 1 5 10 15
 Asn Thr Thr Met Lys Met Val Lys Ser Ile Ala Ala Gly Leu Thr Ala
 20 25 30
 Ala Ala Ala Ile Gly Ala Ala Ala Ala Gly Val Thr Ser Ile Met Ala
 35 40 45
 Gly Gly Pro Val Val Tyr Gln Met Gln Pro Val Val Phe Gly Ala Pro
 50 55 60
 Leu Pro Leu Asp Pro Ala Ser Ala Pro Asp Val Pro Thr Ala Ala Gln
 65 70 75 80
 Leu Thr Ser Leu Leu Asn Ser Leu Ala Asp Pro Asn Val Ser Phe Ala
 85 90 95
 Asn Lys Gly Ser Leu Val Glu Gly Gly Ile Gly Gly Thr Glu Ala Arg
 100 105 110
 Ile Ala Asp His Lys Leu Lys Lys Ala Ala Glu His Gly Asp Leu Pro
 115 120 125
 Leu Ser Phe Ser Val Thr Asn Ile Gln Pro Ala Ala Ala Gly Ser Ala
 130 135 140
 Thr Ala Asp Val Ser Val Ser Gly Pro Lys Leu Ser Ser Pro Val Thr
 145 150 155 160
 Gln Asn Val Thr Phe Val Asn Gln Gly Gly Trp Met Leu Ser Arg Ala
 165 170 175
 Ser Ala Met Glu Leu Leu Gln Ala Ala Gly Xaa
 180 185

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 148 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

```

Asp Glu Val Thr Val Glu Thr Thr Ser Val Phe Arg Ala Asp Phe Leu
1           5           10           15
Ser Glu Leu Asp Ala Pro Ala Gln Ala Gly Thr Glu Ser Ala Val Ser
20           25           30
Gly Val Glu Gly Leu Pro Pro Gly Ser Ala Leu Leu Val Val Lys Arg
35           40           45
Gly Pro Asn Ala Gly Ser Arg Phe Leu Leu Asp Gln Ala Ile Thr Ser
50           55           60
Ala Gly Arg His Pro Asp Ser Asp Ile Phe Leu Asp Asp Val Thr Val
65           70           75           80
Ser Arg Arg His Ala Glu Phe Arg Leu Glu Asn Asn Glu Phe Asn Val
85           90           95
Val Asp Val Gly Ser Leu Asn Gly Thr Tyr Val Asn Arg Glu Pro Val
100          105          110
Asp Ser Ala Val Leu Ala Asn Gly Asp Glu Val Gln Ile Gly Lys Leu
115          120          125
Arg Leu Val Phe Leu Thr Gly Pro Lys Gln Gly Glu Asp Asp Gly Ser
130          135          140
Thr Gly Gly Pro
145

```

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 230 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

```

Thr Ser Asn Arg Pro Ala Arg Arg Gly Arg Arg Ala Pro Arg Asp Thr
1           5           10           15
Gly Pro Asp Arg Ser Ala Ser Leu Ser Leu Val Arg His Arg Arg Gln
20           25           30

```

Gln Arg Asp Ala Leu Cys Leu Ser Ser Thr Gln Ile Ser Arg Gln Ser
 35 40 45
 Asn Leu Pro Pro Ala Ala Gly Gly Ala Ala Asn Tyr Ser Arg Arg Asn
 50 55 60
 Phe Asp Val Arg Ile Lys Ile Phe Met Leu Val Thr Ala Val Val Leu
 65 70 75 80
 Leu Cys Cys Ser Gly Val Ala Thr Ala Ala Pro Lys Thr Tyr Cys Glu
 85 90 95
 Glu Leu Lys Gly Thr Asp Thr Gly Gln Ala Cys Gln Ile Gln Met Ser
 100 105 110
 Asp Pro Ala Tyr Asn Ile Asn Ile Ser Leu Pro Ser Tyr Tyr Pro Asp
 115 120 125
 Gln Lys Ser Leu Glu Asn Tyr Ile Ala Gln Thr Arg Asp Lys Phe Leu
 130 135 140
 Ser Ala Ala Thr Ser Ser Thr Pro Arg Glu Ala Pro Tyr Glu Leu Asn
 145 150 155 160
 Ile Thr Ser Ala Thr Tyr Gln Ser Ala Ile Pro Pro Arg Gly Thr Gln
 165 170 175
 Ala Val Val Leu Xaa Val Tyr His Asn Ala Gly Gly Thr His Pro Thr
 180 185 190
 Thr Thr Tyr Lys Ala Phe Asp Trp Asp Gln Ala Tyr Arg Lys Pro Ile
 195 200 205
 Thr Tyr Asp Thr Leu Trp Gln Ala Asp Thr Asp Pro Leu Pro Val Val
 210 215 220
 Phe Pro Ile Val Ala Arg
 225 230

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 132 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly Phe
 1 5 10 15
 Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg Ser
 20 25 30
 Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu Gly
 35 40 45

98

Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg Val
 50 55 60
 Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp Val
 65 70 75 80
 Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met Ala
 85 90 95
 Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Asn Trp
 100 105 110
 Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala Glu
 115 120 125
 Gly Pro Pro Ala
 130

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 100 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

Val Pro Leu Arg Ser Pro Ser Met Ser Pro Ser Lys Cys Leu Ala Ala
 1 5 10 15
 Ala Gln Arg Asn Pro Val Ile Arg Arg Arg Arg Leu Ser Asn Pro Pro
 20 25 30
 Pro Arg Lys Tyr Arg Ser Met Pro Ser Pro Ala Thr Ala Ser Ala Gly
 35 40 45
 Met Ala Arg Val Arg Arg Arg Ala Ile Trp Arg Gly Pro Ala Thr Xaa
 50 55 60
 Ser Ala Gly Met Ala Arg Val Arg Arg Trp Xaa Val Met Pro Xaa Val
 65 70 75 80
 Ile Gln Ser Thr Xaa Ile Arg Xaa Xaa Gly Pro Phe Asp Asn Arg Gly
 85 90 95
 Ser Glu Arg Lys
 100

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 163 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

```

Met Thr Asp Asp Ile Leu Leu Ile Asp Thr Asp Glu Arg Val Arg Thr
1      5      10      15
Leu Thr Leu Asn Arg Pro Gln Ser Arg Asn Ala Leu Ser Ala Ala Leu
20     25     30
Arg Asp Arg Phe Phe Ala Xaa Leu Xaa Asp Ala Glu Xaa Asp Asp Asp
35     40     45
Ile Asp Val Val Ile Leu Thr Gly Ala Asp Pro Val Phe Cys Ala Gly
50     55     60
Leu Asp Leu Lys Val Ala Gly Arg Ala Asp Arg Ala Ala Gly His Leu
65     70     75     80
Thr Ala Val Gly Gly His Asp Gln Ala Gly Asp Arg Arg Asp Gln Arg
85     90     95
Arg Arg Gly His Arg Arg Ala Arg Thr Gly Ala Val Leu Arg His Pro
100    105    110
Asp Arg Leu Arg Ala Arg Pro Leu Arg Arg His Pro Arg Pro Gly Gly
115    120    125
Ala Ala Ala His Leu Gly Thr Gln Cys Val Leu Ala Ala Lys Gly Arg
130    135    140
His Arg Xaa Gly Pro Val Asp Glu Pro Asp Arg Arg Leu Pro Val Arg
145    150    155    160
Asp Arg Arg

```

(2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 344 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

```

Met Lys Phe Val Asn His Ile Glu Pro Val Ala Pro Arg Arg Ala Gly
1      5      10      15
Gly Ala Val Ala Glu Val Tyr Ala Glu Ala Arg Arg Glu Phe Gly Arg
20     25     30
Leu Pro Glu Pro Leu Ala Met Leu Ser Pro Asp Glu Gly Leu Leu Thr
35     40     45

```

100

Ala Gly Trp Ala Thr Leu Arg Glu Thr Leu Leu Val Gly Gln Val Pro
 50 55 60
 Arg Gly Arg Lys Glu Ala Val Ala Ala Ala Val Ala Ala Ser Leu Arg
 65 70 75 80
 Cys Pro Trp Cys Val Asp Ala His Thr Thr Met Leu Tyr Ala Ala Gly
 85 90 95
 Gln Thr Asp Thr Ala Ala Ala Ile Leu Ala Gly Thr Ala Pro Ala Ala
 100 105 110
 Gly Asp Pro Asn Ala Pro Tyr Val Ala Trp Ala Ala Gly Thr Gly Thr
 115 120 125
 Pro Ala Gly Pro Pro Ala Pro Phe Gly Pro Asp Val Ala Ala Glu Tyr
 130 135 140
 Leu Gly Thr Ala Val Gln Phe His Phe Ile Ala Arg Leu Val Leu Val
 145 150 155 160
 Leu Leu Asp Glu Thr Phe Leu Pro Gly Gly Pro Arg Ala Gln Gln Leu
 165 170 175
 Met Arg Arg Ala Gly Gly Leu Val Phe Ala Arg Lys Val Arg Ala Glu
 180 185 190
 His Arg Pro Gly Arg Ser Thr Arg Arg Leu Glu Pro Arg Thr Leu Pro
 195 200 205
 Asp Asp Leu Ala Trp Ala Thr Pro Ser Glu Pro Ile Ala Thr Ala Phe
 210 215 220
 Ala Ala Leu Ser His His Leu Asp Thr Ala Pro His Leu Pro Pro Pro
 225 230 235 240
 Thr Arg Gln Val Val Arg Arg Val Val Gly Ser Trp His Gly Glu Pro
 245 250 255
 Met Pro Met Ser Ser Arg Trp Thr Asn Glu His Thr Ala Glu Leu Pro
 260 265 270
 Ala Asp Leu His Ala Pro Thr Arg Leu Ala Leu Leu Thr Gly Leu Ala
 275 280 285
 Pro His Gln Val Thr Asp Asp Asp Val Ala Ala Ala Arg Ser Leu Leu
 290 295 300
 Asp Thr Asp Ala Ala Leu Val Gly Ala Leu Ala Trp Ala Ala Phe Thr
 305 310 315 320
 Ala Ala Arg Arg Ile Gly Thr Trp Ile Gly Ala Ala Ala Glu Gly Gln
 325 330 335
 Val Ser Arg Gln Asn Pro Thr Gly
 340

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 485 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

Asp	Asp	Pro	Asp	Met	Pro	Gly	Thr	Val	Ala	Lys	Ala	Val	Ala	Asp	Ala
1				5					10					15	
Leu	Gly	Arg	Gly	Ile	Ala	Pro	Val	Glu	Asp	Ile	Gln	Asp	Cys	Val	Glu
			20					25					30		
Ala	Arg	Leu	Gly	Glu	Ala	Gly	Leu	Asp	Asp	Val	Ala	Arg	Val	Tyr	Ile
		35					40					45			
Ile	Tyr	Arg	Gln	Arg	Arg	Ala	Glu	Leu	Arg	Thr	Ala	Lys	Ala	Leu	Leu
	50					55					60				
Gly	Val	Arg	Asp	Glu	Leu	Lys	Leu	Ser	Leu	Ala	Ala	Val	Thr	Val	Leu
65					70					75					80
Arg	Glu	Arg	Tyr	Leu	Leu	His	Asp	Glu	Gln	Gly	Arg	Pro	Ala	Glu	Ser
				85					90					95	
Thr	Gly	Glu	Leu	Met	Asp	Arg	Ser	Ala	Arg	Cys	Val	Ala	Ala	Ala	Glu
			100					105					110		
Asp	Gln	Tyr	Glu	Pro	Gly	Ser	Ser	Arg	Arg	Trp	Ala	Glu	Arg	Phe	Ala
		115					120					125			
Thr	Leu	Leu	Arg	Asn	Leu	Glu	Phe	Leu	Pro	Asn	Ser	Pro	Thr	Leu	Met
	130					135					140				
Asn	Ser	Gly	Thr	Asp	Leu	Gly	Leu	Leu	Ala	Gly	Cys	Phe	Val	Leu	Pro
145					150					155					160
Ile	Glu	Asp	Ser	Leu	Gln	Ser	Ile	Phe	Ala	Thr	Leu	Gly	Gln	Ala	Ala
				165					170					175	
Glu	Leu	Gln	Arg	Ala	Gly	Gly	Gly	Thr	Gly	Tyr	Ala	Phe	Ser	His	Leu
			180					185					190		
Arg	Pro	Ala	Gly	Asp	Arg	Val	Ala	Ser	Thr	Gly	Gly	Thr	Ala	Ser	Gly
		195					200					205			
Pro	Val	Ser	Phe	Leu	Arg	Leu	Tyr	Asp	Ser	Ala	Ala	Gly	Val	Val	Ser
	210					215						220			
Met	Gly	Gly	Arg	Arg	Arg	Gly	Ala	Cys	Met	Ala	Val	Leu	Asp	Val	Ser
225					230					235					240
His	Pro	Asp	Ile	Cys	Asp	Phe	Val	Thr	Ala	Lys	Ala	Glu	Ser	Pro	Ser
				245					250					255	
Glu	Leu	Pro	His	Phe	Asn	Leu	Ser	Val	Gly	Val	Thr	Asp	Ala	Phe	Leu
			260					265					270		
Arg	Ala	Val	Glu	Arg	Asn	Gly	Leu	His	Arg	Leu	Val	Asn	Pro	Arg	Thr
		275					280					285			

Gly Lys Ile Val Ala Arg Met Pro Ala Ala Glu Leu Phe Asp Ala Ile
 290 295 300
 Cys Lys Ala Ala His Ala Gly Gly Asp Pro Gly Leu Val Phe Leu Asp
 305 310 315 320
 Thr Ile Asn Arg Ala Asn Pro Val Pro Gly Arg Gly Arg Ile Glu Ala
 325 330 335
 Thr Asn Pro Cys Gly Glu Val Pro Leu Leu Pro Tyr Glu Ser Cys Asn
 340 345 350
 Leu Gly Ser Ile Asn Leu Ala Arg Met Leu Ala Asp Gly Arg Val Asp
 355 360 365
 Trp Asp Arg Leu Glu Glu Val Ala Gly Val Ala Val Arg Phe Leu Asp
 370 375 380
 Asp Val Ile Asp Val Ser Arg Tyr Pro Phe Pro Glu Leu Gly Glu Ala
 385 390 395 400
 Ala Arg Ala Thr Arg Lys Ile Gly Leu Gly Val Met Gly Leu Ala Glu
 405 410 415
 Leu Leu Ala Ala Leu Gly Ile Pro Tyr Asp Ser Glu Glu Ala Val Arg
 420 425 430
 Leu Ala Thr Arg Leu Met Arg Arg Ile Gln Gln Ala Ala His Thr Ala
 435 440 445
 Ser Arg Arg Leu Ala Glu Glu Arg Gly Ala Phe Pro Ala Phe Thr Asp
 450 455 460
 Ser Arg Phe Ala Arg Ser Gly Pro Arg Arg Asn Ala Gln Val Thr Ser
 465 470 475 480
 Val Ala Pro Thr Gly
 485

(2) INFORMATION FOR SEQ ID NO:71:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 267 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

Gly Val Ile Val Leu Asp Leu Glu Pro Arg Gly Pro Leu Pro Thr Glu
 1 5 10 15
 Ile Tyr Trp Arg Arg Arg Gly Leu Ala Leu Gly Ile Ala Val Val Val
 20 25 30
 Val Gly Ile Ala Val Ala Ile Val Ile Ala Phe Val Asp Ser Ser Ala
 35 40 45

103

Gly Ala Lys Pro Val Ser Ala Asp Lys Pro Ala Ser Ala Gln Ser His
 50 55 60
 Pro Gly Ser Pro Ala Pro Gln Ala Pro Gln Pro Ala Gly Gln Thr Glu
 65 70 75 80
 Gly Asn Ala Ala Ala Pro Pro Gln Gly Gln Asn Pro Glu Thr Pro
 85 90 95
 Thr Pro Thr Ala Ala Val Gln Pro Pro Pro Val Leu Lys Glu Gly Asp
 100 105 110
 Asp Cys Pro Asp Ser Thr Leu Ala Val Lys Gly Leu Thr Asn Ala Pro
 115 120 125
 Gln Tyr Tyr Val Gly Asp Gln Pro Lys Phe Thr Met Val Val Thr Asn
 130 135 140
 Ile Gly Leu Val Ser Cys Lys Arg Asp Val Gly Ala Ala Val Leu Ala
 145 150 155 160
 Ala Tyr Val Tyr Ser Leu Asp Asn Lys Arg Leu Trp Ser Asn Leu Asp
 165 170 175
 Cys Ala Pro Ser Asn Glu Thr Leu Val Lys Thr Phe Ser Pro Gly Glu
 180 185 190
 Gln Val Thr Thr Ala Val Thr Trp Thr Gly Met Gly Ser Ala Pro Arg
 195 200 205
 Cys Pro Leu Pro Arg Pro Ala Ile Gly Pro Gly Thr Tyr Asn Leu Val
 210 215 220
 Val Gln Leu Gly Asn Leu Arg Ser Leu Pro Val Pro Phe Ile Leu Asn
 225 230 235 240
 Gln Pro Pro Pro Pro Pro Gly Pro Val Pro Ala Pro Gly Pro Ala Gln
 245 250 255
 Ala Pro Pro Pro Glu Ser Pro Ala Gln Gly Gly
 260 265

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

Leu Ile Ser Thr Gly Lys Ala Ser His Ala Ser Leu Gly Val Gln Val
 1 5 10 15
 Thr Asn Asp Lys Asp Thr Pro Gly Ala Lys Ile Val Glu Val Val Ala
 20 25 30

[illegible]

(2) INFORMATION FOR SEQ ID NO:73:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 364 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

Gly 1	Ala	Ala	Val	Ser 5	Leu	Leu	Ala	Ala	Gly 10	Thr	Leu	Val	Leu	Thr 15	Ala
Cys	Gly	Gly	Gly 20	Thr	Asn	Ser	Ser	Ser 25	Ser	Gly	Ala	Gly	Gly 30	Thr	Ser
Gly	Ser	Val 35	His	Cys	Gly	Gly	Lys 40	Lys	Glu	Leu	His	Ser 45	Ser	Gly	Ser
Thr	Ala 50	Gln	Glu	Asn	Ala	Met 55	Glu	Gln	Phe	Val	Tyr 60	Ala	Tyr	Val	Arg
Ser 65	Cys	Pro	Gly	Tyr	Thr 70	Leu	Asp	Tyr	Asn	Ala 75	Asn	Gly	Ser	Gly	Ala 80
Gly	Val	Thr	Gln	Phe 85	Leu	Asn	Asn	Glu	Thr 90	Asp	Phe	Ala	Gly	Ser 95	Asp
Val	Pro	Leu	Asn 100	Pro	Ser	Thr	Gly	Gln 105	Pro	Asp	Arg	Ser	Ala 110	Glu	Arg
Cys	Gly	Ser 115	Pro	Ala	Trp	Asp	Leu 120	Pro	Thr	Val	Phe	Gly 125	Pro	Ile	Ala
Ile	Thr 130	Tyr	Asn	Ile	Lys	Gly 135	Val	Ser	Thr	Leu	Asn 140	Leu	Asp	Gly	Pro
Thr 145	Thr	Ala	Lys	Ile	Phe 150	Asn	Gly	Thr	Ile	Thr 155	Val	Trp	Asn	Asp	Pro 160
Gln	Ile	Gln	Ala	Leu 165	Asn	Ser	Gly	Thr	Asp 170	Leu	Pro	Pro	Thr	Pro 175	Ile

105

Ser Val Ile Phe Arg Ser Asp Lys Ser Gly Thr Ser Asp Asn Phe Gln
 180 185 190
 Lys Tyr Leu Asp Gly Val Ser Asn Gly Ala Trp Gly Lys Gly Ala Ser
 195 200 205
 Glu Thr Phe Ser Gly Gly Val Gly Val Gly Ala Ser Gly Asn Asn Gly
 210 215 220
 Thr Ser Ala Leu Leu Gln Thr Thr Asp Gly Ser Ile Thr Tyr Asn Glu
 225 230 235 240
 Trp Ser Phe Ala Val Gly Lys Gln Leu Asn Met Ala Gln Ile Ile Thr
 245 250 255
 Ser Ala Gly Pro Asp Pro Val Ala Ile Thr Thr Glu Ser Val Gly Lys
 260 265 270
 Thr Ile Ala Gly Ala Lys Ile Met Gly Gln Gly Asn Asp Leu Val Leu
 275 280 285
 Asp Thr Ser Ser Phe Tyr Arg Pro Thr Gln Pro Gly Ser Tyr Pro Ile
 290 295 300
 Val Leu Ala Thr Tyr Glu Ile Val Cys Ser Lys Tyr Pro Asp Ala Thr
 305 310 315 320
 Thr Gly Thr Ala Val Arg Ala Phe Met Gln Ala Ala Ile Gly Pro Gly
 325 330 335
 Gln Glu Gly Leu Asp Gln Tyr Gly Ser Ile Pro Leu Pro Lys Ser Phe
 340 345 350
 Gln Ala Lys Leu Ala Ala Ala Val Asn Ala Ile Ser
 355 360

(2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 309 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

Gln Ala Ala Ala Gly Arg Ala Val Arg Arg Thr Gly His Ala Glu Asp
 1 5 10 15
 Gln Thr His Gln Asp Arg Leu His His Gly Cys Arg Arg Ala Ala Val
 20 25 30
 Val Val Arg Gln Asp Arg Ala Ser Val Ser Ala Thr Ser Ala Arg Pro
 35 40 45
 Pro Arg Arg His Pro Ala Gln Gly His Arg Arg Arg Val Ala Pro Ser
 50 55 60

Gly Gly Arg Arg Arg Pro His Pro His His Val Gln Pro Asp Asp Arg
 65 70 75 80
 Arg Asp Arg Pro Ala Leu Leu Asp Arg Thr Gln Pro Ala Glu His Pro
 85 90 95
 Asp Pro His Arg Arg Gly Pro Ala Asp Pro Gly Arg Val Arg Gly Arg
 100 105 110
 Gly Arg Leu Arg Arg Val Asp Asp Gly Arg Leu Gln Pro Asp Arg Asp
 115 120 125
 Ala Asp His Gly Ala Pro Val Arg Gly Arg Gly Pro His Arg Gly Val
 130 135 140
 Gln His Arg Gly Gly Pro Val Phe Val Arg Arg Val Pro Gly Val Arg
 145 150 155 160
 Cys Ala His Arg Arg Gly His Arg Arg Val Ala Ala Pro Gly Gln Gly
 165 170 175
 Asp Val Leu Arg Ala Gly Leu Arg Val Glu Arg Leu Arg Pro Val Ala
 180 185 190
 Ala Val Glu Asn Leu His Arg Gly Ser Gln Arg Ala Asp Gly Arg Val
 195 200 205
 Phe Arg Pro Ile Arg Arg Gly Ala Arg Leu Pro Ala Arg Arg Ser Arg
 210 215 220
 Ala Gly Pro Gln Gly Arg Leu His Leu Asp Gly Ala Gly Pro Ser Pro
 225 230 235 240
 Leu Pro Ala Arg Ala Gly Gln Gln Gln Pro Ser Ser Ala Gly Gly Arg
 245 250 255
 Arg Ala Gly Gly Ala Glu Arg Ala Asp Pro Gly Gln Arg Gly Arg His
 260 265 270
 His Gln Gly Gly His Asp Pro Gly Arg Gln Gly Ala Gln Arg Gly Thr
 275 280 285
 Ala Gly Val Ala His Ala Ala Ala Gly Pro Arg Arg Ala Ala Val Arg
 290 295 300
 Asn Arg Pro Arg Arg
 305

(2) INFORMATION FOR SEQ ID NO:75:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 580 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

Ser Ala Val Trp Cys Leu Asn Gly Phe Thr Gly Arg His Arg His Gly
 1 5 10 15
 Arg Cys Arg Val Arg Ala Ser Gly Trp Arg Ser Ser Asn Arg Trp Cys
 20 25 30
 Ser Thr Thr Ala Asp Cys Cys Ala Ser Lys Thr Pro Thr Gln Ala Ala
 35 40 45
 Ser Pro Leu Glu Arg Arg Phe Thr Cys Cys Ser Pro Ala Val Gly Cys
 50 55 60
 Arg Phe Arg Ser Phe Pro Val Arg Arg Leu Ala Leu Gly Ala Arg Thr
 65 70 75 80
 Ser Arg Thr Leu Gly Val Arg Arg Thr Leu Ser Gln Trp Asn Leu Ser
 85 90 95
 Pro Arg Ala Gln Pro Ser Cys Ala Val Thr Val Glu Ser His Thr His
 100 105 110
 Ala Ser Pro Arg Met Ala Lys Leu Ala Arg Val Val Gly Leu Val Gln
 115 120 125
 Glu Glu Gln Pro Ser Asp Met Thr Asn His Pro Arg Tyr Ser Pro Pro
 130 135 140
 Pro Gln Gln Pro Gly Thr Pro Gly Tyr Ala Gln Gly Gln Gln Gln Thr
 145 150 155 160
 Tyr Ser Gln Gln Phe Asp Trp Arg Tyr Pro Pro Ser Pro Pro Pro Gln
 165 170 175
 Pro Thr Gln Tyr Arg Gln Pro Tyr Glu Ala Leu Gly Gly Thr Arg Pro
 180 185 190
 Gly Leu Ile Pro Gly Val Ile Pro Thr Met Thr Pro Pro Pro Gly Met
 195 200 205
 Val Arg Gln Arg Pro Arg Ala Gly Met Leu Ala Ile Gly Ala Val Thr
 210 215 220
 Ile Ala Val Val Ser Ala Gly Ile Gly Gly Ala Ala Ala Ser Leu Val
 225 230 235 240
 Gly Phe Asn Arg Ala Pro Ala Gly Pro Ser Gly Gly Pro Val Ala Ala
 245 250 255
 Ser Ala Ala Pro Ser Ile Pro Ala Ala Asn Met Pro Pro Gly Ser Val
 260 265 270
 Glu Gln Val Ala Ala Lys Val Val Pro Ser Val Val Met Leu Glu Thr
 275 280 285
 Asp Leu Gly Arg Gln Ser Glu Glu Gly Ser Gly Ile Ile Leu Ser Ala
 290 295 300
 Glu Gly Leu Ile Leu Thr Asn Asn His Val Ile Ala Ala Ala Ala Lys
 305 310 315 320
 Pro Pro Leu Gly Ser Pro Pro Pro Lys Thr Thr Val Thr Phe Ser Asp

				325						330					335
Gly	Arg	Thr	Ala	Pro	Phe	Thr	Val	Val	Gly	Ala	Asp	Pro	Thr	Ser	Asp
			340					345					350		
Ile	Ala	Val	Val	Arg	Val	Gln	Gly	Val	Ser	Gly	Leu	Thr	Pro	Ile	Ser
		355					360					365			
Leu	Gly	Ser	Ser	Ser	Asp	Leu	Arg	Val	Gly	Gln	Pro	Val	Leu	Ala	Ile
	370					375					380				
Gly	Ser	Pro	Leu	Gly	Leu	Glu	Gly	Thr	Val	Thr	Thr	Gly	Ile	Val	Ser
385					390					395					400
Ala	Leu	Asn	Arg	Pro	Val	Ser	Thr	Thr	Gly	Glu	Ala	Gly	Asn	Gln	Asn
				405					410					415	
Thr	Val	Leu	Asp	Ala	Ile	Gln	Thr	Asp	Ala	Ala	Ile	Asn	Pro	Gly	Asn
			420					425					430		
Ser	Gly	Gly	Ala	Leu	Val	Asn	Met	Asn	Ala	Gln	Leu	Val	Gly	Val	Asn
		435					440					445			
Ser	Ala	Ile	Ala	Thr	Leu	Gly	Ala	Asp	Ser	Ala	Asp	Ala	Gln	Ser	Gly
	450					455					460				
Ser	Ile	Gly	Leu	Gly	Phe	Ala	Ile	Pro	Val	Asp	Gln	Ala	Lys	Arg	Ile
465					470					475					480
Ala	Asp	Glu	Leu	Ile	Ser	Thr	Gly	Lys	Ala	Ser	His	Ala	Ser	Leu	Gly
				485					490					495	
Val	Gln	Val	Thr	Asn	Asp	Lys	Asp	Thr	Pro	Gly	Ala	Lys	Ile	Val	Glu
			500					505					510		
Val	Val	Ala	Gly	Gly	Ala	Ala	Ala	Asn	Ala	Gly	Val	Pro	Lys	Gly	Val
		515					520					525			
Val	Val	Thr	Lys	Val	Asp	Asp	Arg	Pro	Ile	Asn	Ser	Ala	Asp	Ala	Leu
		530				535					540				
Val	Ala	Ala	Val	Arg	Ser	Lys	Ala	Pro	Gly	Ala	Thr	Val	Ala	Leu	Thr
545					550					555					560
Phe	Gln	Asp	Pro	Ser	Gly	Gly	Ser	Arg	Thr	Val	Gln	Val	Thr	Leu	Gly
				565					570					575	
Lys	Ala	Glu	Gln												
			580												

(2) INFORMATION FOR SEQ ID NO:76:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 233 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

Met Asn Asp Gly Lys Arg Ala Val Thr Ser Ala Val Leu Val Val Leu
 1 5 10 15
 Gly Ala Cys Leu Ala Leu Trp Leu Ser Gly Cys Ser Ser Pro Lys Pro
 20 25 30
 Asp Ala Glu Glu Gln Gly Val Pro Val Ser Pro Thr Ala Ser Asp Pro
 35 40 45
 Ala Leu Leu Ala Glu Ile Arg Gln Ser Leu Asp Ala Thr Lys Gly Leu
 50 55 60
 Thr Ser Val His Val Ala Val Arg Thr Thr Gly Lys Val Asp Ser Leu
 65 70 75 80
 Leu Gly Ile Thr Ser Ala Asp Val Asp Val Arg Ala Asn Pro Leu Ala
 85 90 95
 Ala Lys Gly Val Cys Thr Tyr Asn Asp Glu Gln Gly Val Pro Phe Arg
 100 105 110
 Val Gln Gly Asp Asn Ile Ser Val Lys Leu Phe Asp Asp Trp Ser Asn
 115 120 125
 Leu Gly Ser Ile Ser Glu Leu Ser Thr Ser Arg Val Leu Asp Pro Ala
 130 135 140
 Ala Gly Val Thr Gln Leu Leu Ser Gly Val Thr Asn Leu Gln Ala Gln
 145 150 155 160
 Gly Thr Glu Val Ile Asp Gly Ile Ser Thr Thr Lys Ile Thr Gly Thr
 165 170 175
 Ile Pro Ala Ser Ser Val Lys Met Leu Asp Pro Gly Ala Lys Ser Ala
 180 185 190
 Arg Pro Ala Thr Val Trp Ile Ala Gln Asp Gly Ser His His Leu Val
 195 200 205
 Arg Ala Ser Ile Asp Leu Gly Ser Gly Ser Ile Gln Leu Thr Gln Ser
 210 215 220
 Lys Trp Asn Glu Pro Val Asn Val Asp
 225 230

(2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 66 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

Val Ile Asp Ile Ile Gly Thr Ser Pro Thr Ser Trp Glu Gln Ala Ala

110

```

1           5           10           15
Ala Glu Ala Val Gln Arg Ala Arg Asp Ser Val Asp Asp Ile Arg Val
      20           25           30
Ala Arg Val Ile Glu Gln Asp Met Ala Val Asp Ser Ala Gly Lys Ile
      35           40           45
Thr Tyr Arg Ile Lys Leu Glu Val Ser Phe Lys Met Arg Pro Ala Gln
      50           55           60
Pro Arg
65

```

(2) INFORMATION FOR SEQ ID NO:78:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 69 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

```

Val Pro Pro Ala Pro Pro Leu Pro Pro Leu Pro Pro Ser Pro Ile Ser
1           5           10           15
Cys Ala Ser Pro Pro Ser Pro Pro Leu Pro Pro Ala Pro Pro Val Ala
      20           25           30
Pro Gly Pro Pro Met Pro Pro Leu Asp Pro Trp Pro Pro Ala Pro Pro
      35           40           45
Leu Pro Tyr Ser Thr Pro Pro Gly Ala Pro Leu Pro Pro Ser Pro Pro
      50           55           60
Ser Pro Pro Leu Pro
65

```

(2) INFORMATION FOR SEQ ID NO:79:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 355 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

```

Met Ser Asn Ser Arg Arg Arg Ser Leu Arg Trp Ser Trp Leu Leu Ser
1           5           10           15
Val Leu Ala Ala Val Gly Leu Gly Leu Ala Thr Ala Pro Ala Gln Ala
      20           25           30

```

Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
 35 40 45
 Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Ala Pro Gln Val Val
 50 55 60
 Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
 65 70 75 80
 Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
 85 90 95
 Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
 100 105 110
 Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
 115 120 125
 Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
 130 135 140
 Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly
 145 150 155 160
 Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
 165 170 175
 Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
 180 185 190
 Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ser
 195 200 205
 Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
 210 215 220
 Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly Phe Ala
 225 230 235 240
 Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg Ser Gly
 245 250 255
 Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu Gly Leu
 260 265 270
 Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg Val Val
 275 280 285
 Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp Val Ile
 290 295 300
 Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met Ala Asp
 305 310 315 320
 Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Asn Trp Gln
 325 330 335
 Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala Glu Gly
 340 345 350
 Pro Pro Ala

355

(2) INFORMATION FOR SEQ ID NO:80:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 205 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

Ser	Pro	Lys	Pro	Asp	Ala	Glu	Glu	Gln	Gly	Val	Pro	Val	Ser	Pro	Thr	1		5		10		15
Ala	Ser	Asp	Pro	Ala	Leu	Leu	Ala	Glu	Ile	Arg	Gln	Ser	Leu	Asp	Ala	20		25		30		
Thr	Lys	Gly	Leu	Thr	Ser	Val	His	Val	Ala	Val	Arg	Thr	Thr	Gly	Lys	35		40		45		
Val	Asp	Ser	Leu	Leu	Gly	Ile	Thr	Ser	Ala	Asp	Val	Asp	Val	Arg	Ala	50		55		60		
Asn	Pro	Leu	Ala	Ala	Lys	Gly	Val	Cys	Thr	Tyr	Asn	Asp	Glu	Gln	Gly	65		70		75		80
Val	Pro	Phe	Arg	Val	Gln	Gly	Asp	Asn	Ile	Ser	Val	Lys	Leu	Phe	Asp	85		90		95		
Asp	Trp	Ser	Asn	Leu	Gly	Ser	Ile	Ser	Glu	Leu	Ser	Thr	Ser	Arg	Val	100		105		110		
Leu	Asp	Pro	Ala	Ala	Gly	Val	Thr	Gln	Leu	Leu	Ser	Gly	Val	Thr	Asn	115		120		125		
Leu	Gln	Ala	Gln	Gly	Thr	Glu	Val	Ile	Asp	Gly	Ile	Ser	Thr	Thr	Lys	130		135		140		
Ile	Thr	Gly	Thr	Ile	Pro	Ala	Ser	Ser	Val	Lys	Met	Leu	Asp	Pro	Gly	145		150		155		160
Ala	Lys	Ser	Ala	Arg	Pro	Ala	Thr	Val	Trp	Ile	Ala	Gln	Asp	Gly	Ser	165		170		175		
His	His	Leu	Val	Arg	Ala	Ser	Ile	Asp	Leu	Gly	Ser	Gly	Ser	Ile	Gln	180		185		190		
Leu	Thr	Gln	Ser	Lys	Trp	Asn	Glu	Pro	Val	Asn	Val	Asp				195		200		205		

(2) INFORMATION FOR SEQ ID NO:81:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 286 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

Gly Asp Ser Phe Trp Ala Ala Ala Asp Gln Met Ala Arg Gly Phe Val
 1 5 10 15
 Leu Gly Ala Thr Ala Gly Arg Thr Thr Leu Thr Gly Glu Gly Leu Gln
 20 25 30
 His Ala Asp Gly His Ser Leu Leu Leu Asp Ala Thr Asn Pro Ala Val
 35 40 45
 Val Ala Tyr Asp Pro Ala Phe Ala Tyr Glu Ile Gly Tyr Ile Xaa Glu
 50 55 60
 Ser Gly Leu Ala Arg Met Cys Gly Glu Asn Pro Glu Asn Ile Phe Phe
 65 70 75 80
 Tyr Ile Thr Val Tyr Asn Glu Pro Tyr Val Gln Pro Pro Glu Pro Glu
 85 90 95
 Asn Phe Asp Pro Glu Gly Val Leu Gly Gly Ile Tyr Arg Tyr His Ala
 100 105 110
 Ala Thr Glu Gln Arg Thr Asn Lys Xaa Gln Ile Leu Ala Ser Gly Val
 115 120 125
 Ala Met Pro Ala Ala Leu Arg Ala Ala Gln Met Leu Ala Ala Glu Trp
 130 135 140
 Asp Val Ala Ala Asp Val Trp Ser Val Thr Ser Trp Gly Glu Leu Asn
 145 150 155 160
 Arg Asp Gly Val Val Ile Glu Thr Glu Lys Leu Arg His Pro Asp Arg
 165 170 175
 Pro Ala Gly Val Pro Tyr Val Thr Arg Ala Leu Glu Asn Ala Arg Gly
 180 185 190
 Pro Val Ile Ala Val Ser Asp Trp Met Arg Ala Val Pro Glu Gln Ile
 195 200 205
 Arg Pro Trp Val Pro Gly Thr Tyr Leu Thr Leu Gly Thr Asp Gly Phe
 210 215 220
 Gly Phe Ser Asp Thr Arg Pro Ala Gly Arg Arg Tyr Phe Asn Thr Asp
 225 230 235 240
 Ala Glu Ser Gln Val Gly Arg Gly Phe Gly Arg Gly Trp Pro Gly Arg
 245 250 255
 Arg Val Asn Ile Asp Pro Phe Gly Ala Gly Arg Gly Pro Pro Ala Gln
 260 265 270
 Leu Pro Gly Phe Asp Glu Gly Gly Gly Leu Arg Pro Xaa Lys
 275 280 285

(2) INFORMATION FOR SEQ ID NO:82:

114

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 173 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

Thr	Lys	Phe	His	Ala	Leu	Met	Gln	Glu	Gln	Ile	His	Asn	Glu	Phe	Thr
1				5					10					15	
Ala	Ala	Gln	Gln	Tyr	Val	Ala	Ile	Ala	Val	Tyr	Phe	Asp	Ser	Glu	Asp
			20					25					30		
Leu	Pro	Gln	Leu	Ala	Lys	His	Phe	Tyr	Ser	Gln	Ala	Val	Glu	Glu	Arg
			35				40					45			
Asn	His	Ala	Met	Met	Leu	Val	Gln	His	Leu	Leu	Asp	Arg	Asp	Leu	Arg
	50					55					60				
Val	Glu	Ile	Pro	Gly	Val	Asp	Thr	Val	Arg	Asn	Gln	Phe	Asp	Arg	Pro
65					70					75					80
Arg	Glu	Ala	Leu	Ala	Leu	Ala	Leu	Asp	Gln	Glu	Arg	Thr	Val	Thr	Asp
				85					90					95	
Gln	Val	Gly	Arg	Leu	Thr	Ala	Val	Ala	Arg	Asp	Glu	Gly	Asp	Phe	Leu
			100					105					110		
Gly	Glu	Gln	Phe	Met	Gln	Trp	Phe	Leu	Gln	Glu	Gln	Ile	Glu	Glu	Val
			115				120					125			
Ala	Leu	Met	Ala	Thr	Leu	Val	Arg	Val	Ala	Asp	Arg	Ala	Gly	Ala	Asn
			130			135					140				
Leu	Phe	Glu	Leu	Glu	Asn	Phe	Val	Ala	Arg	Glu	Val	Asp	Val	Ala	Pro
145					150					155					160
Ala	Ala	Ser	Gly	Ala	Pro	His	Ala	Ala	Gly	Gly	Arg	Leu			
				165					170						

(2) INFORMATION FOR SEQ ID NO:83:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 107 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

Arg	Ala	Asp	Glu	Arg	Lys	Asn	Thr	Thr	Met	Lys	Met	Val	Lys	Ser	Ile
1					5				10					15	

115

Ala Ala Gly Leu Thr Ala Ala Ala Ala Ile Gly Ala Ala Ala Ala Gly
 20 25 30

Val Thr Ser Ile Met Ala Gly Gly Pro Val Val Tyr Gln Met Gln Pro
 35 40 45

Val Val Phe Gly Ala Pro Leu Pro Leu Asp Pro Xaa Ser Ala Pro Xaa
 50 55 60

Val Pro Thr Ala Ala Gln Trp Thr Xaa Leu Leu Asn Xaa Leu Xaa Asp
 65 70 75 80

Pro Asn Val Ser Phe Xaa Asn Lys Gly Ser Leu Val Glu Gly Gly Ile
 85 90 95

Gly Gly Xaa Glu Gly Xaa Xaa Arg Arg Xaa Gln
 100 105

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 125 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

Val Leu Ser Val Pro Val Gly Asp Gly Phe Trp Xaa Arg Val Val Asn
 1 5 10 15

Pro Leu Gly Gln Pro Ile Asp Gly Arg Gly Asp Val Asp Ser Asp Thr
 20 25 30

Arg Arg Ala Leu Glu Leu Gln Ala Pro Ser Val Val Xaa Arg Gln Gly
 35 40 45

Val Lys Glu Pro Leu Xaa Thr Gly Ile Lys Ala Ile Asp Ala Met Thr
 50 55 60

Pro Ile Gly Arg Gly Gln Arg Gln Leu Ile Ile Gly Asp Arg Lys Thr
 65 70 75 80

Gly Lys Asn Arg Arg Leu Cys Arg Thr Pro Ser Ser Asn Gln Arg Glu
 85 90 95

Glu Leu Gly Val Arg Trp Ile Pro Arg Ser Arg Cys Ala Cys Val Tyr
 100 105 110

Val Gly His Arg Ala Arg Arg Gly Thr Tyr His Arg Arg
 115 120 125

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 117 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

[illegible]

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 103 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

Met 1	Tyr	Arg	Phe	Ala 5	Cys	Arg	Thr	Leu	Met 10	Leu	Ala	Ala	Cys	Ile 15	Leu
Ala	Thr	Gly	Val 20	Ala	Gly	Leu	Gly	Val 25	Gly	Ala	Gln	Ser	Ala 30	Ala	Gln
Thr	Ala	Pro 35	Val	Pro	Asp	Tyr	Tyr 40	Trp	Cys	Pro	Gly	Gln 45	Pro	Phe	Asp
Pro	Ala 50	Trp	Gly	Pro	Asn	Trp 55	Asp	Pro	Tyr	Thr	Cys 60	His	Asp	Asp	Phe
His 65	Arg	Asp	Ser	Asp	Gly 70	Pro	Asp	His	Ser	Arg 75	Asp	Tyr	Pro	Gly	Pro 80
Ile	Leu	Glu	Gly	Pro	Val	Leu	Asp	Asp	Pro	Gly	Ala	Ala	Pro	Pro	Pro

117

85

90

95

Pro Ala Ala Gly Gly Gly Ala
100

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 88 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

Val Gln Cys Arg Val Trp Leu Glu Ile Gln Trp Arg Gly Met Leu Gly
1 5 10 15
Ala Asp Gln Ala Arg Ala Gly Gly Pro Ala Arg Ile Trp Arg Glu His
20 25 30
Ser Met Ala Ala Met Lys Pro Arg Thr Gly Asp Gly Pro Leu Glu Ala
35 40 45
Thr Lys Glu Gly Arg Gly Ile Val Met Arg Val Pro Leu Glu Gly Gly
50 55 60
Gly Arg Leu Val Val Glu Leu Thr Pro Asp Glu Ala Ala Ala Leu Gly
65 70 75 80
Asp Glu Leu Lys Gly Val Thr Ser
85

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 95 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

Thr Asp Ala Ala Thr Leu Ala Gln Glu Ala Gly Asn Phe Glu Arg Ile
1 5 10 15
Ser Gly Asp Leu Lys Thr Gln Ile Asp Gln Val Glu Ser Thr Ala Gly
20 25 30
Ser Leu Gln Gly Gln Trp Arg Gly Ala Ala Gly Thr Ala Ala Gln Ala
35 40 45
Ala Val Val Arg Phe Gln Glu Ala Ala Asn Lys Gln Lys Gln Glu Leu
50 55 60

118

Asp Glu Ile Ser Thr Asn Ile Arg Gln Ala Gly Val Gln Tyr Ser Arg
 65 70 75 80
 Ala Asp Glu Glu Gln Gln Gln Ala Leu Ser Ser Gln Met Gly Phe
 85 90 95

(2) INFORMATION FOR SEQ ID NO:89:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 166 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

Met Thr Gln Ser Gln Thr Val Thr Val Asp Gln Gln Glu Ile Leu Asn
 1 5 10 15
 Arg Ala Asn Glu Val Glu Ala Pro Met Ala Asp Pro Pro Thr Asp Val
 20 25 30
 Pro Ile Thr Pro Cys Glu Leu Thr Xaa Xaa Lys Asn Ala Ala Gln Gln
 35 40 45
 Xaa Val Leu Ser Ala Asp Asn Met Arg Glu Tyr Leu Ala Ala Gly Ala
 50 55 60
 Lys Glu Arg Gln Arg Leu Ala Thr Ser Leu Arg Asn Ala Ala Lys Xaa
 65 70 75 80
 Tyr Gly Glu Val Asp Glu Glu Ala Ala Thr Ala Leu Asp Asn Asp Gly
 85 90 95
 Glu Gly Thr Val Gln Ala Glu Ser Ala Gly Ala Val Gly Gly Asp Ser
 100 105 110
 Ser Ala Glu Leu Thr Asp Thr Pro Arg Val Ala Thr Ala Gly Glu Pro
 115 120 125
 Asn Phe Met Asp Leu Lys Glu Ala Ala Arg Lys Leu Glu Thr Gly Asp
 130 135 140
 Gln Gly Ala Ser Leu Ala His Xaa Gly Asp Gly Trp Asn Thr Xaa Thr
 145 150 155 160
 Leu Thr Leu Gln Gly Asp
 165

(2) INFORMATION FOR SEQ ID NO:90:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

Arg Ala Glu Arg Met
1 5

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 263 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

Val 1	Ala	Trp	Met	Ser 5	Val	Thr	Ala	Gly 10	Gln	Ala	Glu	Leu	Thr	Ala	Ala 15
Gln	Val	Arg	Val 20	Ala	Ala	Ala	Ala	Tyr 25	Glu	Thr	Ala	Tyr	Gly 30	Leu	Thr
Val	Pro	Pro 35	Pro	Val	Ile	Ala	Glu 40	Asn	Arg	Ala	Glu	Leu 45	Met	Ile	Leu
Ile	Ala 50	Thr	Asn	Leu	Leu	Gly 55	Gln	Asn	Thr	Pro	Ala 60	Ile	Ala	Val	Asn
Glu 65	Ala	Glu	Tyr	Gly	Glu 70	Met	Trp	Ala	Gln	Asp 75	Ala	Ala	Ala	Met	Phe 80
Gly	Tyr	Ala	Ala	Ala 85	Thr	Ala	Thr	Ala	Thr 90	Ala	Thr	Leu	Leu	Pro 95	Phe
Glu	Glu	Ala	Pro 100	Glu	Met	Thr	Ser	Ala 105	Gly	Gly	Leu	Leu	Glu 110	Gln	Ala
Ala	Ala	Val 115	Glu	Glu	Ala	Ser	Asp 120	Thr	Ala	Ala	Ala	Asn 125	Gln	Leu	Met
Asn 130	Asn	Val	Pro	Gln	Ala	Leu 135	Lys	Gln	Leu	Ala	Gln 140	Pro	Thr	Gln	Gly
Thr 145	Thr	Pro	Ser	Ser	Lys 150	Leu	Gly	Gly	Leu	Trp 155	Lys	Thr	Val	Ser	Pro 160
His	Arg	Ser	Pro	Ile 165	Ser	Asn	Met	Val	Ser 170	Met	Ala	Asn	Asn	His 175	Met
Ser	Met	Thr	Asn 180	Ser	Gly	Val	Ser	Met 185	Thr	Asn	Thr	Leu	Ser 190	Ser	Met
Leu	Lys	Gly 195	Phe	Ala	Pro	Ala	Ala	Ala 200	Ala	Gln	Ala	Val 205	Gln	Thr	Ala

120

Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
 210 215 220

Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala
 225 230 235 240

Ser Val Arg Tyr Gly His Arg Asp Gly Gly Lys Tyr Ala Xaa Ser Gly
 245 250 255

Arg Arg Asn Gly Gly Pro Ala
 260

(2) INFORMATION FOR SEQ ID NO:92:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 303 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

Met Thr Tyr Ser Pro Gly Asn Pro Gly Tyr Pro Gln Ala Gln Pro Ala
 1 5 10 15

Gly Ser Tyr Gly Gly Val Thr Pro Ser Phe Ala His Ala Asp Glu Gly
 20 25 30

Ala Ser Lys Leu Pro Met Tyr Leu Asn Ile Ala Val Ala Val Leu Gly
 35 40 45

Leu Ala Ala Tyr Phe Ala Ser Phe Gly Pro Met Phe Thr Leu Ser Thr
 50 55 60

Glu Leu Gly Gly Gly Asp Gly Ala Val Ser Gly Asp Thr Gly Leu Pro
 65 70 75 80

Val Gly Val Ala Leu Leu Ala Ala Leu Leu Ala Gly Val Val Leu Val
 85 90 95

Pro Lys Ala Lys Ser His Val Thr Val Val Ala Val Leu Gly Val Leu
 100 105 110

Gly Val Phe Leu Met Val Ser Ala Thr Phe Asn Lys Pro Ser Ala Tyr
 115 120 125

Ser Thr Gly Trp Ala Leu Trp Val Val Leu Ala Phe Ile Val Phe Gln
 130 135 140

Ala Val Ala Ala Val Leu Ala Leu Leu Val Glu Thr Gly Ala Ile Thr
 145 150 155 160

Ala Pro Ala Pro Arg Pro Lys Phe Asp Pro Tyr Gly Gln Tyr Gly Arg
 165 170 175

Tyr Gly Gln Tyr Gly Gln Tyr Gly Val Gln Pro Gly Gly Tyr Tyr Gly
 180 185 190

121

Gln Gln Gly Ala Gln Gln Ala Ala Gly Leu Gln Ser Pro Gly Pro Gln
 195 200 205

Gln Ser Pro Gln Pro Pro Gly Tyr Gly Ser Gln Tyr Gly Gly Tyr Ser
 210 215 220

Ser Ser Pro Ser Gln Ser Gly Ser Gly Tyr Thr Ala Gln Pro Pro Ala
 225 230 235 240

Gln Pro Pro Ala Gln Ser Gly Ser Gln Gln Ser His Gln Gly Pro Ser
 245 250 255

Thr Pro Pro Thr Gly Phe Pro Ser Phe Ser Pro Pro Pro Pro Val Ser
 260 265 270

Ala Gly Thr Gly Ser Gln Ala Gly Ser Ala Pro Val Asn Tyr Ser Asn
 275 280 285

Pro Ser Gly Gly Glu Gln Ser Ser Ser Pro Gly Gly Ala Pro Val
 290 295 300

(2) INFORMATION FOR SEQ ID NO:93:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

Gly Cys Gly Glu Thr Asp Ala Ala Thr Leu Ala Gln Glu Ala Gly Asn
 1 5 10 15

Phe Glu Arg Ile Ser Gly Asp Leu Lys Thr Gln Ile
 20 25

(2) INFORMATION FOR SEQ ID NO:94:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

Asp Gln Val Glu Ser Thr Ala Gly Ser Leu Gln Gly Gln Trp Arg Gly
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:95:

- (i) SEQUENCE CHARACTERISTICS:

122

- (A) LENGTH: 27 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

Gly Cys Gly Ser Thr Ala Gly Ser Leu Gln Gly Gln Trp Arg Gly Ala
1 5 10 15
Ala Gly Thr Ala Ala Gln Ala Ala Val Val Arg
 20 25

(2) INFORMATION FOR SEQ ID NO:96:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

Gly Cys Gly Gly Thr Ala Ala Gln Ala Ala Val Val Arg Phe Gln Glu
1 5 10 15
Ala Ala Asn Lys Gln Lys Gln Glu Leu Asp Glu
 20 25

(2) INFORMATION FOR SEQ ID NO:97:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

Gly Cys Gly Ala Asn Lys Gln Lys Gln Glu Leu Asp Glu Ile Ser Thr
1 5 10 15
Asn Ile Arg Gln Ala Gly Val Gln Tyr Ser Arg
 20 25

(2) INFORMATION FOR SEQ ID NO:98:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 amino acids
 - (B) TYPE: amino acid

123

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

Gly	Cys	Gly	Ile	Arg	Gln	Ala	Gly	Val	Gln	Tyr	Ser	Arg	Ala	Asp	Glu
1				5				10						15	
Glu	Gln	Gln	Gln	Ala	Leu	Ser	Ser	Gln	Met	Gly	Phe				
			20					25							

(2) INFORMATION FOR SEQ ID NO:99:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 507 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

ATGAAGATGG TGAAATCGAT CGCCGCAGGT CTGACCGCCG CGGCTGCAAT CGGCGCCGCT	60
GCGGCCGGTG TGAATTCGAT CATGGCTGGC GGCCCGGTCG TATACCAGAT GCAGCCGGTG	120
GTCTTCGGCG CGCCACTGCC GTTGGACCCG GCATCCGCCC CTGACGTCCC GACCGCCGCC	180
CAGTTGACCA GCCTGCTCAA CAGCCTCGCC GATCCCAACG TGTCGTTTGC GAACAAGGGC	240
AGTCTGGTCG AGGGCGGCAT CGGGGGCACC GAGGCGCGCA TCGCCGACCA CAAGCTGAAG	300
AAGGCCGCCG AGCACGGGGA TCTGCCGCTG TCGTTCAGCG TGACGAACAT CCAGCCGGCG	360
GCCGCCGGTT CGGCCACCGC CGACGTTTCC GTCTCGGGTC CGAAGCTCTC GTCGCCGGTC	420
ACGCAGAACG TCACGTTTCGT GAATCAAGGC GGCTGGATGC TGTCACGCGC ATCGGCGATG	480
GAGTTGCTGC AGGCCGCAGG GAACTGA	507

(2) INFORMATION FOR SEQ ID NO:100:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 168 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

124

Met Lys Met Val Lys Ser Ile Ala Ala Gly Leu Thr Ala Ala Ala Ala
 1 5 10 15
 Ile Gly Ala Ala Ala Ala Gly Val Thr Ser Ile Met Ala Gly Gly Pro
 20 25 30
 Val Val Tyr Gln Met Gln Pro Val Val Phe Gly Ala Pro Leu Pro Leu
 35 40 45
 Asp Pro Ala Ser Ala Pro Asp Val Pro Thr Ala Ala Gln Leu Thr Ser
 50 55 60
 Leu Leu Asn Ser Leu Ala Asp Pro Asn Val Ser Phe Ala Asn Lys Gly
 65 70 75 80
 Ser Leu Val Glu Gly Gly Ile Gly Gly Thr Glu Ala Arg Ile Ala Asp
 85 90 95
 His Lys Leu Lys Lys Ala Ala Glu His Gly Asp Leu Pro Leu Ser Phe
 100 105 110
 Ser Val Thr Asn Ile Gln Pro Ala Ala Ala Gly Ser Ala Thr Ala Asp
 115 120 125
 Val Ser Val Ser Gly Pro Lys Leu Ser Ser Pro Val Thr Gln Asn Val
 130 135 140
 Thr Phe Val Asn Gln Gly Gly Trp Met Leu Ser Arg Ala Ser Ala Met
 145 150 155 160
 Glu Leu Leu Gln Ala Ala Gly Asn
 165

(2) INFORMATION FOR SEQ ID NO:101:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 500 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

CGTGGCAATG TCGTTGACCG TCGGGGCCGG GGTGCGCTCC GCAGATCCCG TGGACGCGGT 60
 CATTAACACC ACCTGCAATT ACGGGCAGGT AGTAGCTGCG CTCAACGCGA CGGATCCGGG 120
 GGCTGCCGCA CAGTTCAACG CCTCACCGGT GGCGCAGTCC TATTTGCGCA ATTTCTCGC 180
 CGCACCGCCA CCTCAGCGCG CTGCCATGGC CGCGCAATTG CAAGCTGTGC CGGGGGCGGC 240
 ACAGTACATC GGCCTTGTCG AGTCGGTTGC CGGCTCCTGC AACAACTATT AAGCCCATGC 300
 GGGCCCCATC CCGCGACCCG GCATCGTCGC CGGGGCTAGG CCAGATTGCC CCGCTCCTCA 360
 ACGGGCCGCA TCCCGCGACC CGGCATCGTC GCCGGGGCTA GGCCAGATTG CCCCCTCCT 420
 CAACGGGCCG CATCTCGTGC CGAATTCCTG CAGCCCGGGG GATCCACTAG TTCTAGAGCG 480

GCCGCCACCG CGGTGGAGCT

500

(2) INFORMATION FOR SEQ ID NO:102:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 96 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

```

Val Ala Met Ser Leu Thr Val Gly Ala Gly Val Ala Ser Ala Asp Pro
1           5           10           15
Val Asp Ala Val Ile Asn Thr Thr Cys Asn Tyr Gly Gln Val Val Ala
20           25           30
Ala Leu Asn Ala Thr Asp Pro Gly Ala Ala Ala Gln Phe Asn Ala Ser
35           40           45
Pro Val Ala Gln Ser Tyr Leu Arg Asn Phe Leu Ala Ala Pro Pro Pro
50           55           60
Gln Arg Ala Ala Met Ala Ala Gln Leu Gln Ala Val Pro Gly Ala Ala
65           70           75           80
Gln Tyr Ile Gly Leu Val Glu Ser Val Ala Gly Ser Cys Asn Asn Tyr
85           90           95

```

(2) INFORMATION FOR SEQ ID NO:103:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 154 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

```

ATGACAGAGC AGCAGTGGAA TTTCGCGGGT ATCGAGGCCG CGGCAAGCGC AATCCAGGGA      60
AATGTCACGT CCATTCATTC CCTCCTTGAC GAGGGGAAGC AGTCCCTGAC CAAGCTCGCA      120
GCGGCCTGGG GCGGTAGCGG TTCGGAAGCG TACC                                154

```

(2) INFORMATION FOR SEQ ID NO:104:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 51 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

```

Met Thr Glu Gln Gln Trp Asn Phe Ala Gly Ile Glu Ala Ala Ala Ser
 1           5           10           15
Ala Ile Gln Gly Asn Val Thr Ser Ile His Ser Leu Leu Asp Glu Gly
          20           25           30
Lys Gln Ser Leu Thr Lys Leu Ala Ala Ala Trp Gly Gly Ser Gly Ser
          35           40           45
Glu Ala Tyr
          50

```

(2) INFORMATION FOR SEQ ID NO:105:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 282 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

```

CGGTCGCGCA CTTCCAGGTG ACTATGAAAG TCGGCTTCCG NCTGGAGGAT TCCTGAACCT      60
TCAAGCGCGG CCGATAACTG AGGTGCATCA TTAAGCGACT TTTCCAGAAC ATCCTGACGC      120
GCTCGAAACG CGGCACAGCC GACGGTGGCT CCGNCGAGGC GCTGNCTCCA AAATCCCTGA      180
GACAATTTCG CGGGGGCGCC TACAAGGAAG TCGGTGCTGA ATTCGNCGNG TATCTGGTCG      240
ACCTGTGTGG TCTGNAGCCG GACGAAGCGG TGCTCGACGT CG                          282

```

(2) INFORMATION FOR SEQ ID NO:106:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 3058 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

```

GATCGTACCC GTGCGAGTGC TCGGGCCGTT TGAGGATGGA GTGCACGTGT CTTTCGTGAT      60
GGCATACCCA GAGATGTTGG CGGCGGCGGC TGACACCCTG CAGAGCATCG GTGCTACCAC      120
TGTGGCTAGC AATGCCGCTG CGGCGGCCCC GACGACTGGG GTGGTGCCCC CCGCTGCCGA      180

```

TGAGGTGTCG	GCGCTGACTG	CGGCGCACTT	CGCCGCACAT	GCGGCGATGT	ATCAGTCCGT	240
GAGCGCTCGG	GCTGCTGCGA	TTCATGACCA	GTTCGTGGCC	ACCCTTGCCA	GCAGCGCCAG	300
CTCGTATGCG	GCCACTGAAG	TCGCCAATGC	GGCGGCGGCC	AGCTAAGCCA	GGAACAGTCG	360
GCACGAGAAA	CCACGAGAAA	TAGGGACACG	TAATGGTGGG	TTTCGGGGCG	TTACCACCGG	420
AGATCAACTC	CGCGAGGATG	TACGCCGGCC	CGGGTTCGGC	CTCGCTGGTG	GCCGCGGCTC	480
AGATGTGGGA	CAGCGTGGCG	AGTGACCTGT	TTTCGGCCGC	GTCGGCGTTT	CAGTCGGTGG	540
TCTGGGGTCT	GACGGTGGGG	TCGTGGATAG	GTTCGTGCGC	GGGTCTGATG	GTGGCGGCGG	600
CCTCGCCGTA	TGTGGCGTGG	ATGAGCGTCA	CCGCGGGGCA	GGCCGAGCTG	ACCGCCGCCC	660
AGGTCCGGGT	TGCTGCGGCG	GCCTACGAGA	CGGCGTATGG	GCTGACGGTG	CCCCCGCCGG	720
TGATCGCCGA	GAACCGTGCT	GAAGTATGTA	TTCTGATAGC	GACCAACCTC	TTGGGGCAAA	780
ACACCCCGGC	GATCGCGGTC	AACGAGGCCG	AATACGGCGA	GATGTGGGCC	CAAGACGCCG	840
CCGCGATGTT	TGGCTACGCC	GCGGCGACGG	CGACGGCGAC	GGCGACGTTG	CTGCCGTTCG	900
AGGAGGCGCC	GGAGATGACC	AGCGCGGGTG	GGCTCCTCGA	GCAGGCCGCC	GCGGTCGAGG	960
AGGCCTCCGA	CACCGCCGCG	GCGAACCAGT	TGATGAACAA	TGTGCCCCAG	GCGCTGCAAC	1020
AGCTGGCCCA	GCCCACGCAG	GGCACCACGC	CTTCTTCCAA	GCTGGGTGGC	CTGTGGAAGA	1080
CGGTCTCGCC	GCATCGGTCT	CCGATCAGCA	ACATGGTGTC	GATGGCCAAC	AACCACATGT	1140
CGATGACCAA	CTCGGTGTG	TCGATGACCA	ACACCTTGAG	CTCGATGTTG	AAGGGCTTTG	1200
CTCCGGCGGC	GGCCGCCCAG	GCCGTGCAAA	CCGCGGCGCA	AAACGGGGTC	CGGGCGATGA	1260
GCTCGCTGGG	CAGCTCGCTG	GGTTCTTCGG	GTCTGGGCGG	TGGGGTGGCC	GCCAAGTTGG	1320
GTCGGGCGGC	CTCGGTCGGT	TCGTTGTCTG	TGCCGCAGGC	CTGGGCCGCG	GCCAACCAGG	1380
CAGTCACCCC	GGCGGCGCGG	GCGCTGCCGC	TGACCAGCCT	GACCAGCGCC	GCGGAAAGAG	1440
GGCCCGGGCA	GATGCTGGGC	GGGCTGCCGG	TGGGGCAGAT	GGGCGCCAGG	GCCGGTGGTG	1500
GGCTCAGTGG	TGTGCTGCGT	GTTCCGCCGC	GACCCTATGT	GATGCCGCAT	TCTCCGGCGG	1560
CCGGCTAGGA	GAGGGGGCGC	AGACTGTCTG	TATTTGACCA	GTGATCGGCG	GTCTCGGTGT	1620
TTCCGCGGCC	GGCTATGACA	ACAGTCAATG	TGCATGACAA	GTTACAGGTA	TTAGGTCCAG	1680
GTTCAACAAG	GAGACAGGCA	ACATGGCCTC	ACGTTTTATG	ACGGATCCGC	ACGCGATGCG	1740
GGACATGGCG	GGCCGTTTTG	AGGTGCACGC	CCAGACGGTG	GAGGACGAGG	CTCGCCGGAT	1800
GTGGGCGTCC	GCGCAAAACA	TTTCCGGTGC	GGGCTGGAGT	GGCATGGCCG	AGGCGACCTC	1860
GCTAGACACC	ATGGCCCAGA	TGAATCAGGC	GTTTCGCAAC	ATCGTGAACA	TGCTGCACGG	1920
GGTGCGTGAC	GGGCTGGTTC	GCGACGCCAA	CAACTACGAG	CAGCAAGAGC	AGGCCTCCCA	1980
GCAGATCCTC	AGCAGCTAAC	GTCAGCCGCT	GCAGCACAAT	ACTTTTACAA	GCGAAGGAGA	2040

```

ACAGGTTCTGA TGACCATCAA CTATCAATTC GGGGATGTCG ACGCTCACGG CGCCATGATC      2100
CGCGCTCAGG CCGGGTTGCT GGAGGCCGAG CATCAGGCCA TCATTCTGTA TGTGTTGACC      2160
GCGAGTGA CT TTTGGGGCGG CGCCGGTTCTG GCGGCCTGCC AGGGGTTTCAT TACCCAGTTG      2220
GGCCGTAACT TCCAGGTGAT CTACGAGCAG GCCAACGCCC ACGGGCAGAA GGTGCAGGCT      2280
GCCGGCAACA ACATGGCGCA AACCGACAGC GCCGTCGGCT CCAGCTGGGC CTGACACCAG      2340
GCCAAGGCCA GGGACGTGGT GTACGAGTGA AGTTCCTCGC GTGATCCTTC GGGTGGCAGT      2400
CTAAGTGGTC AGTGCTGGGG TGTTGGTGGT TTGCTGCTTG GCGGGTTCTT CGGTGCTGGT      2460
CAGTGCTGCT CGGGCTCGGG TGAGGACCTC GAGGCCCAGG TAGCGCCGTC CTTTCGATCCA      2520
TTCGTCGTGT TGTTCTGGCGA GGACGGCTCC GACGAGGCGG ATGATCGAGG CGCGGTCGGG      2580
GAAGATGCCC ACGACGTCGG TTCGGCGTCG TACCTCTCGG TTGAGGCGTT CCTGGGGGTT      2640
GTTGGACCAG ATTTGGCGCC AGATCTGCTT GGGGAAGGCG GTGAACGCCA GCAGGTTCGGT      2700
GCGGGCGGTG TCGAGGTGCT CGGCCACCGC GGGGAGTTTG TCGGTCAGAG CGTCGAGTAC      2760
CCGATCATAT TGGGCAACAA CTGATTCTGGC GTCGGGCTGG TCGTAGATGG AGTGCAGCAG      2820
GGTGCACACC CACGGCCAGG AGGGCTTCGG GGTGGCTGCC ATCAGATTGG CTGCGTAGTG      2880
GGTTCCTGCAG CGCTGCCAGG CCGCTGCGGG CAGGGTGGCG CCGATCGCGG CCACCAGGCC      2940
GGCGTGGGCG TCGCTGGTGA CCAGCGCGAC CCCGGACAGG CCGCGGGCGA CCAGGTTCGG      3000
GAAGAACGCC AGCCAGCCGG CCCCGTCCTC GGCGGAGGTG ACCTGGATGC CCAGGATC      3058

```

(2) INFORMATION FOR SEQ ID NO:107:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 391 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

```

Met Val Asp Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met
1           5           10           15
Tyr Ala Gly Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Gln Met Trp
20           25           30
Asp Ser Val Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser
35           40           45
Val Val Trp Gly Leu Thr Val Gly Ser Trp Ile Gly Ser Ser Ala Gly
50           55           60
Leu Met Val Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr

```


(2) INFORMATION FOR SEQ ID NO:108:

(i). SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1725 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

GACGTCAGCA CCCGCCGTGC AGGGCTGGAG CGTGGTCGGT TTTGATCTGC GGTCAAGGTG	60
ACGTCCCTCG GCGTGTGCGC GCGGTGGATG CAGACTCGAT GCCGCTCTTT AGTGCAACTA	120
ATTTCTGTTGA AGTGCCTGCG AGGTATAGGA CTTACAGATT GGTTAATGTA GCGTTCACCC	180
CGTGTTGGGG TCGATTGGC CGGACCAGTC GTCACCAACG CTTGGCGTGC GCGCCAGGCG	240
GGCGATCAGA TCGCTTGACT ACCAATCAAT CTTGAGCTCC CGGGCCGATG CTCGGGCTAA	300
ATGAGGAGGA GCACGCGTGT CTTTCACTGC GCAACCGGAG ATGTTGGCGG CCGCGGCTGG	360
CGAACTTCGT TCCCTGGGGG CAACGCTGAA GGCTAGCAAT GCCGCCGCAG CCGTGCCGAC	420
GACTGGGGTG GTGCCCCCGG CTGCCGACGA GGTGTCGCTG CTGCTTGCCA CACAATTCCG	480
TACGCATGCG GCGACGTATC AGACGGCCAG CGCCAAGGCC GCGGTGATCC ATGAGCAGTT	540
TGTGACCACG CTGGCCACCA GCGCTAGTTC ATATGCGGAC ACCGAGGCCG CCAACGCTGT	600
GGTCACCGGC TAGCTGACCT GACGGTATTC GAGCGGAAGG ATTATCGAAG TGGTGGATTT	660
CGGGGCGTTA CCACCGGAGA TCAACTCCGC GAGGATGTAC GCCGGCCCGG GTTCGGCCTC	720
GCTGGTGGCC GCCGCGAAGA TGTGGGACAG CGTGGCGAGT GACCTGTTTT CGGCCGCGTC	780
GGCGTTTCAG TCGGTGGTCT GGGGTCTGAC GGTGGGGTCG TGGATAGGTT CGTCGGCGGG	840
TCTGATGGCG GCGGCGGCCT CGCCGTATGT GGCGTGGATG AGCGTCACCG CGGGGCAGGC	900
CCAGCTGACC GCCGCCCAGG TCCGGGTTGC TGCGGCGGCC TACGAGACAG CGTATAGGCT	960
GACGGTGCCC CCGCCGGTGA TCGCCGAGAA CCGTACCGAA CTGATGACGC TGACCGCGAC	1020
CAACCTCTTG GGGCAAAACA CGCCGGCGAT CGAGGCCAAT CAGGCCGCAT ACAGCCAGAT	1080
GTGGGGCCAA GACGCGGAGG CGATGTATGG CTACGCCGCC ACGGCGGCGA CGGCGACCGA	1140
GGCGTTGCTG CCGTTCGAGG ACGCCCCACT GATACCAAC CCCGGCGGGC TCCTTGAGCA	1200
GGCCGTCGCG GTCGAGGAGG CCATCGACAC CGCCGCGGCG AACCAGTTGA TGAACAATGT	1260
GCCCCAAGCG CTGCAACAGC TGGCCCAGCC AGCGCAGGGC GTCGTACCTT CTTCCAAGCT	1320
GGGTGGGCTG TGGACGGCGG TCTCGCCGCA TCTGTGCGCG CTCAGCAACG TCAGTTCGAT	1380
AGCCAACAAC CACATGTCGA TGATGGGCAC GGGTGTGTGCG ATGACCAACA CCTTGCACTC	1440

GATGTTGAAG GGCTTAGCTC CGGCGGCGGC TCAGGCCGTG GAAACCGCGG CGGAAAACGG 1500
 GGTCTGGGCG ATGAGCTCGC TGGGCAGCCA GCTGGGTTCG TCGCTGGGTT CTTCGGGTCT 1560
 GGGCGCTGGG GTGGCCGCCA ACTTGGGTCG GGCGGCCTCG GTCGGTTCGT TGTCGGTGCC 1620
 GCCAGCATGG GCCGCGGCCA ACCAGGCGGT CACCCCGGCG GCGCGGGCGC TGCCGCTGAC 1680
 CAGCCTGACC AGCGCCGCCC AAACCGCCCC CGGACACATG CTGGG 1725

(2) INFORMATION FOR SEQ ID NO:109:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 359 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

Val Val Asp Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met
 1 5 10 15
 Tyr Ala Gly Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Lys Met Trp
 20 25 30
 Asp Ser Val Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser
 35 40 45
 Val Val Trp Gly Leu Thr Val Gly Ser Trp Ile Gly Ser Ser Ala Gly
 50 55 60
 Leu Met Ala Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr
 65 70 75 80
 Ala Gly Gln Ala Gln Leu Thr Ala Ala Gln Val Arg Val Ala Ala Ala
 85 90 95
 Ala Tyr Glu Thr Ala Tyr Arg Leu Thr Val Pro Pro Pro Val Ile Ala
 100 105 110
 Glu Asn Arg Thr Glu Leu Met Thr Leu Thr Ala Thr Asn Leu Leu Gly
 115 120 125
 Gln Asn Thr Pro Ala Ile Glu Ala Asn Gln Ala Ala Tyr Ser Gln Met
 130 135 140
 Trp Gly Gln Asp Ala Glu Ala Met Tyr Gly Tyr Ala Ala Thr Ala Ala
 145 150 155 160
 Thr Ala Thr Glu Ala Leu Leu Pro Phe Glu Asp Ala Pro Leu Ile Thr
 165 170 175
 Asn Pro Gly Gly Leu Leu Glu Gln Ala Val Ala Val Glu Glu Ala Ile
 180 185 190
 Asp Thr Ala Ala Ala Asn Gln Leu Met Asn Asn Val Pro Gln Ala Leu
 195 200 205

132

Gln Gln Leu Ala Gln Pro Ala Gln Gly Val Val Pro Ser Ser Lys Leu
 210 215 220
 Gly Gly Leu Trp Thr Ala Val Ser Pro His Leu Ser Pro Leu Ser Asn
 225 230 235 240
 Val Ser Ser Ile Ala Asn Asn His Met Ser Met Met Gly Thr Gly Val
 245 250 255
 Ser Met Thr Asn Thr Leu His Ser Met Leu Lys Gly Leu Ala Pro Ala
 260 265 270
 Ala Ala Gln Ala Val Glu Thr Ala Ala Glu Asn Gly Val Trp Ala Met
 275 280 285
 Ser Ser Leu Gly Ser Gln Leu Gly Ser Ser Leu Gly Ser Ser Gly Leu
 290 295 300
 Gly Ala Gly Val Ala Ala Asn Leu Gly Arg Ala Ala Ser Val Gly Ser
 305 310 315 320
 Leu Ser Val Pro Pro Ala Trp Ala Ala Ala Asn Gln Ala Val Thr Pro
 325 330 335
 Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser Ala Ala Gln Thr
 340 345 350
 Ala Pro Gly His Met Leu Gly
 355

(2) INFORMATION FOR SEQ ID NO:110:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 3027 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

AGTTCAGTCG AGAATGATAC TGACGGGCTG TATCCACGAT GGCTGAGACA ACCGAACCAC	60
CGTCGGACGC GGGGACATCG CAAGCCGACG CGATGGCGTT GGCCGCCGAA GCCGAAGCCG	120
CCGAAGCCGA AGCGCTGGCC GCCGCGGCGC GGGCCCGTGC CCGTGCCGCC CGGTTGAAGC	180
GTGAGGCGCT GGCGATGGCC CCAGCCGAGG ACGAGAACGT CCCCAGGAT ATGCAGACTG	240
GGAAGACGCC GAAGACTATG ACGACTATGA CGACTATGAG GCCGCAGACC AGGAGGCCGC	300
ACGGTCGGCA TCCTGGCGAC GCGGTTGCG GGTGCGGTTA CCAAGACTGT CCACGATTGC	360
CATGGCGGCC GCAGTCGTCA TCATCTGCGG CTTACCCGGG CTCAGCGGAT ACATTGTGTG	420
GCAACACCAT GAGGCCACCG AACGCCAGCA GCGCGCCGCG GCGTTCGCCG CCGGAGCCAA	480
GCAAGGTGTC ATCAACATGA CCTCGCTGGA CTTCAACAAG GCCAAAGAAG ACGTCGCGCG	540

TGTGATCGAC	AGCTCCACCG	GCGAATTCAG	GGATGACTTC	CAGCAGCGGG	CAGCCGATTT	600
CACCAAGGTT	GTCGAACAGT	CCAAAGTGGT	CACCGAAGGC	ACGGTGAACG	CGACAGCCGT	660
CGAATCCATG	AACGAGCATT	CCGCCGTGGT	GCTCGTCGCG	GCGACTTCAC	GGGTCACCAA	720
TTCCGCTGGG	GCGAAAGACG	AACCACGTGC	GTGGCGGCTC	AAAGTGACCG	TGACCGAAGA	780
GGGGGGACAG	TACAAGATGT	CGAAAGTTGA	GTTTCGTACCG	TGACCGATGA	CGTACGCGAC	840
GTCAACACCG	AAACCACTGA	CGCCACCGAA	GTCGCTGAGA	TCGACTCAGC	CGCAGGCGAA	900
GCCGGTGATT	CGGCGACCGA	GGCATTGTAC	ACCGACTCTG	CAACGGAATC	TACCGCGCAG	960
AAGGGTCAGC	GGCACCCTGA	CCTGTGGCGA	ATGCAGGTTA	CCTTGAAACC	CGTTCCGGTG	1020
ATTCTCATCC	TGCTCATGTT	GATCTCTGGG	GGCGCGACGG	GATGGCTATA	CCTTGAGCAA	1080
TACGACCCGA	TCAGCAGACG	GACTCCGGCG	CCGCCCGTGC	TGCCGTCGCC	GCGGCGTCTG	1140
ACGGGACAAT	CGCGCTGTTG	TGTATTACCC	CGACACGTCG	ACCAAGACTT	CGCTACCGCC	1200
AGGTGCGACC	TCGCCGGCGA	TTTCCTGTCC	TATACGACCA	GTTACGCGAG	CAGATCGTGG	1260
CTCCGGCGGC	CAAACAGAAG	TCACTGAAAA	CCACCGCCAA	GGTGGTGCGC	GCGGCCGTGT	1320
CGGAGCTACA	TCCGGATTCTG	GCCGTCGTTT	TGGTTTTTGT	CGACCAGAGC	ACTACCAGTA	1380
AGGACAGCCC	CAATCCGTCG	ATGGCGGCCA	GCAGCGTGAT	GGTGACCCTA	GCCAAGGTCTG	1440
ACGGCAATTG	GCTGATCACC	AAGTTCACCC	CGGTTTAGGT	TGCCGTAGGC	GGTCGCCAAG	1500
TCTGACGGGG	GCGCGGGTGG	CTGCTCGTGC	GAGATACCGG	CCGTTCTCCG	GACAATCACG	1560
GCCCGACCTC	AAACAGATCT	CGGCCGCTGT	CTAATCGGCC	GGGTTATTTA	AGATTAGTTG	1620
CCACTGTATT	TACCTGATGT	TCAGATTGTT	CAGCTGGATT	TAGCTTCGCG	GCAGGGCGGC	1680
TGGTGCACTT	TGCATCTGGG	GTTGTGACTA	CTTGAGAGAA	TTTGACCTGT	TGCCGACGTT	1740
GTTTGCTGTC	CATCATTGGT	GCTAGTTATG	GCCGAGCGGA	AGGATTATCG	AAGTGGTGGA	1800
CTTCGGGGCG	TTACCACCGG	AGATCAACTC	CGCGAGGATG	TACGCCGGCC	CGGGTTCGGC	1860
CTCGCTGGTG	GCCGCCGCGA	AGATGTGGGA	CAGCGTGCGG	AGTGACCTGT	TTTCGGCCGC	1920
GTCGGCGTTT	CAGTCGGTGG	TCTGGGGTCT	GACGACGGGA	TCGTGGATAG	GTTTCGTCGGC	1980
GGGTCTGATG	GTGGCGGCGG	CCTCGCCGTA	TGTGGCGTGG	ATGAGCGTCA	CCGCGGGGCA	2040
GGCCGAGCTG	ACGCCCGCCC	AGGTCCGGGT	TGCTGCGGCG	GCCTACGAGA	CGGCGTATGG	2100
GCTGACGGTG	CCCCCGCCGG	TGATCGCCGA	GAACCGTGCT	GAAGTATGA	TTCTGATAGC	2160
GACCAACCTC	TTGGGGCAAA	ACACCCCGGC	GATCGCGGTC	AACGAGGCCG	AATACGGGGA	2220
GATGTGGGCC	CAAGACGCCG	CCGCGATGTT	TGGCTACGCC	GCCACGGCGG	CGACGGCGAC	2280
CGAGGCGTTG	CTGCCGTTCTG	AGGACGCCCC	ACTGATCACC	AACCCCGGCG	GGCTCCTTGA	2340
GCAGGCCGTC	GCGGTCGAGG	AGGCCATCGA	CACCGCCGCG	GCGAACCAGT	TGATGAACAA	2400

```

TGTGCCCCAA GCGCTGCAAC AACTGGCCCA GCCCACGAAA AGCATCTGGC CGTTCGACCA      2460
ACTGAGTGAA CTCTGGAAAG CCATCTCGCC GCATCTGTGC CCGCTCAGCA ACATCGTGTC      2520
GATGCTCAAC AACCACGTGT CGATGACCAA CTCGGGTGTG TCGATGGCCA GCACCTTGCA      2580
CTCAATGTTG AAGGGCTTTG CTCCGGCGGC GGCTCAGGCC GTGGAAACCG CGGCGCAAAA      2640
CGGGGTCCAG GCGATGAGCT CGCTGGGCAG CCAGCTGGGT TCGTCGCTGG GTTCTTCGGG      2700
TCTGGGCGCT GGGGTGGCCG CCAACTTGGG TCGGGCGGCC TCGGTCGGTT CGTTGTCGGT      2760
GCCGCAGGCC TGGGCCGCGG CCAACCAGGC GGTACACCCG GCGGCGCGGG CGCTGCCGCT      2820
GACCAGCCTG ACCAGCGCCG CCCAAACCGC CCCC GGACAC ATGCTGGGCG GGCTACCGCT      2880
GGGGCAACTG ACCAATAGCG GCGGCGGGTT CGGCGGGGTT AGCAATGCGT TGCGGATGCC      2940
GCCGCGGGCG TACGTAATGC CCCGTGTGCC CGCCGCCGGG TAACGCCGAT CCGCACGCAA      3000
TGCGGGCCCT CTATCGGGGC AGCGATC                                          3027

```

(2) INFORMATION FOR SEQ ID NO:111:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 396 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:111:

```

Val Val Asp Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met
1           5           10           15
Tyr Ala Gly Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Lys Met Trp
20          25          30
Asp Ser Val Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser
35          40          45
Val Val Trp Gly Leu Thr Thr Gly Ser Trp Ile Gly Ser Ser Ala Gly
50          55          60
Leu Met Val Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr
65          70          75          80
Ala Gly Gln Ala Glu Leu Thr Ala Ala Gln Val Arg Val Ala Ala Ala
85          90          95
Ala Tyr Glu Thr Ala Tyr Gly Leu Thr Val Pro Pro Pro Val Ile Ala
100         105         110
Glu Asn Arg Ala Glu Leu Met Ile Leu Ile Ala Thr Asn Leu Leu Gly
115        120        125
Gln Asn Thr Pro Ala Ile Ala Val Asn Glu Ala Glu Tyr Gly Glu Met

```

135

130	135	140
Trp Ala Gln Asp Ala Ala Ala Met Phe Gly Tyr Ala Ala Thr Ala Ala		
145	150	155 160
Thr Ala Thr Glu Ala Leu Leu Pro Phe Glu Asp Ala Pro Leu Ile Thr		
	165	170 175
Asn Pro Gly Gly Leu Leu Glu Gln Ala Val Ala Val Glu Glu Ala Ile		
	180	185 190
Asp Thr Ala Ala Ala Asn Gln Leu Met Asn Asn Val Pro Gln Ala Leu		
	195	200 205
Gln Gln Leu Ala Gln Pro Thr Lys Ser Ile Trp Pro Phe Asp Gln Leu		
	210	215 220
Ser Glu Leu Trp Lys Ala Ile Ser Pro His Leu Ser Pro Leu Ser Asn		
	225	230 235 240
Ile Val Ser Met Leu Asn Asn His Val Ser Met Thr Asn Ser Gly Val		
	245	250 255
Ser Met Ala Ser Thr Leu His Ser Met Leu Lys Gly Phe Ala Pro Ala		
	260	265 270
Ala Ala Gln Ala Val Glu Thr Ala Ala Gln Asn Gly Val Gln Ala Met		
	275	280 285
Ser Ser Leu Gly Ser Gln Leu Gly Ser Ser Leu Gly Ser Ser Gly Leu		
	290	295 300
Gly Ala Gly Val Ala Ala Asn Leu Gly Arg Ala Ala Ser Val Gly Ser		
	305	310 315 320
Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln Ala Val Thr Pro		
	325	330 335
Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser Ala Ala Gln Thr		
	340	345 350
Ala Pro Gly His Met Leu Gly Gly Leu Pro Leu Gly Gln Leu Thr Asn		
	355	360 365
Ser Gly Gly Gly Phe Gly Gly Val Ser Asn Ala Leu Arg Met Pro Pro		
	370	375 380
Arg Ala Tyr Val Met Pro Arg Val Pro Ala Ala Gly		
	385	390 395

(2) INFORMATION FOR SEQ ID NO:112:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1616 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

CATCGGAGGG AGTGATCACC ATGCTGTGGC ACGCAATGCC ACCGGAGTAA ATACCGCACG	60
GCTGATGGCC GGC GCGGGTTC CGGCTCCAAT GCTTGCGGCG GCCGCGGGAT GGCAGACGCT	120
TTCGGCGGCT CTGGACGCTC AGGCCGTCGA GTTGACCGCG CGCCTGAACT CTCTGGGAGA	180
AGCCTGGACT GGAGGTGGCA GCGACAAGGC GCTTGCGGCT GCAACGCCGA TGGTGGTCTG	240
GCTACAAACC GCGTCAACAC AGGCCAAGAC CCGTGCGATG CAGGCGACGG CGCAAGCCGC	300
GGCATAACACC CAGGCCATGG CCACGACGCC GTCGCTGCCG GAGATCGCCG CCAACCACAT	360
CACCCAGGCC GTCCTTACGG CCACCAACTT CTTGCGTATC AACACGATCC CGATCGCGTT	420
GACCGAGATG GATTATTTCA TCCGTATGTG GAACCAGGCA GCCCTGGCAA TGGAGGTCTA	480
CCAGGCCGAG ACCGCGGTTA ACACGCTTTT CGAGAAGCTC GAGCCGATGG CGTCGATCCT	540
TGATCCCGGC GCGAGCCAGA GCACGACGAA CCCGATCTTC GGAATGCCCT CCCCTGGCAG	600
CTCAACACCG GTTGGCCAGT TGCCGCCGGC GGCTACCCAG ACCCTCGGCC AACTGGGTGA	660
GATGAGCGGC CCGATGCAGC AGCTGACCCA GCCGCTGCAG CAGGTGACGT CGTTGTTCAG	720
CCAGGTGGGC GGCACCGGCG GCGGCAACCC AGCCGACGAG GAAGCCGCGC AGATGGGCCT	780
GCTCGGCACC AGTCCGCTGT CGAACCATCC GCTGGCTGGT GGATCAGGCC CCAGCGCGGG	840
CGCGGGCCTG CTGCGCGCGG AGTCGCTACC TGGCGCAGGT GGGTCGTTGA CCCGCACGCC	900
GCTGATGTCT CAGCTGATCG AAAAGCCGGT TGCCCCCTCG GTGATGCCGG CGGCTGCTGC	960
CGGATCGTCG GCGACGGGTG GCGCCGCTCC GGTGGGTGCG GGAGCGATGG GCCAGGGTGC	1020
GCAATCCGGC GGCTCCACCA GGCCGGGTCT GGTGCGGCCG GCACCGCTCG CGCAGGAGCG	1080
TGAAGAAGAC GACGAGGACG ACTGGGACGA AGAGGACGAC TGGTGAGCTC CCGTAATGAC	1140
AACAGACTTC CCGGCCACCC GGGCCGGAAG ACTTGCCAAC ATTTTGGCGA GGAAGGTAAA	1200
GAGAGAAAGT AGTCCAGCAT GGCAGAGATG AAGACCGATG CCGCTACCCT CGCGCAGGAG	1260
GCAGGTAATT TCGAGCGGAT CTCCGGCGAC CTGAAAACCC AGATCGACCA GGTGGAGTCG	1320
ACGGCAGGTT CGTTGCAGGG CCAGTGCGC GGC GCGGCGG GGACGGCCGC CCAGGCCGCG	1380
GTGGTGCGCT TCCAAGAAGC AGCCAATAAG CAGAAGCAGG AACTCGACGA GATCTCGACG	1440
AATATTCGTC AGGCCGGCGT CCAATACTCG AGGGCCGACG AGGAGCAGCA GCAGGCGCTG	1500
TCCTCGCAA TGGGCTTCTG ACCCGCTAAT ACGAAAAGAA ACGGAGCAA AACATGACAG	1560
AGCAGCAGTG GAATTCGCG GGTATCGAGG CCGCGGCAAG CGCAATCCAG GGAAAT	1616

(2) INFORMATION FOR SEQ ID NO:113:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 432 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

```

CTAGTGGATG GGACCATGGC CATTTTCTGC AGTCTCACTG CCTTCTGTGT TGACATTTTG      60
GCACGCCGGC GGAAACGAAG CACTGGGGTC GAAGAACGGC TCGCTGCCA TATCGTCCGG      120
AGCTTCCATA CCTTCGTGCG GCCGGAAGAG CTTGTCTAG TCGGCCGCCA TGACAACCTC      180
TCAGAGTGCG CTCAAACGTA TAAACACGAG AAAGGGCGAG ACCGACGGAA GGTCGAACTC      240
GCCCCGATCCC GTGTTTCGCT ATTCTACGCG AACTCGGCGT TGCCCTATGC GAACATCCCA      300
GTGACGTTGC CTTGGGTCGA AGCCATTGCC TGACCGGCTT CGCTGATCGT CCGCGCCAGG      360
TTCTGCAGCG CGTTGTTTCAG CTCGGTAGCC GTGGCGTCCC ATTTTGTCTG GACACCCTGG      420
TACGCCTCCG AA                                          432

```

(2) INFORMATION FOR SEQ ID NO:114:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 368 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

```

Met Leu Trp His Ala Met Pro Pro Glu Xaa Asn Thr Ala Arg Leu Met
1           5           10           15

Ala Gly Ala Gly Pro Ala Pro Met Leu Ala Ala Ala Ala Gly Trp Gln
20           25           30

Thr Leu Ser Ala Ala Leu Asp Ala Gln Ala Val Glu Leu Thr Ala Arg
35           40           45

Leu Asn Ser Leu Gly Glu Ala Trp Thr Gly Gly Gly Ser Asp Lys Ala
50           55           60

Leu Ala Ala Ala Thr Pro Met Val Val Trp Leu Gln Thr Ala Ser Thr
65           70           75           80

Gln Ala Lys Thr Arg Ala Met Gln Ala Thr Ala Gln Ala Ala Ala Tyr
85           90           95

Thr Gln Ala Met Ala Thr Thr Pro Ser Leu Pro Glu Ile Ala Ala Asn
100          105          110

His Ile Thr Gln Ala Val Leu Thr Ala Thr Asn Phe Phe Gly Ile Asn
115          120          125

```

138

```

Thr Ile Pro Ile Ala Leu Thr Glu Met Asp Tyr Phe Ile Arg Met Trp
130                      135                      140

Asn Gln Ala Ala Leu Ala Met Glu Val Tyr Gln Ala Glu Thr Ala Val
145                      150                      155                      160

Asn Thr Leu Phe Glu Lys Leu Glu Pro Met Ala Ser Ile Leu Asp Pro
165                      170                      175

Gly Ala Ser Gln Ser Thr Thr Asn Pro Ile Phe Gly Met Pro Ser Pro
180                      185                      190

Gly Ser Ser Thr Pro Val Gly Gln Leu Pro Pro Ala Ala Thr Gln Thr
195                      200                      205

Leu Gly Gln Leu Gly Glu Met Ser Gly Pro Met Gln Gln Leu Thr Gln
210                      215                      220

Pro Leu Gln Gln Val Thr Ser Leu Phe Ser Gln Val Gly Gly Thr Gly
225                      230                      235                      240

Gly Gly Asn Pro Ala Asp Glu Glu Ala Ala Gln Met Gly Leu Leu Gly
245                      250                      255

Thr Ser Pro Leu Ser Asn His Pro Leu Ala Gly Gly Ser Gly Pro Ser
260                      265                      270

Ala Gly Ala Gly Leu Leu Arg Ala Glu Ser Leu Pro Gly Ala Gly Gly
275                      280                      285

Ser Leu Thr Arg Thr Pro Leu Met Ser Gln Leu Ile Glu Lys Pro Val
290                      295                      300

Ala Pro Ser Val Met Pro Ala Ala Ala Ala Gly Ser Ser Ala Thr Gly
305                      310                      315                      320

Gly Ala Ala Pro Val Gly Ala Gly Ala Met Gly Gln Gly Ala Gln Ser
325                      330                      335

Gly Gly Ser Thr Arg Pro Gly Leu Val Ala Pro Ala Pro Leu Ala Gln
340                      345                      350

Glu Arg Glu Glu Asp Asp Glu Asp Asp Trp Asp Glu Glu Asp Asp Trp
355                      360                      365

```

(2) INFORMATION FOR SEQ ID NO:115:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 100 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

```

Met Ala Glu Met Lys Thr Asp Ala Ala Thr Leu Ala Gln Glu Ala Gly
1           5           10           15

```


[illegible]

(2) INFORMATION FOR SEQ ID NO:116:

(i) SEQUENCE CHARACTERISTICS:

- SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 396 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

GATCTCCGGC	GACCTGAAAA	CCCAGATCGA	CCAGGTGGAG	TCGACGGCAG	GTTCTGTTGCA	60
GGGCCAGTGG	CGCGGCGCGG	CGGGGACGGC	CGCCCAGGCC	GCGGTGGTGC	GCTTCCAAGA	120
AGCAGCCAAT	AAGCAGAAGC	AGGAACTCGA	CGAGATCTCG	ACGAATATTC	GTCAGGCCGG	180
CGTCCAATAC	TCGAGGGCCG	ACGAGGAGCA	GCAGCAGGCG	CTGTCTCTGC	AAATGGGCTT	240
CTGACCCGCT	AATACGAAAA	GAAACGGAGC	AAAAACATGA	CAGAGCAGCA	GTGGAATTTT	300
GCGGGTATCG	AGGCCGCGGC	AAGCGCAATC	CAGGGAAATG	TCACGTCCAT	TCATTCCCTC	360
CTTGACGAGG	GGAAGCAGTC	CCTGACCAAG	CTCGCA			396

(2) INFORMATION FOR SEQ ID NO:117:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 80 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

Ile Ser Gly Asp Leu Lys Thr Gln Ile Asp Gln Val Glu Ser Thr Ala

140

1	5	10	15
Gly Ser Leu Gln Gly Gln Trp Arg Gly Ala Ala Gly Thr Ala Ala Gln	20	25	30
Ala Ala Val Val Arg Phe Gln Glu Ala Ala Asn Lys Gln Lys Gln Glu	35	40	45
Leu Asp Glu Ile Ser Thr Asn Ile Arg Gln Ala Gly Val Gln Tyr Ser	50	55	60
Arg Ala Asp Glu Glu Gln Gln Gln Ala Leu Ser Ser Gln Met Gly Phe	65	70	75
			80

(2) INFORMATION FOR SEQ ID NO:118:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 387 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

.(xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

GTGGATCCCG ATCCCGTGTT TCGCTATTCT ACGCGAACTC GCGGTTGCCC TATGCGAACA	60
TCCCAGTGAC GTTGCCCTTCG GTCGAAGCCA TTGCCTGACC GGCTTCGCTG ATCGTCCGCG	120
CCAGGTTCTG CAGCGCGTTG TTCAGCTCGG TAGCCGTGGC GTCCCATTTT TGCTGGACAC	180
CCTGGTACGC CTCCGAACCG CTACCGCCCC AGGCCGCTGC GAGCTTGGTC AGGGACTGCT	240
TCCCCTCGTC AAGGAGGGAA TGAATGGACG TGACATTTCC CTGGATTGCG CTTGCCGCGG	300
CCTCGATACC CGCGAAATTC CACTGCTGCT CTGTCATGTT TTTGCTCCGT TTCTTTTCGT	360
ATTAGCGGGT CAGAAGCCCA TTTGCGA	387

(2) INFORMATION FOR SEQ ID NO:119:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 272 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:119:

CGGCACGAGG ATCTCGGTTG GCCCAACGGC GCTGGCGAGG GCTCCGTTCC GGGGGCGAGC	60
TGCGCGCCGG ATGCTTCCTC TGCCCGCAGC CGCGCCTGGA TGGATGGACC AGTTGCTACC	120
TTCCCGACGT TTCGTTCCGT GTCTGTGCGA TAGCGGTGAC CCCGGCGCGC ACGTCGGGAG	180

TGTTGGGGGG CAGGCCGGGT CGGTGGTTCG GCCGGGGACG CAGACGGTCT GGACGGAACG 240
GGCGGGGGTT CGCCGATTGG CATCTTTGCC CA 272

(2) INFORMATION FOR SEQ ID NO:120:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:120:

Asp Pro Val Asp Ala Val Ile Asn Thr Thr Cys Asn Tyr Gly Gln Val
1 5 10 15
Val Ala Ala Leu
20

(2) INFORMATION FOR SEQ ID NO:121:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:121:

Ala Val Glu Ser Gly Met Leu Ala Leu Gly Thr Pro Ala Pro Ser
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:122:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:122:

Ala Ala Met Lys Pro Arg Thr Gly Asp Gly Pro Leu Glu Ala Ala Lys
1 5 10 15
Glu Gly Arg

(2) INFORMATION FOR SEQ ID NO:123:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:123:

Tyr Tyr Trp Cys Pro Gly Gln Pro Phe Asp Pro Ala Trp Gly Pro
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:124:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:124:

Asp Ile Gly Ser Glu Ser Thr Glu Asp Gln Gln Xaa Ala Val
1 5 10

(2) INFORMATION FOR SEQ ID NO:125:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:125:

Ala Glu Glu Ser Ile Ser Thr Xaa Glu Xaa Ile Val Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:126:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:126:

Asp Pro Glu Pro Ala Pro Pro Val Pro Thr Thr Ala Ala Ser Pro Pro
1 5 10 15
Ser

(2) INFORMATION FOR SEQ ID NO:127:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:127:

Ala Pro Lys Thr Tyr Xaa Glu Glu Leu Lys Gly Thr Asp Thr Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:128:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:128:

Asp Pro Ala Ser Ala Pro Asp Val Pro Thr Ala Ala Gln Leu Thr Ser
1 5 10 15
Leu Leu Asn Ser Leu Ala Asp Pro Asn Val Ser Phe Ala Asn
20 25 30

(2) INFORMATION FOR SEQ ID NO:129:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:129:

Asp Pro Pro Asp Pro His Gln Xaa Asp Met Thr Lys Gly Tyr Tyr Pro
1 5 10 15

Gly Gly Arg Arg Xaa Phe
20

(2) INFORMATION FOR SEQ ID NO:130:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:130:

Asp Pro Gly Tyr Thr Pro Gly
1 5

(2) INFORMATION FOR SEQ ID NO:131:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: /note= "The Second Residue Can Be Either a Pro or Thr"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:131:

Xaa Xaa Gly Phe Thr Gly Pro Gln Phe Tyr
1 5 10

(2) INFORMATION FOR SEQ ID NO:132:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: /note= "The Third Residue Can Be Either a Gln or Leu"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:132:

Xaa Pro Xaa Val Thr Ala Tyr Ala Gly
1 5

(2) INFORMATION FOR SEQ ID NO:133:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:133:

Xaa Xaa Xaa Glu Lys Pro Phe Leu Arg
1 5

(2) INFORMATION FOR SEQ ID NO:134:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:134:

Xaa Asp Ser Glu Lys Ser Ala Thr Ile Lys Val Thr Asp Ala Ser
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:135:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:135:

Ala Gly Asp Thr Xaa Ile Tyr Ile Val Gly Asn Leu Thr Ala Asp
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:136:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:136:

Ala Pro Glu Ser Gly Ala Gly Leu Gly Gly Thr Val Gln Ala Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:137:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:137:

Xaa Tyr Ile Ala Tyr Xaa Thr Thr Ala Gly Ile Val Pro Gly Lys Ile
1 5 10 15
Asn Val His Leu Val
20

(2) INFORMATION FOR SEQ ID NO:138:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 882 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:138:

GCAACGCTGT CGTGGCCTTT GCGGTGATCG GTTTCGCCTC GCTGGCGGTG GCGGTGGCGG	60
TCACCATCCG ACCGACCGCG GCCTCAAAAC CGGTAGAGGG ACACCAAAAC GCCCAGCCAG	120
GGAAGTTCAT GCCGTTGTTG CCGACGCAAC AGCAGGCGCC GGTCCCGCCG CCTCCGCCCCG	180
ATGATCCCAC CGCTGGATTC CAGGGCGGCA CCATTCCGGC TGTACAGAAC GTGGTGCCGC	240
GGCCGGGTAC CTCACCCGGG GTGGGTGGGA CGCCGGCTTC GCCTGCGCCG GAAGCGCCGG	300
CCGTGCCCCG TGTTGTGCCT GCCCCGGTGC CAATCCCGGT CCCGATCATC ATTCCCCCGT	360
TCCCGGGTTG GCAGCCTGGA ATGCCGACCA TCCCCACCGC ACCGCCGACG ACGCCGGTGA	420
CCACGTCCGC GACGACGCCG CCGACCACGC CGCCGACCAC GCCGGTGACC ACGCCGCCAA	480
CGACGCCGCC GACCACGCCG GTGACCACGC CGCCAACGAC GCCGCCGACC ACGCCGGTGA	540
CCACGCCACC AACGACCGTC GCCCCGACGA CCGTCGCCCC GACGACGGTC GCTCCGACCA	600
CCGTGCCCC GACCACGGTC GCTCCAGCCA CCGCCACGCC GACGACCGTC GCTCCGCAGC	660

.147

CGACGCAGCA GCCCAGCAA CAACCAACCC AACAGATGCC AACCCAGCAG CAGACCGTGG 720
 CCCCAGAC GGTGGCGCCG GCTCCGCAGC CGCCGTCCGG TGGCCGCAAC GGCAGCGGCG 780
 GGGGCGACTT ATTCGGCGGG TTCTGATCAC GGTCGCGGCT TCACTACGGT CGGAGGACAT 840
 GGCCGGTGAT GCGGTGACGG TGGTGCTGCC CTGTCTCAAC GA 882

(2) INFORMATION FOR SEQ ID NO:139:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 815 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:139:

CCATCAACCA ACCGCTCGCG CCGCCCGCGC CGCCGGATCC GCCGTGCGCG CCACGCCCCG 60
 CGGTGCCTCC GGTGCCCCCG TTGCCGCCGT CGCCGCCGTC GCCGCCGACC GGCTGGGTGC 120
 CTAGGGCGCT GTTACGCCC TGTTGGCGG GGACGCCGCC GGCACCACCG GTACCGCCGA 180
 TGGCGCCGTT GCCGCCGGCG GCACCGTTGC CACCGTTGCC ACCGTTGCCA CCGTTGCCGA 240
 CCAGCCACCC GCCGCGACCA CCGGCACCGC CGGCGCCGCC CGCACCGCCG GCGTGCCCGT 300
 TCGTGCCCGT ACCGCCGGCA CCGCCGTTGC CGCCGTCACC GCCGACGGAA CTACCGGCGG 360
 ACGCGGCCTG CCCGCCGGCG CCGCCCGCAC CGCCATTGGC ACCGCCGTCA CCGCCGGCTG 420
 GGAGTGCCG GATTAGGGCA CTGACCGGCG CAACCAGCGC AAGTACTCTC GGTCACCGAG 480
 CACTTCCAGA CGACACCACA GCACGGGGTT GTCGGCGGAC TGGGTGAAAT GGCAGCCGAT 540
 AGCGGCTAGC TGTCGGCTGC GGTCAACCTC GATCATGATG TCGAGGTGAC CGTGACCGCG 600
 CCCCCGAAG GAGGCGCTGA ACTCGGCGTT GAGCCGATCG GCGATCGGTT GGGGCAGTGC 660
 CCAGGCCAAT ACGGGGATAC CGGGTGTCNA AGCCGCCGCG AGCGCAGCTT CGGTTGCGCG 720
 ACNGTGGTCG GGGTGGCCTG TTACGCCGTT GTCNTCGAAC ACGAGTAGCA GGTCTGCTCC 780
 GGCGAGGGCA TCCACCACGC GTTGCCTCAG CTCGT 815

(2) INFORMATION FOR SEQ ID NO:140:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1152 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:140:

```

ACCAGCCGCC GGCTGAGGTC TCAGATCAGA GAGTCTCCGG ACTCACC GGG GCGGTT CAGC      60
CTTCTCCCAG AACAACTGCT GAAGATCCTC GCCC GCGAAA CAGGCGCTGA TTTGACGCTC      120
TATGACCGGT TGAACGACGA GATCATCCGG CAGATTGATA TGGCACCGCT GGGCTAACAG      180
GTGCGCAAGA TGGTGACGCT GTATGTCTCG GACTCCGTGT CGCGGATCAG CTTTGCCGAC      240
GGCCGGGTGA TCGTGTGGAG CGAGGAGCTC GGCGAGAGCC AGTATCCGAT CGAGACGCTG      300
GACGGCATCA CGCTGTTTGG GCGGCCGACG ATGACAACGC CCTTCATCGT TGAGATGCTC      360
AAGCGTGAGC GCGACATCCA GCTCTTCACG ACCGACGGCC ACTACCAGGG CCGGATCTCA      420
ACACCCGACG TGTCATACGC GCCGCGGCTC CGTCAGCAAG TTCACCGCAC CGACGATCCT      480
GCGTTCTGCC TGTCGTTAAG CAAGCGGATC GTGTCGAGGA AGATCCTGAA TCAGCAGGCC      540
TTGATTCGGG CACACACGTC GGGGCAAGAC GTTGCTGAGA GCATCCGCAC GATGAAGCAC      600
TCGCTGGCCT GGGTCGATCG ATCGGGCTCC CTGGCGGAGT TGAACGGGTT CGAGGGAAAT      660
GCCGCAAAGG CATACTTCAC CGCGCTGGGG CATCTCGTCC CGCAGGAGTT CGCATTCCAG      720
GGCCGCTCGA CTCGGCCGCC GTTGGACGCC TTCAACTCGA TGGTCAGCCT CGGCTATTCTG      780
CTGCTGTACA AGAACATCAT AGGGGCGATC GAGCGTCACA GCCTGAACGC GTATATCGGT      840
TTCCTACACC AGGATTTCACG AGGGCACGCA ACGTCTCGTG CCGAATTCGG CACGAGCTCC      900
GCTGAAACCG CTGGCCGGCT GCTCAGTGCC CGTACGTAAT CCGCTGCGCC CAGGCCGGCC      960
CGCCGGCCGA ATACCAGCAG ATCGGACAGC GAATTGCCGC CCAGCCGGTT GGAGCCGTGC     1020
ATACCGCCGG CAACTCACC GGCAGCGAAC AGGCCTGGCA CCGTGGCGGC GCCGGTGTCC     1080
GCGTCTACTT CGACACCGCC CATCACGTAG TGACACGTCG GCCCGACTTC CATTGCCTGC     1140
GTTCCGGCACG AG                                                         1152

```

(2) INFORMATION FOR SEQ ID NO:141:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 655 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:141:

```

CTCGTGCCGA TTCGGCAGGG TGTA CTTGCC GGTGGTGTAN GCCGCATGAG TGCCGACGAC      60
CAGCAATGCG GCAACAGCAC GGATCCCGGT CAACGACGCC ACCCGGTCCA CGTGGGCGAT      120

```

```

CCGCTCGAGT CCGCCCTGGG CGGCTCTTTC CTTGGGCAGG GTCATCCGAC GTGTTTCCGC      180
CGTGGTTTGC CGCCATTATG CCGGCGCGCC GCGTCGGGCG GCCGGTATGG CCGAANGTCG      240
ATCAGCACAC CCGAGATACG GGTCTGTGCA AGCTTTTTTGA GCGTCGCGCG GGGCAGCTTC      300
GCCGGCAATT CTACTAGCGA GAAGTCTGGC CCGATACGGA TCTGACCGAA GTCGCTGCGG      360
TGCAGCCCAC CCTCATTGGC GATGGCGCCG ACGATGGCGC CTGGACCGAT CTTGTGCCGC      420
TTGCCGACGG CGACGCGGTA GGTGGTCAAG TCCGGTCTAC GCTTGGGCCT TTGCGGACGG      480
TCCCGACGCT GGTGCGGGTT GCGCCGCGAA AGCGGCGGGT CGGGTGCCAT CAGGAATGCC      540
TCACCGCCGC GGCAGTGCAC GGCCAGTGCC GCGGCGATGT CAGCCATCGG GACATCATGC      600
TCGCGTTCAT ACTCCTCGAC CAGTCGGCGG AACAGCTCGA TTCCCGGACC GCCCA          655

```

(2) INFORMATION FOR SEQ ID NO:142:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 267 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:142:

```

Asn Ala Val Val Ala Phe Ala Val Ile Gly Phe Ala Ser Leu Ala Val
1           5           10           15
Ala Val Ala Val Thr Ile Arg Pro Thr Ala Ala Ser Lys Pro Val Glu
20          25          30
Gly His Gln Asn Ala Gln Pro Gly Lys Phe Met Pro Leu Leu Pro Thr
35          40          45
Gln Gln Gln Ala Pro Val Pro Pro Pro Pro Pro Asp Asp Pro Thr Ala
50          55          60
Gly Phe Gln Gly Gly Thr Ile Pro Ala Val Gln Asn Val Val Pro Arg
65          70          75          80
Pro Gly Thr Ser Pro Gly Val Gly Gly Thr Pro Ala Ser Pro Ala Pro
85          90          95
Glu Ala Pro Ala Val Pro Gly Val Val Pro Ala Pro Val Pro Ile Pro
100         105         110
Val Pro Ile Ile Ile Pro Pro Phe Pro Gly Trp Gln Pro Gly Met Pro
115        120        125
Thr Ile Pro Thr Ala Pro Pro Thr Thr Pro Val Thr Thr Ser Ala Thr
130        135        140
Thr Pro Pro Thr Thr Pro Pro Thr Thr Pro Val Thr Thr Pro Pro Thr

```

150

145		150		155		160
Thr	Pro	Pro	Thr	Thr	Pro	Thr
		165		170		175
Thr	Pro	Val	Thr	Thr	Val	Ala
		180		185		190
Pro	Thr	Thr	Val	Ala	Pro	Thr
		195		200		205
Ala	Thr	Ala	Thr	Pro	Thr	Thr
		210		215		220
Thr	Gln	Gln	Pro	Thr	Gln	Gln
			230		235	
Pro	Gln	Thr	Val	Ala	Pro	Ala
			245		250	
Gly	Ser	Gly	Gly	Gly	Asp	Leu
			260			

(2) INFORMATION FOR SEQ ID NO:143:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 174 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:143:

Ile	Asn	Gln	Pro	Leu	Ala	Pro	Pro	Ala	Pro	Pro	Asp	Pro	Pro	Ser	Pro
1				5				10						15	
Pro	Arg	Pro	Pro	Val	Pro	Pro	Val	Pro	Pro	Leu	Pro	Pro	Ser	Pro	Pro
			20				25					30			
Ser	Pro	Pro	Thr	Gly	Trp	Val	Pro	Arg	Ala	Leu	Leu	Pro	Pro	Trp	Leu
		35				40					45				
Ala	Gly	Thr	Pro	Pro	Ala	Pro	Pro	Val	Pro	Pro	Met	Ala	Pro	Leu	Pro
		50			55						60				
Pro	Ala	Ala	Pro	Leu	Pro	Pro	Leu	Pro	Pro	Leu	Pro	Pro	Leu	Pro	Thr
		65			70					75					80
Ser	His	Pro	Pro	Arg	Pro	Pro	Ala	Pro	Pro	Ala	Pro	Pro	Ala	Pro	Pro
				85				90						95	
Ala	Cys	Pro	Phe	Val	Pro	Val	Pro	Pro	Ala	Pro	Pro	Leu	Pro	Pro	Ser
			100				105						110		
Pro	Pro	Thr	Glu	Leu	Pro	Ala	Asp	Ala	Ala	Cys	Pro	Pro	Ala	Pro	Pro
			115				120						125		

151

Ala Pro Pro Leu Ala Pro Pro Ser Pro Pro Ala Gly Ser Ala Ala Ile
 130 135 140

Arg Ala Leu Thr Gly Ala Thr Ser Ala Ser Thr Leu Gly His Arg Ala
 145 150 155 160

Leu Pro Asp Asp Thr Thr Ala Arg Gly Cys Arg Arg Thr Gly
 165 170

(2) INFORMATION FOR SEQ ID NO:144:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:144:

Gln Pro Pro Ala Glu Val Ser Asp Gln Arg Val Ser Gly Leu Thr Gly
 1 5 10 15

Ala Val Gln Pro Ser Pro Arg Thr Thr Ala Glu Asp Pro Arg Pro Arg
 20 25 30

Asn Arg Arg
 35

(2) INFORMATION FOR SEQ ID NO:145:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 104 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:145:

Arg Ala Asp Ser Ala Gly Cys Thr Cys Arg Trp Cys Xaa Pro His Glu
 1 5 10 15

Cys Arg Arg Pro Ala Met Arg Gln Gln His Gly Ser Arg Ser Thr Thr
 20 25 30

Pro Pro Gly Pro Arg Gly Arg Ser Ala Arg Val Arg Pro Gly Arg Leu
 35 40 45

Phe Pro Trp Ala Gly Ser Ser Asp Val Phe Pro Pro Trp Phe Ala Ala
 50 55 60

Ile Met Pro Ala Arg Arg Val Gly Arg Pro Val Trp Pro Xaa Val Asp
 65 70 75 80

152

Gln His Thr Arg Asp Thr Gly Leu Cys Lys Leu Phe Glu Arg Arg Ala
85 90 95
Gly Gln Leu Arg Arg Gln Phe Tyr
100

(2) INFORMATION FOR SEQ ID NO:146:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 53 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "PCR primer"
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:146:

GGATCCATAT GGGCCATCAT CATCATCATC ACGTGATCGA CATCATCGGG ACC

53

(2) INFORMATION FOR SEQ ID NO:147:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "PCR Primer"
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:147:

CCTGAATTCA GGCCTCGGTT GCGCCGGCCT CATCTTGAAC GA

42

(2) INFORMATION FOR SEQ ID NO:148:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "PCR Primer"
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:148:

GGATCCTGCA GGCTCGAAAC CACCGAGCGG T

31

(2) INFORMATION FOR SEQ ID NO:149:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:149:

CTCTGAATTC AGCGCTGGAA ATCGTCGCGA T

31

(2) INFORMATION FOR SEQ ID NO:150:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:150:

GGATCCAGCG CTGAGATGAA GACCGATGCC GCT

33

(2) INFORMATION FOR SEQ ID NO:151:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR primer"

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mycobacterium tuberculosis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:151:

GAGAGAATTC TCAGAAGCCC ATTTGCGAGG ACA

33

(2) INFORMATION FOR SEQ ID NO:152:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1993 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Mycobacterium tuberculosis

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 152..1273

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:152:

TGTTCTTCGA CGGCAGGCTG GTGGAGGAAG GGCCACCGA ACAGCTGTTT TCCTCGCCGA	60
AGCATGCGGA AACCGCCCGA TACGTCGCCG GACTGTCGGG GGACGTCAAG GACGCCAAGC	120
GCGGAAATTG AAGAGCACAG AAAGGTATGG C GTG AAA ATT CGT TTG CAT ACG	172
Val Lys Ile Arg Leu His Thr	
1 5	
CTG TTG GCC GTG TTG ACC GCT GCG CCG CTG CTG CTA GCA GCG GCG GGC	220
Leu Leu Ala Val Leu Thr Ala Ala Pro Leu Leu Leu Ala Ala Ala Gly	
10 15 20	
TGT GGC TCG AAA CCA CCG AGC GGT TCG CCT GAA ACG GGC GCC GGC GCC	268
Cys Gly Ser Lys Pro Pro Ser Gly Ser Pro Glu Thr Gly Ala Gly Ala	
25 30 35	
GGT ACT GTC GCG ACT ACC CCC GCG TCG TCG CCG GTG ACG TTG GCG GAG	316
Gly Thr Val Ala Thr Thr Pro Ala Ser Ser Pro Val Thr Leu Ala Glu	
40 45 50 55	
ACC GGT AGC ACG CTG CTC TAC CCG CTG TTC AAC CTG TGG GGT CCG GCC	364
Thr Gly Ser Thr Leu Leu Tyr Pro Leu Phe Asn Leu Trp Gly Pro Ala	
60 65 70	
TTT CAC GAG AGG TAT CCG AAC GTC ACG ATC ACC GCT CAG GGC ACC GGT	412
Phe His Glu Arg Tyr Pro Asn Val Thr Ile Thr Ala Gln Gly Thr Gly	
75 80 85	
TCT GGT GCC GGG ATC GCG CAG GCC GCC GCC GGG ACG GTC AAC ATT GGG	460
Ser Gly Ala Gly Ile Ala Gln Ala Ala Ala Gly Thr Val Asn Ile Gly	
90 95 100	
GCC TCC GAC GCC TAT CTG TCG GAA GGT GAT ATG GCC GCG CAC AAG GGG	508
Ala Ser Asp Ala Tyr Leu Ser Glu Gly Asp Met Ala Ala His Lys Gly	
105 110 115	
CTG ATG AAC ATC GCG CTA GCC ATC TCC GCT CAG CAG GTC AAC TAC AAC	556
Leu Met Asn Ile Ala Leu Ala Ile Ser Ala Gln Gln Val Asn Tyr Asn	
120 125 130 135	

[illegible]

GCATGCTGGC CCGTGCCGGT GAAGTCGGCC GCGCTGGCCC GGCCATCCGG TGGTTGGGTG 1393
 GGATAGGTGC GGTGATCCCG CTGCTTGCGC TGGTCTTGGT GCTGGTGGTG CTGGTCATCG 1453
 AGGCGATGGG TGCATCAGG CTCAACGGGT TGCATTTCTT CACCGCCACC GAATGGAATC 1513
 CAGGCAACAC CTACGGCGAA ACCGTTGTCA CCGACGCGTC GCCCATCCGG TCGGCGCCTA 1573
 CTACGGGGCG TTGCCGCTGA TCGTCGGGAC GCTGGCGACC TCGGCAATCG CCCTGATCAT 1633
 CGCGGTGCCG GTCTCTGTAG GAGCGGCGCT GGTGATCGTG GAACGGCTGC CGAAACGGTT 1693
 GGCCGAGGCT GTGGGAATAG TCCTGGAATT GCTCGCCGGA ATCCCCAGCG TGGTCGTCGG 1753
 TTTGTGGGGG GCAATGACGT TCGGGCCGTT CATCGCTCAT CACATCGCTC CGGTGATCGC 1813
 TCACAACGCT CCCGATGTGC CGGTGCTGAA CTACTTGCGC GGCGACCCGG GCAACGGGGA 1873
 GGGCATGTTG GTGTCCGGTC TGGTGTGGC GGTGATGGTC GTTCCCATTA TCGCCACCAC 1933
 CACTCATGAC CTGTTCCGGC AGGTGCCGGT GTTGCCCCGG GAGGGCGCGA TCGGGAATTC 1993

(2) INFORMATION FOR SEQ ID NO:153:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 374 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:153:

Val Lys Ile Arg Leu His Thr Leu Leu Ala Val Leu Thr Ala Ala Pro
 1 5 10 15
 Leu Leu Leu Ala Ala Ala Gly Cys Gly Ser Lys Pro Pro Ser Gly Ser
 20 25 30
 Pro Glu Thr Gly Ala Gly Ala Gly Thr Val Ala Thr Thr Pro Ala Ser
 35 40 45
 Ser Pro Val Thr Leu Ala Glu Thr Gly Ser Thr Leu Leu Tyr Pro Leu
 50 55 60
 Phe Asn Leu Trp Gly Pro Ala Phe His Glu Arg Tyr Pro Asn Val Thr
 65 70 75 80
 Ile Thr Ala Gln Gly Thr Gly Ser Gly Ala Gly Ile Ala Gln Ala Ala
 85 90 95
 Ala Gly Thr Val Asn Ile Gly Ala Ser Asp Ala Tyr Leu Ser Glu Gly
 100 105 110
 Asp Met Ala Ala His Lys Gly Leu Met Asn Ile Ala Leu Ala Ile Ser
 115 120 125
 Ala Gln Gln Val Asn Tyr Asn Leu Pro Gly Val Ser Glu His Leu Lys
 130 135 140

157

Leu Asn Gly Lys Val Leu Ala Ala Met Tyr Gln Gly Thr Ile Lys Thr
 145 150 155 160
 Trp Asp Asp Pro Gln Ile Ala Ala Leu Asn Pro Gly Val Asn Leu Pro
 165 170 175
 Gly Thr Ala Val Val Pro Leu His Arg Ser Asp Gly Ser Gly Asp Thr
 180 185 190
 Phe Leu Phe Thr Gln Tyr Leu Ser Lys Gln Asp Pro Glu Gly Trp Gly
 195 200 205
 Lys Ser Pro Gly Phe Gly Thr Thr Val Asp Phe Pro Ala Val Pro Gly
 210 215 220
 Ala Leu Gly Glu Asn Gly Asn Gly Gly Met Val Thr Gly Cys Ala Glu
 225 230 235 240
 Thr Pro Gly Cys Val Ala Tyr Ile Gly Ile Ser Phe Leu Asp Gln Ala
 245 250 255
 Ser Gln Arg Gly Leu Gly Glu Ala Gln Leu Gly Asn Ser Ser Gly Asn
 260 265 270
 Phe Leu Leu Pro Asp Ala Gln Ser Ile Gln Ala Ala Ala Gly Phe
 275 280 285
 Ala Ser Lys Thr Pro Ala Asn Gln Ala Ile Ser Met Ile Asp Gly Pro
 290 295 300
 Ala Pro Asp Gly Tyr Pro Ile Ile Asn Tyr Glu Tyr Ala Ile Val Asn
 305 310 315 320
 Asn Arg Gln Lys Asp Ala Ala Thr Ala Gln Thr Leu Gln Ala Phe Leu
 325 330 335
 His Trp Ala Ile Thr Asp Gly Asn Lys Ala Ser Phe Leu Asp Gln Val
 340 345 350
 His Phe Gln Pro Leu Pro Pro Ala Val Val Lys Leu Ser Asp Ala Leu
 355 360 365
 Ile Ala Thr Ile Ser Ser
 370

(2) INFORMATION FOR SEQ ID NO:154:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1993 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:154:

TGTTCTTCGA CGGCAGGCTG GTGGAGGAAG GGCCACCGA ACAGCTGTTC TCCTCGCCGA

60

AGCATGCGGA AACCGCCCGA TACGTCGCCG GACTGTCTGGG GGACGTCAAG GACGCCAAGC	120
GCGGAAATTG AAGAGCACAG AAAGGTATGG CGTGAAAATT CGTTTGCATA CGCTGTTGGC	180
CGTGTTGACC GCTGCGCCGC TGCTGCTAGC AGCGGCGGGC TGTGGCTCGA AACCACCGAG	240
CGGTTGCGCT GAAACGGGCG CCGGCGCCGG TACTGTCTCG ACTACCCCCG CGTCGTCTGCC	300
GGTGACGTTG GCGGAGACCG GTAGCACGCT GCTCTACCCG CTGTTCAACC TGTGGGGTCC	360
GGCCTTTTAC GAGAGGTATC CGAACGTCAC GATCACCGCT CAGGGCACCG GTTCTGGTGC	420
CGGGATCTCG CAGGCCGCCG CCGGGACGGT CAACATTGGG GCCTCCGACG CCTATCTGTC	480
GGAAGGTGAT ATGGCCGCGC ACAAGGGGCT GATGAACATC GCGCTAGCCA TCTCCGCTCA	540
GCAGGTCAAC TACAACCTGC CCGGAGTGAG CGAGCACCTC AAGCTGAACG GAAAAGTCCT	600
GGCGGCCATG TACCAGGGCA CCATCAAAAC CTGGGACGAC CCGCAGATCG CTGCGCTCAA	660
CCCCGGCGTG AACCTGCCCC GCACCGCGGT AGTTCCGCTG CACCGCTCCG ACGGGTCCGG	720
TGACACCTTC TTGTTACCC AGTACCTGTC CAAGCAAGAT CCCGAGGGCT GGGGCAAGTC	780
GCCCGGCTTC GGCACCACCG TCGACTTCCC GGCGGTGCCG GGTGCGCTGG GTGAGAACGG	840
CAACGGCGGC ATGGTGACCG GTTGCGCCGA GACACCGGGC TCGGTGGCCT ATATCGGCAT	900
CAGCTTCCTC GACCAGGCCA GTCAACGGGG ACTCGGCGAG GCCCAACTAG GCAATAGCTC	960
TGGCAATTTT TTGTTGCCCG ACGCGCAAAG CATTAGGCC GCGGCGGCTG GCTTCGCATC	1020
GAAAACCCCG GCGAACCAGG CGATTTTCAT GATCGACGGG CCCGCCCCGG ACGGCTACCC	1080
GATCATCAAC TACGAGTACG CCATCGTCAA CAACCGGCAA AAGGACGCCG CCACCGCGCA	1140
GACCTTGACG GCATTTCTGC ACTGGGCGAT CACCGACGGC AACAAGGCCT CGTTCCTCGA	1200
CCAGGTTTAT TTCCAGCCGC TGCCGCCCGC GGTGGTGAAG TTGTCTGACG CGTTGATCGC	1260
GACGATTTCC AGCTAGCCTC GTTGACCACC ACGCGACAGC AACCTCCGTC GGGCCATCGG	1320
GCTGCTTTGC GGAGCATGCT GGCCCGTGCC GGTGAAGTCG GCCGCGCTGG CCCGGCCATC	1380
CGGTGGTTGG GTGGGATAGG TGCGGTGATC CCGCTGCTTG CGCTGGTCTT GGTGCTGGTG	1440
GTGCTGGTCA TCGAGGCGAT GGGTGCGATC AGGCTCAACG GGTGTCATTT CTTACCGCC	1500
ACCGAATGGA ATCCAGGCAA CACCTACGGC GAAACCGTTG TCACCGACGC GTCGCCCATC	1560
CGGTGCGGCG CTAATACGGG GCGTTGCCGC TGATCGTCGG GACGCTGGCG ACCTCGGCAA	1620
TCGCCCTGAT CATCGCGGTG CCGGTCTCTG TAGGAGCGGC GCTGGTGATC GTGGAACGGC	1680
TGCCGAAACG GTTGGCCGAG GCTGTGGGAA TAGTCCTGGA ATTGCTCGCC GGAATCCCCA	1740
GCGTGGTCGT CGGTTTGTGG GGGGCAATGA CGTTCGGGCC GTTCATCGCT CATCACATCG	1800
CTCCGGTGAT CGCTCACAAC GCTCCCGATG TGCCGGTGCT GAACTACTTG CGCGGCGACC	1860
CGGGCAACGG GGAGGGCATG TTGGTGTCG GTCTGGTGTT GGCGGTGATG GTCGTTCCCA	1920

TTATCGCCAC CACCACTCAT GACCTGTTCC GGCAGGTGCC GGTGTTGCCC CGGGAGGGCG 1980
 CGATCGGGAA TTC 1993

(2) INFORMATION FOR SEQ ID NO:155:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 374 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:155:

Met	Lys	Ile	Arg	Leu	His	Thr	Leu	Leu	Ala	Val	Leu	Thr	Ala	Ala	Pro	1	5	10	15
Leu	Leu	Leu	Ala	Ala	Ala	Gly	Cys	Gly	Ser	Lys	Pro	Pro	Ser	Gly	Ser	20	25	30	
Pro	Glu	Thr	Gly	Ala	Gly	Ala	Gly	Thr	Val	Ala	Thr	Thr	Pro	Ala	Ser	35	40	45	
Ser	Pro	Val	Thr	Leu	Ala	Glu	Thr	Gly	Ser	Thr	Leu	Leu	Tyr	Pro	Leu	50	55	60	
Phe	Asn	Leu	Trp	Gly	Pro	Ala	Phe	His	Glu	Arg	Tyr	Pro	Asn	Val	Thr	65	70	75	80
Ile	Thr	Ala	Gln	Gly	Thr	Gly	Ser	Gly	Ala	Gly	Ile	Ala	Gln	Ala	Ala	85	90	95	
Ala	Gly	Thr	Val	Asn	Ile	Gly	Ala	Ser	Asp	Ala	Tyr	Leu	Ser	Glu	Gly	100	105	110	
Asp	Met	Ala	Ala	His	Lys	Gly	Leu	Met	Asn	Ile	Ala	Leu	Ala	Ile	Ser	115	120	125	
Ala	Gln	Gln	Val	Asn	Tyr	Asn	Leu	Pro	Gly	Val	Ser	Glu	His	Leu	Lys	130	135	140	
Leu	Asn	Gly	Lys	Val	Leu	Ala	Ala	Met	Tyr	Gln	Gly	Thr	Ile	Lys	Thr	145	150	155	160
Trp	Asp	Asp	Pro	Gln	Ile	Ala	Ala	Leu	Asn	Pro	Gly	Val	Asn	Leu	Pro	165	170	175	
Gly	Thr	Ala	Val	Val	Pro	Leu	His	Arg	Ser	Asp	Gly	Ser	Gly	Asp	Thr	180	185	190	
Phe	Leu	Phe	Thr	Gln	Tyr	Leu	Ser	Lys	Gln	Asp	Pro	Glu	Gly	Trp	Gly	195	200	205	
Lys	Ser	Pro	Gly	Phe	Gly	Thr	Thr	Val	Asp	Phe	Pro	Ala	Val	Pro	Gly	210	215	220	
Ala	Leu	Gly	Glu	Asn	Gly	Asn	Gly	Gly	Met	Val	Thr	Gly	Cys	Ala	Glu	225	230	235	240

160

Thr Pro Gly Cys Val Ala Tyr Ile Gly Ile Ser Phe Leu Asp Gln Ala
 245 250 255
 Ser Gln Arg Gly Leu Gly Glu Ala Gln Leu Gly Asn Ser Ser Gly Asn
 260 265 270
 Phe Leu Leu Pro Asp Ala Gln Ser Ile Gln Ala Ala Ala Gly Phe
 275 280 285
 Ala Ser Lys Thr Pro Ala Asn Gln Ala Ile Ser Met Ile Asp Gly Pro
 290 295 300
 Ala Pro Asp Gly Tyr Pro Ile Ile Asn Tyr Glu Tyr Ala Ile Val Asn
 305 310 315 320
 Asn Arg Gln Lys Asp Ala Ala Thr Ala Gln Thr Leu Gln Ala Phe Leu
 325 330 335
 His Trp Ala Ile Thr Asp Gly Asn Lys Ala Ser Phe Leu Asp Gln Val
 340 345 350
 His Phe Gln Pro Leu Pro Pro Ala Val Val Lys Leu Ser Asp Ala Leu
 355 360 365
 Ile Ala Thr Ile Ser Ser
 370

(2) INFORMATION FOR SEQ ID NO:156:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1777 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:156:

GGTCTTGACC ACCACCTGGG TGTCGAAGTC GGTGCCCCGA TTGAAGTCCA GGTACTCGTG	60
GGTGCGGCGG GCGAAACAAT AGCGACAAGC ATGCGAGCAG CCGCGGTAGC CGTTGACGGT	120
GTAGCGAAAC GGCAACGCGG CCGCGTTGGG CACCTTGTTT AGCGCTGATT TGCACAACAC	180
CTCGTGGAAG GTGATGCCGT CGAATTGTGG CGCGCGAAG CTGCGGACCA GGCCGATCCG	240
CTGCAACCCG GCAGCGCCCG TCGTCAACGG GCATCCCGTT CACCGCGACG GCTTGCCGGG	300
CCCAACGCAT ACCATTATTC GAACAACCGT TCTATACTTT GTCAACGCTG GCCGCTACCG	360
AGCGCCGCAC AGGATGTGAT ATGCCATCTC TGCCCGCACA GACAGGAGCC AGGCCTTATG	420
ACAGCATTCG GCGTCGAGCC CTACGGGCAG CCGAAGTACC TAGAAATCGC CGGGAAGCGC	480
ATGGCGTATA TCGACGAAGG CAAGGGTGAC GCCATCGTCT TTCAGCACGG CAACCCACG	540
TCGTCTTACT TGTGGCGCAA CATCATGCCG CACTTGGAAG GGCTGGGCGG GCTGGTGGCC	600

TGCGATCTGA	TCGGGATGGG	CGCGTCGGAC	AAGCTCAGCC	CATCGGGACC	CGACCGCTAT	660
AGCTATGGCG	AGCAACGAGA	CTTTTTGTTC	GCGCTCTGGG	ATGCGCTCGA	CCTCGGCGAC	720
CACGTGGTAC	TGGTGCTGCA	CGACTGGGGC	TCGGCGCTCG	GCTTCGACTG	GGCTAACCAG	780
CATCGCGACC	GAGTGCAGGG	GATCGCGTTC	ATGGAAGCGA	TCGTCACCCC	GATGACGTGG	840
GCGGACTGGC	CGCCGGCCGT	GCGGGGTGTG	TTCCAGGGTT	TCCGATCGCC	TCAAGGCGAG	900
CCAATGGCGT	TGGAGCACAA	CATCTTTGTC	GAACGGGTGC	TGCCCGGGGC	GATCCTGCGA	960
CAGCTCAGCG	ACGAGGAAAT	GAACCACTAT	CGGCGGCCAT	TCGTGAACGG	CGGCGAGGAC	1020
CGTCGCCCCA	CGTTGTCTGT	GCCACGAAAC	CTTCCAATCG	ACGGTGAGCC	CGCCGAGGTC	1080
GTCGCGTTGG	TCAACGAGTA	CCGGAGCTGG	CTCGAGGAAA	CCGACATGCC	GAAACTGTTC	1140
ATCAACGCCG	AGCCCGGGCG	GATCATCACC	GGCCGCATCC	GTGACTATGT	CAGGAGCTGG	1200
CCCAACCAGA	CCGAAATCAC	AGTGCCCGGC	GTGCATTTTC	TTCAGGAGGA	CAGCGATGGC	1260
GTCGTATCGT	GGGCGGGCGC	TCGGCAGCAT	CGGCGACCTG	GGAGCGCTCT	CATTTACGA	1320
GACCAAGAAT	GTGATTTCCG	GCGAAGGCGG	CGCCCTGCTT	GTCAACTCAT	AAGACTTCCT	1380
GCTCCGGGCA	GAGATTCTCA	GGGAAAAGGG	CACCAATCGC	AGCCGCTTCC	TTCGCAACGA	1440
GGTCGACAAA	TATACGTGGC	AGGACAAAGG	TCTTCCTATT	TGCCCAGCGA	ATTAGTCGCT	1500
GCCTTTCTAT	GGGCTCAGTT	CGAGGAAGCC	GAGCGGATCA	CGCGTATCCG	ATTGGACCTA	1560
TGGAACCGGT	ATCATGAAAG	CTTCGAATCA	TTGGAACAGC	GGGGGCTCCT	GCGCCGTCCG	1620
ATCATCCCAC	AGGGCTGCTC	TCACAACGCC	CACATGTACT	ACGTGTACT	AGCGCCCAGC	1680
GCCGATCGGG	AGGAGGTGCT	GGCGCGTCTG	ACGAGCGAAG	GTATAGGCGC	GGTCTTTCAT	1740
TACGTGCCGC	TTCACGATTC	GCCGGCCGGG	CGTCGCT			1777

(2) INFORMATION FOR SEQ ID NO:157:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 324 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:157:

GAGATTGAAT	CGTACCGGTC	TCCTTAGCGG	CTCCGTCCCG	TGAATGCCCA	TATCACGCAC	60
GGCCATGTTC	TGGCTGTCGA	CCTTCGCCCC	ATGCCCGGAC	GTTGGTAAAC	CCAGGGTTTG	120
ATCAGTAATT	CCGGGGGACG	GTTGCGGGAA	GGCGGCCAGG	ATGTGCGTGA	GCCGCGGCGC	180
CGCCGTCGCC	CAGGCGACCG	CTGGATGCTC	AGCCCCGGTG	CGGCGACGTA	GCCAGCGTTT	240

GGCGCGTGTC GTCCACAGTG GTACTCCGGT GACGACGCGG CGCGGTGCCT GGGTGAAGAC 300
 CGTGACCGAC GCCGCCGATT CAGA 324

(2) INFORMATION FOR SEQ ID NO:158:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1338 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:158:

GCGGTACCGC CGCGTTGCGC TGGCACGGGA CCTGTACGAC CTGAACCACT TCGCCTCGCG 60
 AACGATTGAC GAACCGCTCG TCGGGCGGCT GTGGGTGCTC AAGGTGTGGG GTGATGTCGT 120
 CGATGACCGG CGCGGCACCC GGCCACTACG CGTCGAAGAC GTCCTCGCCG CCCGCAGCGA 180
 GCACGACTTC CAGCCCGACT CGATCGGCGT GCTGACCCGT CCTGTGCTGTA TGGCTGCCTG 240
 GGAAGCTCGC GTTCGGAAGC GATTTGCGTT CCTCACTGAC CTCGACGCCG ACGAGCAGCG 300
 GTGGGCCGCC TGCACGAAC GGCACCGCCG CGAAGTGGAG AACGCGCTGG CGGTGCTGCG 360
 GTCCTGATCA ACCTGCCGGC GATCGTGCCG TTCCGCTGGC ACGGTTGCGG CTGGACGCGG 420
 CTGAATCGAC TAGATGAGAG CAGTTGGGCA CGAATCCGGC TGTGGTGGTG AGCAAGACAC 480
 GAGTACTGTC ATCACTATTG GATGCACTGG ATGACCGGCC TGATTGAGCA GGACCAATGG 540
 AACTGCCCCG GGCAAAACGT CTCGGAGATG ATCGGCGTCC CCTCGGAACC CTGCGGTGCT 600
 GCGGTCATTC GGACATCGGT CCGGCTCGCG GGATCGTGGT GACGCCAGCG CTGAAGGAGT 660
 GGAGCGCGGC GGTGCACGCG CTGCTGGACG GCCGGCAGAC GGTGCTGCTG CGTAAGGGCG 720
 GGATCGGCGA GAAGCGCTTC GAGGTGGCGG CCCACGAGTT CTTGTTGTTC CCGACGGTCG 780
 CGCACAGCCA CGCCGAGCGG GTTCGCCCCG AGCACCGCGA CCTGCTGGGC CCGGCGGCCG 840
 CCGACAGCAC CGACGAGTGT GTGCTACTGC GGGCCGCAGC GAAAGTTGTT GCCGCACTGC 900
 CGGTTAACCG GCCAGAGGGT CTGGACGCCA TCGAGGATCT GCACATCTGG ACCGCCGAGT 960
 CGGTGCGCGC CGACCGGCTC GACTTTCGGC CCAAGCACAA ACTGGCCGTC TTGGTGGTCT 1020
 CGGCGATCCC GCTGGCCGAG CCGGTCCGGC TGGCGCGTAG GCCCGAGTAC GGCGGTTGCA 1080
 CCAGCTGGGT GCAGCTGCCG GTGACGCCGA CGTTGGCGGC GCCGGTGAC GACGAGGCCG 1140
 CGCTGGCCGA GGTGCGGCCG CGGGTCCGCG AGGCCGTGGG TTGACTGGGC GGCATCGCTT 1200
 GGGTCTGAGC TGTACGCCCA GTCGGCGCTG CGAGTGATCT GCTGTCGGTT CGGTCCCTGC 1260
 TGGCGTCAAT TGACGGCGCG GGCAACAGCA GCATTGGCGG CGCCATCCTC CGCGCGGCCG 1320

GCGCCACCG CTACAACC

1338

(2) INFORMATION FOR SEQ ID NO:159:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 321 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:159:

CCGGCGGCAC CGGCGGCACC GGCGGTACCG GCGGCAACGG CGCTGACGCC GCTGCTGTGG	60
TGGGCTTCGG CGCGAACGGC GACCCTGGCT TCGCTGGCGG CAAAGGCGGT AACGGCGGAA	120
TAGGTGGGGC CGCGGTGACA GGCGGGGTCG CCGGCGACGG CGGCACCGGC GGCAAAGGTG	180
GCACCGGCGG TGCCGGCGGC GCCGGCAACG ACGCCGGCAG CACCGGCAAT CCCGGCGGTA	240
AGGGCGGCGA CGGCGGGATC GGCGGTGCCG GCGGGGCCGG CGGCGCGGCC GGCACCGGCA	300
ACGGCGGCCA TGCCGGCAAC C	321

(2) INFORMATION FOR SEQ ID NO:160:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 492 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:160:

GAAGACCCGG CCCC GCCATA TCGATCGGCT CGCCGACTAC TTTCGCCGAA CGTGACACGG	60
GCGGCGTCGG GCTGATCATC ACCGGTGGCT ACGCGCCCAA CCGCACCGGA TGGCTGCTGC	120
CGTTCGCCTC CGAACTCGTC ACTTCGGCGC AAGCCCGACG GCACCGCCGA ATCACCAGGG	180
CGGTCCACGA TTCGGGTGCA AAGATCCTGC TGCAAATCCT GCACGCCGGA CGCTACGCCT	240
ACCACCCACT TGCGGTCAGC GCCTCGCCGA TCAAGGCGCC GATCACCCCG TTTCGTCCGC	300
GAGCACTATC GGCTCGCGGG GTCGAAGCGA CCATCGCGGA TTTCGCCCCG TGCGCGCAGT	360
TGGCCCGCGA TGCCGGCTAC GACGGCGTCG AAATCATGGG CAGCGAAGGG TATCTGCTCA	420
ATCAGTTCCT GGCGCCGCGC ACCAACAAGC GCACCGACTC GTGGGGCGGC ACACCGGCCA	480
ACCGTCGCCG GT	492

(2) INFORMATION FOR SEQ ID NO:161:

164

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 536 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:161:

```

Phe Ala Gln His Leu Val Glu Gly Asp Ala Val Glu Leu Trp Arg Ala
1           5           10           15
Asn Ala Ala Asp Gln Ala Asp Pro Leu Gln Pro Gly Ser Ala Arg Arg
20           25           30
Gln Arg Ala Ser Arg Ser Pro Arg Arg Leu Ala Gly Pro Asn Ala Tyr
35           40           45
His Tyr Ser Asn Asn Arg Ser Ile Leu Cys Gln Arg Trp Pro Leu Pro
50           55           60
Ser Ala Ala Gln Asp Val Ile Cys His Leu Cys Pro His Arg Gln Glu
65           70           75           80
Pro Gly Leu Met Thr Ala Phe Gly Val Glu Pro Tyr Gly Gln Pro Lys
85           90           95
Tyr Leu Glu Ile Ala Gly Lys Arg Met Ala Tyr Ile Asp Glu Gly Lys
100          105          110
Gly Asp Ala Ile Val Phe Gln His Gly Asn Pro Thr Ser Ser Tyr Leu
115          120          125
Trp Arg Asn Ile Met Pro His Leu Glu Gly Leu Gly Arg Leu Val Ala
130          135          140
Cys Asp Leu Ile Gly Met Gly Ala Ser Asp Lys Leu Ser Pro Ser Gly
145          150          155          160
Pro Asp Arg Tyr Ser Tyr Gly Glu Gln Arg Asp Phe Leu Phe Ala Leu
165          170          175
Trp Asp Ala Leu Asp Leu Gly Asp His Val Val Leu Val Leu His Asp
180          185          190
Trp Gly Ser Ala Leu Gly Phe Asp Trp Ala Asn Gln His Arg Asp Arg
195          200          205
Val Gln Gly Ile Ala Phe Met Glu Ala Ile Val Thr Pro Met Thr Trp
210          215          220
Ala Asp Trp Pro Pro Ala Val Arg Gly Val Phe Gln Gly Phe Arg Ser
225          230          235          240
Pro Gln Gly Glu Pro Met Ala Leu Glu His Asn Ile Phe Val Glu Arg
245          250          255
Val Leu Pro Gly Ala Ile Leu Arg Gln Leu Ser Asp Glu Glu Met Asn
260          265          270

```

165

His Tyr Arg Arg Pro Phe Val Asn Gly Gly Glu Asp Arg Arg Pro Thr
 275 280 285
 Leu Ser Trp Pro Arg Asn Leu Pro Ile Asp Gly Glu Pro Ala Glu Val
 290 295 300
 Val Ala Leu Val Asn Glu Tyr Arg Ser Trp Leu Glu Glu Thr Asp Met
 305 310 315 320
 Pro Lys Leu Phe Ile Asn Ala Glu Pro Gly Ala Ile Ile Thr Gly Arg
 325 330 335
 Ile Arg Asp Tyr Val Arg Ser Trp Pro Asn Gln Thr Glu Ile Thr Val
 340 345 350
 Pro Gly Val His Phe Val Gln Glu Asp Ser Asp Gly Val Val Ser Trp
 355 360 365
 Ala Gly Ala Arg Gln His Arg Arg Pro Gly Ser Ala Leu Ile Ser Arg
 370 375 380
 Asp Gln Glu Cys Asp Phe Arg Arg Arg Arg Arg Pro Ala Cys Gln Leu
 385 390 395 400
 Ile Arg Leu Pro Ala Pro Gly Arg Asp Ser Gln Gly Lys Gly His Gln
 405 410 415
 Ser Gln Pro Leu Pro Ser Gln Arg Gly Arg Gln Ile Tyr Val Ala Gly
 420 425 430
 Gln Arg Ser Ser Tyr Leu Pro Ser Glu Leu Val Ala Ala Phe Leu Trp
 435 440 445
 Ala Gln Phe Glu Glu Ala Glu Arg Ile Thr Arg Ile Arg Leu Asp Leu
 450 455 460
 Trp Asn Arg Tyr His Glu Ser Phe Glu Ser Leu Glu Gln Arg Gly Leu
 465 470 475 480
 Leu Arg Arg Pro Ile Ile Pro Gln Gly Cys Ser His Asn Ala His Met
 485 490 495
 Tyr Tyr Val Leu Leu Ala Pro Ser Ala Asp Arg Glu Glu Val Leu Ala
 500 505 510
 Arg Leu Thr Ser Glu Gly Ile Gly Ala Val Phe His Tyr Val Pro Leu
 515 520 525
 His Asp Ser Pro Ala Gly Arg Arg
 530 535

(2) INFORMATION FOR SEQ ID NO:162:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 284 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

166

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:162:

```

Asn Glu Ser Ala Pro Arg Ser Pro Met Leu Pro Ser Ala Arg Pro Arg
1          5          10          15
Tyr Asp Ala Ile Ala Val Leu Leu Asn Glu Met His Ala Gly His Cys
20          25          30
Asp Phe Gly Leu Val Gly Pro Ala Pro Asp Ile Val Thr Asp Ala Ala
35          40          45
Gly Asp Asp Arg Ala Gly Leu Gly Val Asp Glu Gln Phe Arg His Val
50          55          60
Gly Phe Leu Glu Pro Ala Pro Val Leu Val Asp Gln Arg Asp Asp Leu
65          70          75          80
Gly Gly Leu Thr Val Asp Trp Lys Val Ser Trp Pro Arg Gln Arg Gly
85          90          95
Ala Thr Val Leu Ala Ala Val His Glu Trp Pro Pro Ile Val Val His
100         105         110
Phe Leu Val Ala Glu Leu Ser Gln Asp Arg Pro Gly Gln His Pro Phe
115         120         125
Asp Lys Asp Val Val Leu Gln Arg His Trp Leu Ala Leu Arg Arg Ser
130         135         140
Glu Thr Leu Glu His Thr Pro His Gly Arg Arg Pro Val Arg Pro Arg
145         150         155         160
His Arg Gly Asp Asp Arg Phe His Glu Arg Asp Pro Leu His Ser Val
165         170         175
Ala Met Leu Val Ser Pro Val Glu Ala Glu Arg Arg Ala Pro Val Val
180         185         190
Gln His Gln Tyr His Val Val Ala Glu Val Glu Arg Ile Pro Glu Arg
195         200         205
Glu Gln Lys Val Ser Leu Leu Ala Ile Ala Ile Ala Val Gly Ser Arg
210         215         220
Trp Ala Glu Leu Val Arg Arg Ala His Pro Asp Gln Ile Ala Gly His
225         230         235         240
Gln Pro Ala Gln Pro Phe Gln Val Arg His Asp Val Ala Pro Gln Val
245         250         255
Arg Arg Arg Gly Val Ala Val Leu Lys Asp Asp Gly Val Thr Leu Ala
260         265         270
Phe Val Asp Ile Arg His Ala Leu Pro Gly Asp Phe
275         280

```

(2) INFORMATION FOR SEQ ID NO:163:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 264 base pairs

167

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:163:

ATGAACATGT CGTCGGTGGT GGGTCGCAAG GCCTTTGCGC GATTCGCCCG CTACTCCTCC	60
GCCATGCACG CGATCGCCGG TTTCTCCGAT GCGTTGCGCC AAGAGCTGCG GGGTAGCGGA	120
ATCGCCGTCT CGGTGATCCA CCCGGCGCTG ACCCAGACAC CGCTGTTGGC CAACGTCGAC	180
CCCGCCGACA TGCCGCCGCC GTTTCGCAGC CTCACGCCCA TTCCCGTTCA CTGGGTCGCG	240
GCAGCGGTGC TTGACGGTGT GGCG	264

(2) INFORMATION FOR SEQ ID NO:164:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1171 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:164:

TAGTCGGCGA CGATGACGTC GCGGTCCAGG CCGACCGCTT CAAGCACCAG CGCGACCACG	60
AAGCCGGTGC GATCCTTACC CGCGAAGCAG TGGGTGAGCA CCGGGCGTCC GCGGGCAAGC	120
AGTGTGACGA CACGATGTAG CGCGCGCTGT GCTCCATTGC GCGTTGGGAA TTGGCGATAC	180
TCGTGCGTCA TGTAGCGGGT GGCCGCGTCA TTTATCGACT GGCTGGATTG GCCGGACTCG	240
CCGTTGGACC CGTCATTGGT TAGCAGCCTC TTGAATGCGG TTTCGTGCGG CGCTGAGTCG	300
TCGGCGTCAT CATCGGCGAG GTCGGGGAAC GGCAGCAGGT GGACGTCGAT GCCGTCCGGA	360
ACCCGTCTTG GACCGCGGCG GGCAACCTCC CGGGACGACC GCAGGTCGGC AACGTCGGTG	420
ATCCCCAGCC GGCGCAGCGT TGCCCCCTCG GCCGAATTTC GCACGAGGCT GGCGAGCCAC	480
CGGGCATCAC CAAGCAACGC TTGCCAGTA CGGATCGTCA CTTCCGCATC CGGCAGACCA	540
ATCTCCTCGC CGCCCATCGT CAGATCCCGC TCGTGCGTTG ACAAGAACGG CCGCAGATGT	600
GCCAGCGGGT ATCGGAGATT GAACCGCGCA CGCAGTTCTT CAATCGCTGC GCGTGCCGC	660
ACTATTGGCA CTTTCCGGCG GTCGCGGTAT TCAGCAAGCA TGCGAGTCTC GACGAAGTCG	720
CCCCACGTAA CCCACGGCGT AGCTCCCGGC GTGACGCGGA GGATCGGCGG GTGATCTTTG	780
CCGCCACGCT CGTAGCCGTT GATCCACCGC TTCGCGGTGC CGGCGGGGAG GCCGATCAGC	840

168

```

TTATCGACCT CGGCGTATGC CGACGGCAAG CTGGGCGCGT TCGTCGAGGT CAAGAACTCC      900
ACCATCGGCA CCGGCACCAA GGTGCCGCAC CTGACCTACG TCGGCGACGC CGACATCGGC      960
GAGTACAGCA ACATCGGCGC CTCCAGCGTG TTCGTCAACT ACGACGGTAC GTCCAAACGG      1020
CGCACCACCG TCGGTTTCGCA CGTACGGACC GGGTCCGACA CCATGTTCGT GGCCCCAGTA      1080
ACCATCGGCG ACGGCGCGTA TACCGGGGCC GGCACAGTGG TGCGGGAGGA TGTCCCGCCG      1140
GGGGCGCTGG CAGTGTCTGGC GGGTCCGCAA C                                1171

```

(2) INFORMATION FOR SEQ ID NO:165:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 227 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:165:

```

GCAAAGGCGG CACCGGCGGG GCCGGCATGA ACAGCCTCGA CCCGCTGCTA GCCGCCCAAG      60
ACGGCGGCCA AGGCGGCACC GGCGGCACCG GCGGCAACGC CGGCGCCGGC GGCACCAGCT      120
TCACCCAAGG CGCCGACGGC AACGCCGGCA ACGGCGGTGA CGGCGGGGTC GGCGGCAACG      180
CGGGAAACGG CGGAAACGGC GCAGACAACA CCACCACCGC CGCCGCC                                227

```

(2) INFORMATION FOR SEQ ID NO:166:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 304 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:166:

```

CCTCGCCACC ATGGGCGGGG AGGGCGGTAG CGGTGGCGCC GGCTCTACCC CAGGCGCCAA      60
GGGCGCCAC GGCTTCACTC CAACCAGCGG CGGCGACGGC GGCGACGGCG GCAACGGCGG      120
CAACTCCCAA GTGGTCGGCG GCAACGGCGG CGACGGCGGC AATGGCGGCA ACGGCGGCAG      180
CGCCGGCACG GGCGGCAACG GCGGCCCGGG CGGCGACGGC GCGTTTGGTG GCATGAGTGC      240
CAACGCCACC AACCCTGGTG AAAACGGGCC AAACGGTAAC CCCGGCGGCA ACGGTGGCGC      300
CGGC                                304

```

(2) INFORMATION FOR SEQ ID NO:167:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1439 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:167:

GTGGGACGCT GCCGAGGCTG TATAACAAGG ACAACATCGA CCAGCGCCGG CTCGGTGAGC	60
TGATCGACCT ATTTAACAGT GCGCGCTTCA GCCGGCAGGG CGAGCACCGC GCCCGGGATC	120
TGATGGGTGA GGTCTACGAA TACTTCCTCG GCAATTTTCG TCGCGCGGAA GGGAAGCGGG	180
GTGGCGAGTT CTTTACCCCG CCCAGCGTGG TCAAGGTGAT CGTGGAGGTG CTGGAGCCGT	240
CGAGTGGGCG GGTGTATGAC CCGTGCTGCG GTTCCGGAGG CATGTTTGTG CAGACCGAGA	300
AGTTCATCTA CGAACACGAC GGCGATCCGA AGGATGTCTC GATCTATGGC CAGGAAAGCA	360
TTGAGGAGAC CTGGCGGATG GCGAAGATGA ACCTCGCCAT CCACGGCATC GACAACAAGG	420
GGCTCGGCGC CCGATGGAGT GATACCTTCG CCCGCGACCA GCACCCGGAC GTGCAGATGG	480
ACTACGTGAT GGCCAATCCG CCGTTCAACA TCAAAGACTG GGCCCGCAAC GAGGAAGACC	540
CACGCTGGCG CTTCGGTGTT CCGCCCGCCA ATAACGCCAA CTACGCATGG ATTCAGCACA	600
TCCTGTACAA CTTGGCGCCG GGAGGTCGGG CGGGCGTGGT GATGGCCAAC GGGTCGATGT	660
CGTCGAACTC CAACGGCAAG GGGGATATTC GCGCGCAAAT CGTGGAGGCG GATTTGGTTT	720
CCTGCATGGT CGCGTTACCC ACCCAGCTGT TCCGCAGCAC CGGAATCCCG GTGTGCCTGT	780
GGTTTTTCGC CAAAACAAG GCGGCAGGTA AGCAAGGGTC TATCAACCGG TCGGGGCAGG	840
TGCTGTTCAT CGACGCTCGT GAACTGGGCG ACCTAGTGGA CCGGGCCGAG CGGGCGCTGA	900
CCAACGAGGA GATCGTCCGC ATCGGGGATA CCTTCCACGC GAGCACGACC ACCGGCAACG	960
CCGGCTCCGG TGGTGCCGGC GGTAATGGGG GCACTGGCCT CAACGGCGCG GCGGTGCTG	1020
GCGGGGCCGG CGGCAACGCG GGTGTCGCCG GCGTGTCTT CGGCAACGCT GTGGGCGGCG	1080
ACGGCGGCAA CGGCGGCAAC GGCGGCCACG GCGGCGACGG CACGACGGGC GGCGCCGGCG	1140
GCAAGGGCGG CAACGGCAGC AGCGGTGCCG CCAGCGGCTC AGGCGTCGTC AACGTCACCG	1200
CCGGCCACGG CGGCAACGGC GGCAATGGCG GCAACGGCGG CAACGGCTCC GCGGGCGCCG	1260
GCGGCCAGGG CGGTGCCGGC GGCAGCGCCG GCAACGGCGG CCACGGCGGC GGTGCCACCG	1320
GCGGCGCCAG CGGCAAGGGC GGCAACGGCA CCAGCGGTGC CGCCAGCGGC TCAGGCGTCA	1380
TCAACGTCAC CGCCGGCCAC GGCGGCAACG GCGGCAATGG CCGCAACGGC GGCAACGGC	1439

(2) INFORMATION FOR SEQ ID NO:168:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 329 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:168:

GGGCCGGCGG GGCCGGATTT TCTCGTGCCT TGATTGTCGC TGGGGATAAC GGCGGTGATG	60
GTGGTAACGG CGGGATGGGC GGGGCTGGCG GGGCTGGCGG CCCCGGCGGG GCCGGCGGCC	120
TGATCAGCCT GCTGGGCGGC CAAGGCGCCG GCGGGGCGCG CGGGACCGGC GGGGCCGGCG	180
GTGTTGGCGG TGACGGCGGG GCCGGCGGCC CCGGCAACCA GGCCTTCAAC GCAGGTGCCG	240
GCGGGGCGCG CGGCCTGATC AGCCTGCTGG GCGGCCAAGG CGCCGGCGGG GCCGGCGGGA	300
CCGGCGGGGC CGGCGGTGTT GCGGCTGAC	329

(2) INFORMATION FOR SEQ ID NO:169:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 80 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:169:

GCAACGGTGG CAACGGCGGC ACCAGCACGA CCGTGGGGAT GGCCGGAGGT AACTGTGGTG	60
CCGCCGGGCT GATCGGCAAC	80

(2) INFORMATION FOR SEQ ID NO:170:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 392 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:170:

GGGCTGTGTC GCACTCACAC CGCCGCATTC GGCGACGTTG GCCGCCCAAT ATCCAGCTCA	60
AGGCCTACTA CTTACCGTCG GAGGACCGCC GCATCAAGGT GCGGGTCAGC GCCCAAGGAA	120
TCAAGGTCAT CGACCGCGAC GGGCATCGAG GCCGTCGTCG CGCGGCTCGG GCAGGATCCG	180

171

CCCCGGCGCA CTTCGCGCGC CAAGCGGGCT CATCGCTCCG AACGGCGGCG ATCCTGTGAG 240
 CACAAC TGAT GGCGCGCAAC GAGATTCGTC CAATTGTCAA GCCGTGTTCG ACCGCAGGGA 300
 CCGGTTATAC GTATGTCAAC CTATGTCACT CGCAAGAACC GGCATAACGA TCCCGTGATC 360
 CGCCGACAGC CCACGAGTGC AAGACCGTTA CA 392

(2) INFORMATION FOR SEQ ID NO:171:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 535 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:171:

ACCGGCGCCA CCGGCGGCAC CGGGTTCGCC GGTGGCGCCG GCGGGGCCGG CGGGCAGGGC 60
 GGTATCAGCG GTGCCGGCGG CACCAACGGC TCTGGTGGCG CTGGCGGCAC CGGCGGACAA 120
 GGCGGCGCCG GGGGCGCTGG CGGGGCCGGC GCCGATAACC CCACCGGCAT CGGCGGCGCC 180
 GGCGGCACCG GCGGCACCGG CGGAGCGGCC GGAGCCGGCG GGGCCGGTGG CGCCATCGGT 240
 ACCGGCGGCA CCGGCGGCGC GGTGGGCAGC GTCGGTAACG CCGGGATCGG CGGTACCGGC 300
 GGTACGGGTG GTGTCGGTGG TGCTGGTGGT GCAGGTGCGG CTGCGGCCGC TGGCAGCAGC 360
 GCTACCGGTG GCGCCGGGTT CGCCGGCGGC GCCGGCGGAG AAGGCGGACC GGGCGGCAAC 420
 AGCGGTGTGG GCGGCACCAA CGGCTCCGGC GGCGCCGGCG GTGCAGGCGG CAAGGGCGGC 480
 ACCGGAGGTG CCGGCGGGTC CGGCGCGGAC AACCCACCG GTGCTGGTTT CGCCG 535

(2) INFORMATION FOR SEQ ID NO:172:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 690 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:172:

CCGACGTCGC CGGGGCGATA CGGGGGTCAC CGACTACTAC ATCATCCGCA CCGAGAATCG 60
 GCCGCTGCTG CAACCGCTGC GGGCGGTGCC GGTCAATCGGA GATCCGCTGG CCGACCTGAT 120
 CCAGCCGAAC CTGAAGGTGA TCGTCAACCT GGGCTACGGC GACCCGAACCT ACGGCTACTC 180
 GACGAGCTAC GCCGATGTGC GAACGCCGTT CGGGCTGTGG CCGAACGTGC CGCCTCAGGT 240

172

CATCGCCGAT GCCCTGGCCG CCGGAACACA AGAAGGCATC CTTGACTTCA CGGCCGACCT	300
GCAGGCGCTG TCCGCGCAAC CGCTCACGCT CCCGCAGATC CAGCTGCCGC AACCCGCCGA	360
TCTGGTGGCC GCGGTGGCCG CCGCACCGAC GCCGGCCGAG GTGGTGAACA CGCTCGCCAG	420
GATCATCTCA ACCAACTACG CCGTCCTGCT GCCCACCCTG GACATCGCCC TCGCCTGGTC	480
ACCACCCTGC CGCTGTACAC CACCCAACTG TTCGTCAGGC AACTCGCTGC GGGCAATCTG	540
ATCAACGCGA TCGGCTATCC CCTGGCGGCC ACCGTAGGTT TAGGCACGAT CGATAGCGGG	600
CGGCGTGGA TTGCTACCC TCCTCGCGGC GGCCTCGGAC ACCGTTTCGAA ACATCGAGGG	660
CCTCGTCACC TAACGGATTC CCGACGGCAT	690

(2) INFORMATION FOR SEQ ID NO:173:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 407 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:173:

ACGGTGACGG CGGTACTGGC GCGGCCACG GCGGCAACGG CGGGAATCCC GGGTGGCTCT	60
TGGGCACAGC CGGGGGTGGC GGCAACGGTG GCGCCGGCAG CACCGGTACT GCAGGTGGCG	120
GCTCTGGGGG CACCGGCGGC GACGGCGGGA CCGGCGGGCG TGGCGGCCTG TTAATGGGCG	180
CCGGCGCCGG CGGGCACGGT GGCACTGGCG GCGCGGGCGG TGCCGGTGTC GACGGTGGCG	240
GCGCCGGCGG GGCCGGCGGG GCCGGCGGCA ACGGCGGCGC CGGGGGTCAA GCCGCCCTGC	300
TGTTGCGGGC CGGCGGCACC GGCGGAGCCG GCGGCTACGG CGGCGATGGC GGTGGCGGCG	360
GTGACGGCTT CGACGGCACG ATGGCCGGCC TGGGTGGTAC CGGTGGC	407

(2) INFORMATION FOR SEQ ID NO:174:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 468 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:174:

GATCGGTCAG CGCATCGCCC TCGGCGGCAA GCGATTCCGC GGTCTACCG AAGAACATCG	60
TGCACGCGGC GGCGCGGACC AGCCCGCTGC GCTGCGGCGC GTCGAACGCC TCCAGCAGGC	120

173

ACAGCCAGTC CTTGGCGGCC TCGAGGCGA ACACGTCGGT GTCACCGGTG TAGATCGCCG	180
GGATGCCCCG CTCCGCCAAC GCATTCCGGC ACGCCCGCGC GTCTTTGTGA TGCTCGACGA	240
TCACCGCGAT GTCTGCGGCC ACCACGGGCC GCCCGGCGAA GGTGGCCCCG CTGGCCAGTA	300
GCGCCGCGAC GTCGGCGGCC AGGTCGTCGG GGATGTGCCG GCGCAGCGCT CCGGCGCGAC	360
GCCCGAAAAA CGACCCTCA CCCAGCTGGG TCCCGCTGGC ATATCCCTTG CCGTCTGGG	420
CGATATTGGA CGCGCATGCC CCGACCGCGT ACAGGCCGGC CACCACCG	468

(2) INFORMATION FOR SEQ ID NO:175:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 219 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:175:

GGTGGTAACG GCGGCCAGGG TGGCATCGGC GGCGCCGGCG AGAGAGGCGC CGACGGCGCC	60
GGCCCCAATG CTAACGGCGC AAACGGCGAG AACGGCGGTA GCGGTGGTAA CCGTGGCGAC	120
GGCGGCGCCG GCGGCAATGG CGGCGCGGGC GGCAACGCGC AGGCGGCCGG GTACACCGAC	180
GGCGCCACGG GCACCGCGCG CGACGGCGGC AACGGCGGC	219

(2) INFORMATION FOR SEQ ID NO:176:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 494 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:176:

TAGCTCCGGC GAGGGCGGCA AGGGCGGCGA EGGTGCCAC GGCGGTGACG GCGTCGGCGG	60
CAACAGTTCC GTCACCCAAG GCGGCAGCGG CCGTGGCGGC GGCGCCGGCG GCGCCGGCGG	120
CAGCGGCTTT TTCGGCGGCA AGGGCGGCTT CGGCGGCGAC GGCGGTCAGG GCGGCCCCAA	180
CGGCGGCGGT ACCGTCGGCA CCGTGGCCGG TGGCGGCGGC AACGGCGGTG TCGGCGGCCG	240
GGGCGGCGAC GGCGTCTTTG CCGGTGCCGG CGGCCAGGGC GGCCTCGGTG GGCAGGGCGG	300
CAATGGCGGC GGCTCCACCG GCGGCAACGG CGGCCTTGGC GGCGCGGGCG GTGGCGGAGG	360
CAACGCCCCG GCTCGTGCCG AATCCGGGCT GACCATGGAC AGCGCGGCCA AGTTCGCTGC	420

CATCGCATCA GCGCGTACT GCGCCGAACA CCTGGAACAT CACCCGAGTT AGCGGGGCGC 480
ATTTCCTGAT CACC 494

(2) INFORMATION FOR SEQ ID NO:177:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 220 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:177:

GGGCCGGTGG TGCCGCGGGC CAGCTCTTCA GCGCCGGAGG CGCGGCGGGT GCCGTTGGGG 60
TTGGCGGCAC CGGCGGCCAG GGTGGGGCTG GCGGTGCCGG AGCGGCCGGC GCCACGCCCC 120
CCGCCAGCAC AGGTCTAACC GGTGGTACCG GGTTCGCTGG CGGGGCCGGC GGCGTCGGCG 180
GCCAGAGCGG CAACGCCATT GCCGGCGGCA TCAACGGCTC 220

(2) INFORMATION FOR SEQ ID NO:178:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 388 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:178:

ATGGCGGCAA CGGGGGCCCC GCGGGTGCTG GCGGGGCCGG CGACTACAAT TTCCAACGGC 60
GGGCAGGGTG GTGCCGGCGG CCAAGGCGGC CAAGGCGGCC TGGGCGGGGC AAGCACCACC 120
TGATCGGCCT AGCCGCACCC GGGAAAGCCG ATCCAACAGG CGACGATGCC GCCTTCCTTG 180
CCGCGTTGGA CCAGGCCGGC ATCACCTACG CTGACCCAGG CCACGCCATA ACGGCCGCCA 240
AGGCGATGTG TGGGCTGTGT GCTAACGGCG TAACAGGTCT ACAGCTGGTC GCGGACCTGC 300
GGGACTACAA TCCCGGGCTG ACCATGGACA GCGCGGCCAA GTTCGCTGCC ATCGCATCAG 360
GCGCGTACTG CCGCGAACAC CTGGAACA 388

(2) INFORMATION FOR SEQ ID NO:179:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 400 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:179:

GCAAAGGCGG CACCGGCGGG GCCGGCATGA ACAGCCTCGA CCCGCTGCTA GCCGCCCAAG	60
ACGGCGGCCA AGGCGGCACC GGCGGCACCG GCGGCAACGC CGGCGCCGGC GGCACCAGCT	120
TCACCCAAGG CGCCGACGGC AACGCCGGCA ACGGCGGTGA CGGCGGGGTC GGCGGCAACG	180
GCGGAAACGG CGGAAACGGC GCAGACAACA CCACCACCGC CGCCGCCGGC ACCACAGGCG	240
GCGACGGCGG GGCCGGCGGG GCCGGCGGAA CCGGCGGAAC CGGCGGAGCC GCCGGCACCG	300
GCACCGGCGG CCAACAAGGC AACGGCGGCA ACGGCGGCAC CGGCGGCAAA GGCGGCACCG	360
GCGGCGACGG TGCACTCTCA GGCAGCACCG GTGGTGCCGG	400

(2) INFORMATION FOR SEQ ID NO:180:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 538 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:180:

GGCAACGGCG GCAACGGCGG CATCGCCGGC ATTGGGCGGC AACGGCGTTC CGGGACGGGC	60
AGCGGCAACG GCGGCCAACG GCGGCAGCGG CGGCAACGGC GGCAACGCCG GCATGGGCGG	120
CAACAGCGGC ACCGGCAGCG GCGACGGCGG TGCCGGCGGG AACGGCGGCG CGGCGGGCAC	180
GGGCGGCACC GGCGGCGACG GCGGCCTCAC CGGTACTGGC GGCACCGGCG GCAGCGGTGG	240
CACCGGCGGT GACGGCGGTA ACGGCGGCAA CGGAGCAGAT AACACCGCAA ACATGACTGC	300
GCAGGCGGGC GGTGACGGTG GCAACGGCGG CGACGGTGCC TTCGGCGGCG GGGCCGGGGC	360
CGGCGGCGGT GGCTTGACCG CTGGCGCCAA CGGCACCGGC GGGCAAGGCG GCGCCGGCGG	420
CGATGGCGGC AACGGGGCCA TCGGCGGCCA CGGCCCACTC ACTGACGAEC CCGGCGGCAA	480
CGGGGGCACC GCGGCAACG GCGGCACCGG CGGCACCGGC GGCGCGGGCA TCGGCAGC	538

(2) INFORMATION FOR SEQ ID NO:181:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 239 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:181:

GGGCCGGTGG	TGCCGCGGGC	CAGCTCTTCA	GCGCCGGAGG	CGCGGCGGGT	GCCGTTGGGG	60
TTGGCGGCAC	CGGCGGCCAG	GGTGGGGCTG	GCGGTGCCCG	AGCGGCCGGC	GCCGACGCCC	120
CCGCCAGCAC	AGGTCTAACC	GGTGGTACCG	GGTTCGCTGG	CGGGGCCGGC	GGCGTCGGCG	180
GCCACGGCGG	CAACGCCATT	GCCGGCGGCA	TCAACGGCTC	CGGTGGTGCC	GGCGGCACC	239

(2) INFORMATION FOR SEQ ID NO:182:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 985 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:182:

AGCAGCGCTA	CCGGTGGCGC	CGGGTTCGCC	GGCGGCGCCG	GCGGAGAAGG	CGGAGCGGGC	60
GGCAACAGCG	GTGTGGGCGG	CACCAACGGC	TCCGGCGGCG	CCGGCGGTGC	AGGCGGCAAG	120
GGCGGCACCG	GAGGTGCCCG	CGGGTCCGGC	GCGGACAACC	CCACCGGTGC	TGGTTTCGCC	180
GGTGGCGCCG	GCGGCACAGG	TGGCGCGGCC	GGCGCCGGCG	GGGCCGGCGG	GGCGACCGGT	240
ACCGGCGGCA	CCGGCGGCGT	TGTCGGCGCC	ACCGGTAGTG	CAGGCATCGG	CGGGGCGGCG	300
GGCCGCGGCG	GTGACGGCGG	CGATGGGGCC	AGCGGTCTCG	GCCTGGGCCT	CTCCGGCTTT	360
GACGGCGGCC	AAGGCGGCCA	AGGCGGGGCC	GGCGGCAGCG	CCGGCGCCGG	CGGCATCAAC	420
GGGGCCGGCG	GGGCCGGCGG	CAACGGCGGC	GACGGCGGGG	ACGGCGCAAC	CGGTGCCGCA	480
GGTCTCGGCG	ACAACGGCGG	GGTCGGCGGT	GACGGTGGGG	CCGGTGGCGC	CGCCGGCAAC	540
GGCGGCAACG	CGGGCGTCGG	CCTGACAGCC	AAGGCCGGCG	ACGGCGGCGC	CGCGGGCAAT	600
GGCGGCAACG	GGGGCGCCGG	CGGTGCTGGC	GGGGCCGGCG	ACAACAATTT	CAACGGCGGC	660
CAGGGTGGTG	CCGGCGGCCA	AGGCGGCCAA	GGCGGCTTGG	GCGGGGCAAG	CACCACCTGA	720
TCGGCCTAGC	CGCACCCGGG	AAAGCCGATC	CAACAGGCGA	CGATGCCCGC	TTCTTGCCG	780
CGTTGGACCA	GGCCGGCATC	ACCTACGCTG	ACCCAGGCCA	CGCCATAACG	GCCGCCAAGG	840
CGATGTGTGG	GCTGTGTGCT	AACGGCGTAA	CAGGTCTACA	GCTGGTCGCG	GACCTGCGGG	900
AATACAATCC	CGGGCTGACC	ATGGACAGCG	CGGCCAAGTT	CGCTGCCATC	GCATCAGGCG	960
CGTACTGCCC	CGAACACCTG	GAACA				985

(2) INFORMATION FOR SEQ ID NO:183:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2138 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:183:

CGGCACGAGG ATCGGTACCC CGCGGCATCG GCAGCTGCCG ATTCGCCGGG TTTCCCCACC	60
CGAGGAAAGC CGCTACCAGA TGGCGCTGCC GAAGTAGGGC GATCCGTTTCG CGATGCCGGC	120
ATGAACGGGC GGCATCAAAT TAGTGAGGA ACCTTTCAGT TTAGCGACGA TAATGGCTAT	180
AGCACTAAGG AGGATGATCC GATATGACGC AGTCGCAGAC CGTGACGGTG GATCAGCAAG	240
AGATTTTGAA CAGGGCCAAC GAGGTGGAGG CCCCAGTGGC GGACCCACCG ACTGATGTCC	300
CCATCACACC GTGCGAACTC ACGGCGGCTA AAAACGCCGC CCAACAGCTG GTATTGTCCG	360
CCGACAACAT GCGGGAATAC CTGGCGGCCG GTGCCAAAGA GCGGCAGCGT CTGGCGACCT	420
CGCTGCGCAA CGCGGCCAAG GCGTATGGCG AGGTTGATGA GGAGGCTGCG ACCGCGCTGG	480
ACAACGACGG CGAAGGAACT GTGCAGGCAG AATCGGCCGG GGCCGTCGGA GGGGACAGTT	540
CGGCCGAAC TAAACGATACG CCGAGGGTGG CCACGGCCGG TGAACCCAAC TTCATGGATC	600
TCAAAGAAGC GGCAAGGAAG CTCGAAACGG GCGACCAAGG CGCATCGCTC GCGCACTTTG	660
CGGATGGGTG GAACACTTTC AACCTGACGC TGCAAGGCGA CGTCAAGCGG TTCCGGGGGT	720
TTGACAACTG GGAAGGCGAT GCGGCTACCG CTTGCGAGGC TTCGCTCGAT CAACAACGGC	780
AATGGATACT CCACATGGCC AAATTGAGCG CTGCGATGGC CAAGCAGGCT CAATATGTCTG	840
CGCAGCTGCA CGTGTGGGCT AGGCGGGAAC ATCCGACTTA TGAAGACATA GTCGGGCTCG	900
AACGGCTTTA CGCGGAAAAC CCTTCGGCCC GCGACCAAAT TCTCCCGGTG TACGCGGAGT	960
ATCAGCAGAG GTCGGAGAAG GTGCTGACCG AATACAACAA CAAGGCAGCC CTGGAACCGG	1020
TAAACCCGCC GAAGCCTCCC CCCGCCATCA AGATCGACCC GCCCCGCCT CCGCAAGAGC	1080
AGGGATTGAT CCCTGGCTTC GTGATGEGGC CGTCTGACGG CTCCGGTGTG ACTCCCGGTA	1140
CCGGGATGCC AGCCGCACCG ATGGTTCCGC CTACCGGATC GCCGGGTGGT GGCCTCCCGG	1200
CTGACACGGC GGCGCAGCTG ACGTCGGCTG GGCGGGAAGC CGCAGCGCTG TCGGGCGACG	1260
TGGCGGTCAA AGCGGCATCG CTCGGTGGCG GTGGAGGCGG CGGGGTGCCG TCGGCGCCGT	1320
TGGGATCCGC GATCGGGGGC GCCGAATCGG TGCGGCCCGC TGGCGCTGGT GACATTGCCG	1380
GCTTAGGCCA GGAAGGGGCC GCGGCGGGCG CCGCGCTGGG CGGCGGTGGC ATGGGAATGC	1440
CGATGGGTGC CGCGCATCAG GGACAAGGGG GCGCCAAGTC CAAGGGTTCT CAGCAGGAAG	1500

```

ACGAGGCGCT CTACACCGAG GATCGGGCAT GGACCGAGGC CGTCATTGGT AACCGTCGGC      1560
GCCAGGACAG TAAGGAGTCG AAGTGAGCAT GGACGAATTG GACCCGCATG TCGCCCGGGC      1620
GTTGACGCTG GCGGCGCGGT TTCAGTCGGC CCTAGACGGG ACGCTCAATC AGATGAACAA      1680
CGGATCCTTC CGCGCCACCG ACGAAGCCGA GACCGTCGAA GTGACGATCA ATGGGCACCA      1740
GTGGCTCACC GGCCTGCGCA TCGAAGATGG TTTGCTGAAG AAGCTGGGTG CCGAGGCGGT      1800
GGCTCAGCGG GTCAACGAGG CGCTGCACAA TGCGCAGGCC GCGGCGTCCG CGTATAACGA      1860
CGCGGCGGGC GAGCAGCTGA CCGCTGCGTT ATCGGCCATG TCCCGCGCGA TGAACGAAGG      1920
AATGGCCTAA GCCCATTGTT GCGGTGGTAG CGACTACGCA CCGAATGAGC GCCGCAATGC      1980
GGTCATTTCAG CGCGCCCGAC ACGGCGTGAG TACGCATTGT CAATGTTTTG ACATGGATCG      2040
GCCGGGTTCG GAGGGCGCCA TAGTCCTGGT CGCCAATATT GCCGCAGCTA GCTGGTCTTA      2100
GGTTCGGTTA CGCTGGTTAA TTATGACGTC CGTTACCA      2138

```

(2) INFORMATION FOR SEQ ID NO:184:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 460 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:184:

```

Met Thr Gln Ser Gln Thr Val Thr Val Asp Gln Gln Glu Ile Leu Asn
1           5           10           15
Arg Ala Asn Glu Val Glu Ala Pro Met Ala Asp Pro Pro Thr Asp Val
20          25          30
Pro Ile Thr Pro Cys Glu Leu Thr Ala Ala Lys Asn Ala Ala Gln Gln
35          40          45
Leu Val Leu Ser Ala Asp Asn Met Arg Glu Tyr Leu Ala Ala Gly Ala
50          55          60
Lys Glu Arg Gln Arg Leu Ala Thr Ser Leu Arg Asn Ala Ala Lys Ala
65          70          75          80
Tyr Gly Glu Val Asp Glu Glu Ala Ala Thr Ala Leu Asp Asn Asp Gly
85          90          95
Glu Gly Thr Val Gln Ala Glu Ser Ala Gly Ala Val Gly Gly Asp Ser
100         105         110
Ser Ala Glu Leu Thr Asp Thr Pro Arg Val Ala Thr Ala Gly Glu Pro
115         120         125
Asn Phe Met Asp Leu Lys Glu Ala Ala Arg Lys Leu Glu Thr Gly Asp
130         135         140

```


179

Gln Gly Ala Ser Leu Ala His Phe Ala Asp Gly Trp Asn Thr Phe Asn
 145 150 155 160
 Leu Thr Leu Gln Gly Asp Val Lys Arg Phe Arg Gly Phe Asp Asn Trp
 165 170 175
 Glu Gly Asp Ala Ala Thr Ala Cys Glu Ala Ser Leu Asp Gln Gln Arg
 180 185 190
 Gln Trp Ile Leu His Met Ala Lys Leu Ser Ala Ala Met Ala Lys Gln
 195 200 205
 Ala Gln Tyr Val Ala Gln Leu His Val Trp Ala Arg Arg Glu His Pro
 210 215 220
 Thr Tyr Glu Asp Ile Val Gly Leu Glu Arg Leu Tyr Ala Glu Asn Pro
 225 230 235 240
 Ser Ala Arg Asp Gln Ile Leu Pro Val Tyr Ala Glu Tyr Gln Gln Arg
 245 250 255
 Ser Glu Lys Val Leu Thr Glu Tyr Asn Asn Lys Ala Ala Leu Glu Pro
 260 265 270
 Val Asn Pro Pro Lys Pro Pro Pro Ala Ile Lys Ile Asp Pro Pro Pro
 275 280 285
 Pro Pro Gln Glu Gln Gly Leu Ile Pro Gly Phe Leu Met Pro Pro Ser
 290 295 300
 Asp Gly Ser Gly Val Thr Pro Gly Thr Gly Met Pro Ala Ala Pro Met
 305 310 315 320
 Val Pro Pro Thr Gly Ser Pro Gly Gly Gly Leu Pro Ala Asp Thr Ala
 325 330 335
 Ala Gln Leu Thr Ser Ala Gly Arg Glu Ala Ala Ala Leu Ser Gly Asp
 340 345 350
 Val Ala Val Lys Ala Ala Ser Leu Gly Gly Gly Gly Gly Gly Gly Val
 355 360 365
 Pro Ser Ala Pro Leu Gly Ser Ala Ile Gly Gly Ala Glu Ser Val Arg
 370 375 380
 Pro Ala Gly Ala Gly Asp Ile Ala Gly Leu Gly Gln Gly Arg Ala Gly
 385 390 395 400
 Gly Gly Ala Ala Leu Gly Gly Gly Gly Met Gly Met Pro Met Gly Ala
 405 410 415
 Ala His Gln Gly Gln Gly Gly Ala Lys Ser Lys Gly Ser Gln Gln Glu
 420 425 430
 Asp Glu Ala Leu Tyr Thr Glu Asp Arg Ala Trp Thr Glu Ala Val Ile
 435 440 445
 Gly Asn Arg Arg Arg Gln Asp Ser Lys Glu Ser Lys
 450 455 460

(2) INFORMATION FOR SEQ ID NO:185:

180

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 277 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:185:

Ala	Gly	Asn	Val	Thr	Ser	Ala	Ser	Gly	Pro	His	Arg	Phe	Gly	Ala	Pro	1	5	10	15
Asp	Arg	Gly	Ser	Gln	Arg	Arg	Arg	Arg	His	Pro	Ala	Ala	Ser	Thr	Ala	20	25	30	
Thr	Glu	Arg	Cys	Arg	Phe	Asp	Arg	His	Val	Ala	Arg	Gln	Arg	Cys	Gly	35	40	45	
Phe	Pro	Pro	Ser	Arg	Arg	Gln	Leu	Arg	Arg	Arg	Val	Ser	Arg	Glu	Ala	50	55	60	
Thr	Thr	Arg	Arg	Ser	Gly	Arg	Arg	Asn	His	Arg	Cys	Gly	Trp	His	Pro	65	70	75	80
Gly	Thr	Gly	Ser	His	Thr	Gly	Ala	Val	Arg	Arg	Arg	His	Gln	Glu	Ala	85	90	95	
Arg	Asp	Gln	Ser	Leu	Leu	Leu	Arg	Arg	Arg	Gly	Arg	Val	Asp	Leu	Asp	100	105	110	
Gly	Gly	Gly	Arg	Leu	Arg	Arg	Val	Tyr	Arg	Phe	Gln	Gly	Cys	Leu	Val	115	120	125	
Val	Val	Phe	Gly	Gln	His	Leu	Leu	Arg	Pro	Leu	Leu	Ile	Leu	Arg	Val	130	135	140	
His	Arg	Glu	Asn	Leu	Val	Ala	Gly	Arg	Arg	Val	Phe	Arg	Val	Lys	Pro	145	150	155	160
Phe	Glu	Pro	Asp	Tyr	Val	Phe	Ile	Ser	Arg	Met	Phe	Pro	Pro	Ser	Pro	165	170	175	
His	Val	Gln	Leu	Arg	Asp	Ile	Leu	Ser	Leu	Leu	Gly	His	Arg	Ser	Ala	180	185	190	
Gln	Phe	Gly	His	Val	Glu	Tyr	Pro	Leu	Pro	Leu	Leu	Ile	Glu	Arg	Ser	195	200	205	
Leu	Ala	Ser	Gly	Ser	Arg	Ile	Ala	Phe	Pro	Val	Val	Lys	Pro	Pro	Glu	210	215	220	
Pro	Leu	Asp	Val	Ala	Leu	Gln	Arg	Gln	Val	Glu	Ser	Val	Pro	Pro	Ile	225	230	235	240
Arg	Lys	Val	Arg	Glu	Arg	Cys	Ala	Leu	Val	Ala	Arg	Phe	Glu	Leu	Pro	245	250	255	
Cys	Arg	Phe	Phe	Glu	Ile	His	Glu	Val	Gly	Phe	Thr	Gly	Arg	Gly	His				

181

260 265 270

Pro Arg Arg Ile Gly
275

(2) INFORMATION FOR SEQ ID NO:186:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 192 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:186:

Arg	Val	Ala	Ala	Ser	Phe	Ile	Asp	Trp	Leu	Asp	Ser	Pro	Asp	Ser	Pro			
1				5					10					15				
Leu	Asp	Pro	Ser	Leu	Val	Ser	Ser	Leu	Leu	Asn	Ala	Val	Ser	Cys	Gly			
			20					25					30					
Ala	Glu	Ser	Ser	Ala	Ser	Ser	Ser	Ala	Arg	Ser	Gly	Asn	Gly	Ser	Arg			
		35					40					45						
Trp	Thr	Ser	Met	Pro	Ser	Gly	Thr	Arg	Pro	Gly	Pro	Arg	Arg	Ala	Thr			
	50					55					60							
Ser	Arg	Asp	Asp	Arg	Arg	Ser	Ala	Thr	Ser	Val	Ile	Pro	Ser	Arg	Arg			
65				70					75					80				
Ser	Val	Ala	Pro	Arg	Ala	Glu	Phe	Gly	Thr	Arg	Leu	Ala	Ser	His	Arg			
			85					90						95				
Ala	Ser	Pro	Ser	Asn	Ala	Cys	Pro	Val	Arg	Ile	Val	Thr	Ser	Ala	Ser			
			100					105					110					
Gly	Arg	Pro	Ile	Ser	Ser	Pro	Pro	Ile	Val	Arg	Ser	Arg	Ser	Cys	Val			
		115				120						125						
Asp	Lys	Asn	Gly	Arg	Arg	Cys	Ala	Ser	Gly	Tyr	Arg	Arg	Leu	Asn	Arg			
	130					135					140							
Ala	Arg	Ser	Ser	Ser	Ile	Ala	Ala	Arg	Cys	Arg	Thr	Ile	Gly	Thr	Phe			
145					150				155						160			
Arg	Arg	Ser	Arg	Tyr	Ser	Ala	Ser	Met	Arg	Val	Ser	Thr	Asn	Ser	Pro			
				165					170					175				
His	Val	Thr	His	Gly	Val	Ala	Pro	Gly	Val	Thr	Arg	Arg	Ile	Gly	Gly			
			180					185					190					

(2) INFORMATION FOR SEQ ID NO:187:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 196 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:

182

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:187:

Gln Glu Arg Pro Gln Met Cys Gln Arg Val Ser Glu Ile Glu Pro Arg
 1 5 10 15
 Thr Gln Phe Phe Asn Arg Cys Ala Leu Pro His Tyr Trp His Phe Pro
 20 25 30
 Ala Val Ala Val Phe Ser Lys His Ala Ser Leu Asp Glu Leu Ala Pro
 35 40 45
 Arg Asn Pro Arg Arg Ser Ser Arg Arg Asp Ala Glu Asp Arg Arg Val
 50 55 60
 Ile Phe Ala Ala Thr Leu Val Ala Val Asp Pro Pro Leu Arg Gly Ala
 65 70 75 80
 Gly Gly Glu Ala Asp Gln Leu Ile Asp Leu Gly Val Cys Arg Arg Gln
 85 90 95
 Ala Gly Arg Val Arg Arg Gly Gln Glu Leu His His Arg His Arg His
 100 105 110
 Gln Gly Ala Ala Pro Asp Leu Arg Arg Arg Arg Arg His Arg Arg Val
 115 120 125
 Gln Gln His Arg Arg Leu Gln Arg Val Arg Gln Leu Arg Arg Tyr Val
 130 135 140
 Gln Thr Ala His His Arg Arg Phe Ala Arg Thr Asp Arg Val Arg His
 145 150 155 160
 His Val Arg Gly Pro Ser Asn His Arg Arg Arg Arg Val Tyr Arg Gly
 165 170 175
 Arg His Ser Gly Ala Gly Gly Cys Pro Ala Gly Gly Ala Gly Ser Val
 180 185 190
 Gly Gly Ser Ala
 195

(2) INFORMATION FOR SEQ ID NO:188:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 311 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:188:

Val Arg Cys Gly Thr Leu Val Pro Val Pro Met Val Glu Phe Leu Thr

183

1	5	10	15
Ser Thr Asn Ala Pro Ser Leu Pro Ser Ala Tyr Ala Glu Val Asp Lys	20	25	30
Leu Ile Gly Leu Pro Ala Gly Thr Ala Lys Arg Trp Ile Asn Gly Tyr	35	40	45
Glu Arg Gly Gly Lys Asp His Pro Pro Ile Leu Arg Val Thr Pro Gly	50	55	60
Ala Thr Pro Trp Val Thr Trp Gly Glu Phe Val Glu Thr Arg Met Leu	65	70	75
Ala Glu Tyr Arg Asp Arg Arg Lys Val Pro Ile Val Arg Gln Arg Ala	85	90	95
Ala Ile Glu Glu Leu Arg Ala Arg Phe Asn Leu Arg Tyr Pro Leu Ala	100	105	110
His Leu Arg Pro Phe Leu Ser Thr His Glu Arg Asp Leu Thr Met Gly	115	120	125
Gly Glu Glu Ile Gly Leu Pro Asp Ala Glu Val Thr Ile Arg Thr Gly	130	135	140
Gln Ala Leu Leu Gly Asp Ala Arg Trp Leu Ala Ser Leu Val Pro Asn	145	150	155
Ser Ala Arg Gly Ala Thr Leu Arg Arg Leu Gly Ile Thr Asp Val Ala	165	170	175
Asp Leu Arg Ser Ser Arg Glu Val Ala Arg Arg Gly Pro Gly Arg Val	180	185	190
Pro Asp Gly Ile Asp Val His Leu Leu Pro Phe Pro Asp Leu Ala Asp	195	200	205
Asp Asp Ala Asp Asp Ser Ala Pro His Glu Thr Ala Phe Lys Arg Leu	210	215	220
Leu Thr Asn Asp Gly Ser Asn Gly Glu Ser Gly Glu Ser Ser Gln Ser	225	230	235
Ile Asn Asp Ala Ala Thr Arg Tyr Met Thr Asp Glu Tyr Arg Gln Phe	245	250	255
Pro Thr Arg Asn Gly Ala Gln Arg Ala Leu His Arg Val Val Thr Leu	260	265	270
Leu Ala Ala Gly Arg Pro Val Leu Thr His Cys Phe Ala Gly Lys Asp	275	280	285
Arg Thr Gly Phe Val Val Ala Leu Val Leu Glu Ala Val Gly Leu Asp	290	295	300
Arg Asp Val Ile Val Ala Asp	305	310	

(2) INFORMATION FOR SEQ ID NO:189:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2072 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:189:

CTCGTGCCGA TTCGGCACGA GCTGAGCAGC CCAAGGGGCC GTTCGGCGAA GTCATCGAGG	60
CATTCGCCGA CGGGCTGGCC GGCAAGGGTA AGCAAATCAA CACCACGCTG AACAGCCTGT	120
CGCAGGCGTT GAACGCCTTG AATGAGGGCC GCGGCGACTT CTTGCGGGTG GTACGCAGCC	180
TGGCGCTATT CGTCAACGCG CTACATCAGG ACGACCAACA GTTCGTGCGG TTGAACAAGA	240
ACCTTGCGGA GTTCACCGAC AGGTTGACCC ACTCCGATGC GGACCTGTCT AACGCCATCC	300
AGCAATTCGA CAGCTTGCTC GCCGTGCGCG GCCCGTTCTT CGCCAAGAAC CGCGAGGTGC	360
TGACGCATGA CGTCAATAAT CTCGCGACCG TGACCACCAC GTTGCTGCAG CCCGATCCGT	420
TGGATGGGTT GGAGACCGTC CTGCACATCT TCCCGACGCT GGCGGCGAAC ATTAACCAGC	480
TTTACCATCC GACACACGGT GGCGTGGTGT CGCTTTCCGC GTTCACGAAT TTCGCCAACC	540
CGATGGAGTT CATCTGCAGC TCGATTCAGG CGGGTAGCCG GCTCGGTTAT CAAGAGTCGG	600
CCGAACCTCTG TGCGCAGTAT CTGGCGCCAG TCCTCGATGC GATCAAGTTC AACTACTTTC	660
CGTTCGGCCT GAACGTGGCC AGCACCGCCT CGACACTGCC TAAAGAGATC GCGTACTCCG	720
AGCCCCGCTT GCAGCCGCCC AACGGGTACA AGGACACCAC GGTGCCCCGGC ATCTGGGTGC	780
CGGATACGCC GTTGTACAC ACCAACACGC AGCCCGGTTG GGTGGTGGCA CCCGGGATGC	840
AAGGGGTTCA GGTGGGACCG ATCACGCAGG GTTTGCTGAC GCCGGAGTCC CTGGCCGAAC	900
TCATGGGTGG TCCCGATATC GCCCCTCCGT CGTCAGGGCT GCAAACCCCG CCCGGACCCC	960
CGAATGCGTA CGACGAGTAC CCCGTGCTGC CGCCGATCGG TTTACAGGCC CCACAGGTGC	1020
CGATACCACC GCCGCCTCCT GGGCCCCGACG TAATCCCGGG TCCGGTGCCA CCGGTCTTGG	1080
CGGCGATCGT GTTCCCAAGA GATCGCCCCG CAGCGTCGGA AAACCTCGAC TACATGGGCC	1140
TCTTGTGTGCT GTCGCCGGGC CTGGCGACCT TCCTGTTCGG GGTGTCATCT AGCCCCGCCC	1200
GTGGAACGAT GGCCGATCGG CACGTGTTGA TACCGGCGAT CACCGGCCTG GCGTTGATCG	1260
CGGCATTCTG CGCACATTCT TGGTACCGCA CAGAACATCC GTCATAGAC ATGCGCTTGT	1320
TCCAGAACCG AGCGGTGCGG CAGGCCAACA TGACGATGAC GGTGCTCTCC CTCGGGCTGT	1380
TTGGCTCCTT CTTGCTGCTC CCGAGCTACC TCCAGCAAGT GTTGACCAA TCACCGATGC	1440
AATCGGGGGT GCATATCATC CCACAGGGCC TCGGTGCCAT GCTGGCGATG CCGATCGCCG	1500
GAGCGATGAT GGACCGACGG GGACCGCCA AGATCGTGCT GGTGGGATC ATGCTGATCG	1560

CTGCGGGGTT	GGGCACCTTC	GCCTTTGGTG	TCGCGCGGCA	AGCGGACTAC	TTACCCATTC	1620
TGCCGACCGG	GCTGGCAATC	ATGGGCATGG	GCATGGGCTG	CTCCATGATG	CCACTGTCCG	1680
GGGCGGCAGT	GCAGACCCTG	GCCCCACATC	AGATCGCTCG	CGGTTTCGACG	CTGATCAGCG	1740
TCAACCAGCA	GGTGGGCGGT	TCGATAGGGA	CCGCACTGAT	GTCGGTGCTG	CTCACCTACC	1800
AGTTCAATCA	CAGCGAAATC	ATCGCTACTG	CAAAGAAAGT	CGCACTGACC	CCAGAGAGTG	1860
GCGCCGGGCG	GGGGGCGGCG	GTTGACCCTT	CCTCGCTACC	GCGCCAAACC	AACTTCGCGG	1920
CCCAACTGCT	GCATGACCTT	TCGCACGCCT	ACGCGGTGGT	ATTCTGTGATA	GCGACCGCGC	1980
TAGTGGTCTC	GACGCTGATC	CCCGCGGCAT	TCCTGCCGAA	ACAGCAGGCT	AGTCATCGAA	2040
GAGCACCGTT	GCTATCCGCA	TGACGTCTGC	TT			2072

(2) INFORMATION FOR SEQ ID NO:190:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1923 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:190:

TCACCCCGGA	GAAGTCGTTC	GTCGACGACC	TGGACATCGA	CTCGCTGTCTG	ATGGTCGAGA	60
TCGCCGTGCA	GACCGAGGAC	AAGTACGGCG	TCAAGATCCC	CGACGAGGAC	CTCGCCGGTC	120
TGCGTACCGT	CGGTGACGTT	GTCGCCTACA	TCCAGAAGCT	CGAGGAAGAA	AACCCGGAGG	180
CGGCTCAGGC	GTTGCGCGCG	AAGATTGAGT	CGGAGAACCC	CGATGCGGCA	CGAGCAGATC	240
GGTGCGTTTC	ACCCACATCG	CAAGCTCGAG	ACGCCCCTCG	TCCTCTTGCA	CGCTCAGCCA	300
GGTTGGCGTG	TCGCCGCCTT	CCAGCAAGTG	TTCCCACCAC	ACGAAGGGAC	CCTCGCGAAA	360
GGTGACTGAT	CCGCGGACCA	CATAGTCGAT	GCCACCGTGG	CTGACAATTG	CGCCGGGTCC	420
GAGTTGGCGG	GGGCCGAATT	GCGGCATTGC	GTCGAAGGCC	AGCGGATCCC	GGCGCCCGCC	480
CGGCGTGGCT	GGTGTTTTGG	GCCGEEGGAT	GGCCACGACG	AGAACGACGA	TGGCGGCGAT	540
GAACAGCGCC	ACGGCAATCA	CGACCAGCAG	ATTTCCCACG	CATACCCTCT	CGTACCGCTG	600
CGCCGCGGTT	GGTCGATCGG	TCGCATATCG	ATGGCGCCGT	TTAACGTAAC	AGCTTTCGCG	660
GGACCGGGGG	TCACAACGGG	CGAGTTGTCC	GGCCGGGAAC	CCGGCAGGTC	TCGGCCGCGG	720
TCACCCAGC	TCACTGGTGC	ACCATCCGGG	TGTCGGTGAG	CGTGCAACTC	AAACACACTC	780
AACGGCAACG	GTTTCTCAGG	TCACCAGCTC	AACCTCGACC	CGCAATCGCT	CGTACGTTTC	840
GACCGCGCGC	AGGTCGCGAG	TCAGCAGCTT	TGCGCCGGCA	GCTTTCGCCG	TGAAGCCGAC	900

CAGGGCATCG TAGGTTGCGC CACCGGTGAC ATCGTGCTCG GCGAGGTGGT CGGTCAAGCC 960
 GCGATATGAG CAGGCATCCA GTGCCAGGTA GTTGCTGGAG GTGATGTCCG CCAAGTAGGC 1020
 GTGGACGGCA ACAGGGGCAA TACGATGCGG CGGTGGTAGC CGGGTCAAGA CCGAATAGGT 1080
 TTCCACAGCC GCGTGCGCGA TCAGATGGAC GCCACGGTTG AGCGCGCGCA CGGCGGCCTC 1140
 GTGCCCTTCG TGCCAGGTCG CGAATCCGGC AACCAGCACG CTGGTGTCTG GTGCGATCAC 1200
 CGCCGTGTGC GATCGAGCGT TTCCCGAACG ATTTCGTCGG TCAACGGGGG CAGGGGACGT 1260
 TCTGGCCGTG CGACGAGAAC CGAGCCTTCC CGAACGAGTT CGACACCGGT CGGGGCCGGC 1320
 TCAATCTCGA TGCGCCCATC GCGCTCGGTG ATCTCCACCT GGTCGTTCCC GCGCAAGCCA 1380
 AGGCGCTCGC GAATCCGCTT GGAATCACC AGACGTCCTG CGACATCGAT GGTTGTTCGC 1440
 ATGGTAGGAA ATTTACCATC GCACGTTCCA TAGGCGTGTC CTGCGCGGGA TGTCGGGACG 1500
 ATCCGCTAGC GTATCGAACG ATTGTTTCGG AAATGGCTGA GGGAGCGTGC GGTGCGGGTG 1560
 ATGGGTGTCG ATCCCGGGTT GACCCGATGC GGGCTGTGCG TCATCGAGAG TGGGCGTGGT 1620
 CGGCAGCTCA CCGCGCTGGA TGTCGACGTG GTGCGCACAC CGTCGGATGC GGCCTTGGCG 1680
 CAGCGCCTGT TGGCCATCAG CGATGCCGTC GAGCACTGGC TGGACACCCA TCATCCGGAG 1740
 GTGGTGGCTA TCGAACGGGT GTTCTCTCAG CTCAACGTGA CCACGGTGAT GGGCACCGCG 1800
 CAGGCCGGCG GCGTGATCGC CCTGGCGGCG GCCAAACGTG GTGTCGACGT GCATTTCCAT 1860
 ACCCCCAGCG AGGTCAAGGC GCGGTCACCT GGCAACGGTT CCGCAGACAA GGCTCAGGTC 1920
 ACC 1923

(2) INFORMATION FOR SEQ ID NO:191:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1055 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:191:

CTGGCGTGCC AGTGTCACCG GCGATATGAC GTCGGCATTC AATTTGCGCG CCCC GCCGGA 60
 CCCGTCGCCA CCAATCTGG ACCACCCGGT CCGTCAATTG CCGAAGGTCG CCAAGTGCGT 120
 GCCCAATGTG GTGCTGGGTT TCTTGAACGA AGGCCTGCCG TATCGGGTGC CCTACCCCCA 180
 AACAACGCCA GTCCAGGAAT CCGGTCCCGC GCGGCCGATT CCCAGCGGCA TCTGCTAGCC 240
 GGGGATGGTT CAGACGTAAC GGTGGCTAG GTCGAAACCC GCGCCAGGGC CGCTGGACGG 300
 GCTCATGGCA GCGAAATTAG AAAACCCGGG ATATTGTCCG CGGATTGTCA TACGATGCTG 360

AGTGCTTGGT GGTTCGTGTT TAGCCATTGA GTGTGGATGT GTTGAGACCC TGGCCTGGAA	420
GGGGACAACG TGCTTTTGCC TCTTGGTCCG CCTTTGCCGC CCGACGCGGT GGTGGCGAAA	480
CGGGCTGAGT CGGGAATGCT CGGCGGGTTG TCGGTTCCGC TCAGCTGGGG AGTGGCTGTG	540
CCACCCGATG ATTATGACCA CTGGGCGCCT GCGCCGGAGG ACGGCGCCGA TGTCGATGTC	600
CAGGCGGCCG AAGGGGCGGA CGCAGAGGCC GCGGCCATGG ACAGTGGA TGAGTGGCAG	660
GCGTGGAACG AGTGGGTGGC GGAGAACGCT GAACCCCGCT TTGAGGTGCC ACGGAGTAGC	720
AGCAGCGTGA TTCCGCATTC TCCGGCGGCC GGCTAGGAGA GGGGGCGCAG ACTGTCGTTA	780
TTTGACCACT GATCGGCGGT CTCGGTGTTT CCGCGGCCGG CTATGACAAC AGTCAATGTG	840
CATGACAAGT TACAGGTATT AGGTCCAGGT TCAACAAGGA GACAGGCAAC ATGGCAACAC	900
GTTTTATGAC GGATCCGCAC GCGATGCGGG ACATGGCGGG CCGTTTTGAG GTGCACGCCC	960
AGACGGTGGA GGACGAGGCT CGCCGGATGT GGGCGTCCGC GCAAAACATC TCGGGNGCGG	1020
GCTGGAGTGG CATGGCCGAG GCGACCTCGC TAGAC	1055

(2) INFORMATION FOR SEQ ID NO:192:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 359 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:192:

CCGCCTCGTT GTTGGCATA TCCGCCGCGG CCGCCTCGAC CGCACTGGCC GTGGCGTGTG	60
TCCGGGCTGA CCACCGGGAT CGCCGAACCA TCCGAGATCA CCTCGCAATG ATCCACCTCG	120
CGCAGCTGGT CACCCAGCCA CCGGGCGGTG TGCACAGCG CCTGCATCAC CTTGGTATAG	180
CCGTCGCGCC CCAGCCGCAG GAAGTTGTAG TACTGGCCCA CCACCTGGTT ACCGGGACGG	240
GAGAAGTTCA GGGTGAAGGT CGGCATGTCG CCGCCGAGGT AGTTGACCCG GAAAACCAGA	300
TCCTCCGGCA GGTGCTCGGG CCCGCGCCAC ACGACAAACC CGACGCGGG ATAGGTCAG	359

(2) INFORMATION FOR SEQ ID NO:193:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 350 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:193:

```

AACGGGCCCCG TGGGCACCGC TCCTCTAAGG GCTCTCGTTG GTCGCATGAA GTGCTGGAAG      60
GATGCATCTT  GGCAGATTCC CGCCAGAGCA AAACAGCCGC TAGTCCTAGT CCGAGTCGCC      120
CGCAAAGTTC CTCGAATAAC TCCGTACCCG GAGCGCCAAA CCGGGTCTCC TTCGCTAAGC      180
TGCGCGAACC ACTTGAGGTT CCGGGACTCC TTGACGTCCA GACCGATTCTG TTCGAGTGGC      240
TGATCGGTTT GCCGCGCTGG CGCGAATCCG CCGCCGAGCG GGGTGATGTC AACCCAGTGG      300
GTGGCCTGGA AGAGGTGCTC TACGAGCTGT CTCCGATCGA GGACTTCTCC      350

```

(2) INFORMATION FOR SEQ ID NO:194:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 679 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:194:

```

Glu Gln Pro Lys Gly Pro Phe Gly Glu Val Ile Glu Ala Phe Ala Asp
1           5           10           15
Gly Leu Ala Gly Lys Gly Lys Gln Ile Asn Thr Thr Leu Asn Ser Leu
20        25        30
Ser Gln Ala Leu Asn Ala Leu Asn Glu Gly Arg Gly Asp Phe Phe Ala
35        40        45
Val Val Arg Ser Leu Ala Leu Phe Val Asn Ala Leu His Gln Asp Asp
50        55        60
Gln Gln Phe Val Ala Leu Asn Lys Asn Leu Ala Glu Phe Thr Asp Arg
65        70        75        80
Leu Thr His Ser Asp Ala Asp Leu Ser Asn Ala Ile Gln Gln Phe Asp
85        90        95
Ser Leu Leu Ala Val Ala Arg Pro Phe Phe Ala Lys Asn Arg Glu Val
100       105       110
Leu Thr His Asp Val Asn Asn Leu Ala Thr Val Thr Thr Thr Leu Leu
115       120       125
Gln Pro Asp Pro Leu Asp Gly Leu Glu Thr Val Leu His Ile Phe Pro
130       135       140
Thr Leu Ala Ala Asn Ile Asn Gln Leu Tyr His Pro Thr His Gly Gly
145       150       155       160
Val Val Ser Leu Ser Ala Phe Thr Asn Phe Ala Asn Pro Met Glu Phe
165       170       175

```

189

Ile	Cys	Ser	Ser	Ile	Gln	Ala	Gly	Ser	Arg	Leu	Gly	Tyr	Gln	Glu	Ser	
			180					185					190			
Ala	Glu	Leu	Cys	Ala	Gln	Tyr	Leu	Ala	Pro	Val	Leu	Asp	Ala	Ile	Lys	
		195					200					205				
Phe	Asn	Tyr	Phe	Pro	Phe	Gly	Leu	Asn	Val	Ala	Ser	Thr	Ala	Ser	Thr	
	210					215					220					
Leu	Pro	Lys	Glu	Ile	Ala	Tyr	Ser	Glu	Pro	Arg	Leu	Gln	Pro	Pro	Asn	
225					230					235					240	
Gly	Tyr	Lys	Asp	Thr	Thr	Val	Pro	Gly	Ile	Trp	Val	Pro	Asp	Thr	Pro	
				245					250					255		
Leu	Ser	His	Arg	Asn	Thr	Gln	Pro	Gly	Trp	Val	Val	Ala	Pro	Gly	Met	
			260					265					270			
Gln	Gly	Val	Gln	Val	Gly	Pro	Ile	Thr	Gln	Gly	Leu	Leu	Thr	Pro	Glu	
		275					280					285				
Ser	Leu	Ala	Glu	Leu	Met	Gly	Gly	Pro	Asp	Ile	Ala	Pro	Pro	Ser	Ser	
	290					295					300					
Gly	Leu	Gln	Thr	Pro	Pro	Gly	Pro	Pro	Asn	Ala	Tyr	Asp	Glu	Tyr	Pro	
305					310					315					320	
Val	Leu	Pro	Pro	Ile	Gly	Leu	Gln	Ala	Pro	Gln	Val	Pro	Ile	Pro	Pro	
				325					330					335		
Pro	Pro	Pro	Gly	Pro	Asp	Val	Ile	Pro	Gly	Pro	Val	Pro	Pro	Val	Leu	
			340					345					350			
Ala	Ala	Ile	Val	Phe	Pro	Arg	Asp	Arg	Pro	Ala	Ala	Ser	Glu	Asn	Phe	
		355					360					365				
Asp	Tyr	Met	Gly	Leu	Leu	Leu	Leu	Ser	Pro	Gly	Leu	Ala	Thr	Phe	Leu	
	370					375					380					
Phe	Gly	Val	Ser	Ser	Ser	Pro	Ala	Arg	Gly	Thr	Met	Ala	Asp	Arg	His	
385					390					395					400	
Val	Leu	Ile	Pro	Ala	Ile	Thr	Gly	Leu	Ala	Leu	Ile	Ala	Ala	Phe	Val	
				405					410					415		
Ala	His	Ser	Trp	Tyr	Arg	Thr	Glu	His	Pro	Leu	Ile	Asp	Met	Arg	Leu	
			420					425					430			
Phe	Gln	Asn	Arg	Ala	Val	Ala	Gln	Ala	Asn	Met	Thr	Met	Thr	Val	Leu	
		435					440					445				
Ser	Leu	Gly	Leu	Phe	Gly	Ser	Phe	Leu	Leu	Leu	Pro	Ser	Tyr	Leu	Gln	
	450					455					460					
Gln	Val	Leu	His	Gln	Ser	Pro	Met	Gln	Ser	Gly	Val	His	Ile	Ile	Pro	
465					470					475					480	
Gln	Gly	Leu	Gly	Ala	Met	Leu	Ala	Met	Pro	Ile	Ala	Gly	Ala	Met	Met	
				485					490					495		
Asp	Arg	Arg	Gly	Pro	Ala	Lys	Ile	Val	Leu	Val	Gly	Ile	Met	Leu	Ile	
			500					505					510			

190

Ala Ala Gly Leu Gly Thr Phe Ala Phe Gly Val Ala Arg Gln Ala Asp
 515 520 525

Tyr Leu Pro Ile Leu Pro Thr Gly Leu Ala Ile Met Gly Met Gly Met
 530 535 540

Gly Cys Ser Met Met Pro Leu Ser Gly Ala Ala Val Gln Thr Leu Ala
 545 550 555 560

Pro His Gln Ile Ala Arg Gly Ser Thr Leu Ile Ser Val Asn Gln Gln
 565 570 575

Val Gly Gly Ser Ile Gly Thr Ala Leu Met Ser Val Leu Leu Thr Tyr
 580 585 590

Gln Phe Asn His Ser Glu Ile Ile Ala Thr Ala Lys Lys Val Ala Leu
 595 600 605

Thr Pro Glu Ser Gly Ala Gly Arg Gly Ala Ala Val Asp Pro Ser Ser
 610 615 620

Leu Pro Arg Gln Thr Asn Phe Ala Ala Gln Leu Leu His Asp Leu Ser
 625 630 635 640

His Ala Tyr Ala Val Val Phe Val Ile Ala Thr Ala Leu Val Val Ser
 645 650 655

Thr Leu Ile Pro Ala Ala Phe Leu Pro Lys Gln Gln Ala Ser His Arg
 660 665 670

Arg Ala Pro Leu Leu Ser Ala
 675

(2) INFORMATION FOR SEQ ID NO:195:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 120 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:195:

Thr Pro Glu Lys Ser Phe Val Asp Asp Leu Asp Ile Asp Ser Leu Ser
 1 5 10 15

Met Val Glu Ile Ala Val Gln Thr Glu Asp Lys Tyr Gly Val Lys Ile
 20 25 30

Pro Asp Glu Asp Leu Ala Gly Leu Arg Thr Val Gly Asp Val Val Ala
 35 40 45

Tyr Ile Gln Lys Leu Glu Glu Asn Pro Glu Ala Ala Gln Ala Leu
 50 55 60

Arg Ala Lys Ile Glu Ser Glu Asn Pro Asp Ala Ala Arg Ala Asp Arg
 65 70 75 80

Cys Val Ser Pro Thr Ser Gln Ala Arg Asp Ala Arg Arg Pro Leu Ala
85 90 95

Arg Ser Ala Arg Leu Ala Cys Arg Arg Leu Pro Ala Ser Val Pro Thr
100 105 110

Thr Arg Arg Asp Pro Arg Glu Arg
115 120

(2) INFORMATION FOR SEQ ID NO:196:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 89 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:196:

```

Leu Ala Cys Gln Cys His Arg Arg Tyr Asp Val Gly Ile Gln Phe Arg
1      5      10     15
Gly Pro Ala Gly Pro Val Ala Thr Gln Ser Gly Pro Pro Gly Pro Ser
20    25    30
Ile Ala Glu Gly Arg Gln Val Arg Ala Gln Cys Gly Ala Gly Phe Leu
35    40    45
Glu Arg Arg Pro Ala Val Ser Gly Ala Leu Pro Pro Asn Asn Ala Ser
50    55    60
Pro Gly Ile Arg Ser Arg Ala Ala Asp Ser Gln Arg His Leu Leu Ala
65    70    75    80
Gly Asp Gly Ser Asp Val Thr Val Gly
85

```

(2) INFORMATION FOR SEQ ID NO:197:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 119 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:197:

Ala Ser Leu Leu Ala Tyr Ser Ala Ala Ala Ala Ser Thr Ala Leu Ala
1 5 10 15
Val Ala Cys Val Arg Ala Asp His Arg Asp Arg Arg Thr Ile Arg Asp
20 25 30

192

His Leu Ala Met Ile His Leu Ala Gln Leu Val Thr Gln Pro Pro Gly
 35 40 45
 Gly Val Arg Gln Arg Leu His His Leu Gly Ile Ala Val Ala Pro Gln
 50 55 60
 Pro Gln Glu Val Val Val Leu Ala His His Leu Val Thr Gly Thr Gly
 65 70 75 80
 Glu Val Gln Gly Glu Gly Arg His Val Ala Ala Glu Val Val Asp Pro
 85 90 95
 Glu Asn Gln Ile Leu Arg Gln Val Leu Gly Pro Ala Pro His Asp Lys
 100 105 110
 Pro Asp Ala Gly Ile Gly Gln
 115

(2) INFORMATION FOR SEQ ID NO:198:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 116 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:198:

Arg Ala Arg Gly His Arg Ser Ser Lys Gly Ser Arg Trp Ser His Glu
 1 5 10 15
 Val Leu Glu Gly Cys Ile Leu Ala Asp Ser Arg Gln Ser Lys Thr Ala
 20 25 30
 Ala Ser Pro Ser Pro Ser Arg Pro Gln Ser Ser Ser Asn Asn Ser Val
 35 40 45
 Pro Gly Ala Pro Asn Arg Val Ser Phe Ala Lys Leu Arg Glu Pro Leu
 50 55 60
 Glu Val Pro Gly Leu Leu Asp Val Gln Thr Asp Ser Phe Glu Trp Leu
 65 70 75 80
 Ile Gly Ser Pro Arg Trp Arg Glu Ser Ala Ala Glu Arg Gly Asp Val
 85 90 95
 Asn Pro Val Gly Gly Leu Glu Glu Val Leu Tyr Glu Leu Ser Pro Ile
 100 105 110
 Glu Asp Phe Ser
 115

(2) INFORMATION FOR SEQ ID NO:199:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 811 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:199:

TGCTACGCAG CAATCGCTTT GGTGACAGAT GTGGATGCCG GCGTCGCTGC TGGCGATGGC	60
GTGAAAGCCG CCGACGTGTT CGCCGCATTC GGGGAGAACA TCGAACTGCT CAAAAGGCTG	120
GTGCGGGCCG CCATCGATCG GGTGCGCGAC GAGCGCACGT GCACGCACTG TCAACACCAC	180
GCCGGTGTTT CGTTGCCGTT CGAGCTGCCA TGAGGGTGCT GCTGACCGGC GCGGCCGGCT	240
TCATCGGGTC GCGCGTGGAT GCGGCGTTAC GGGCTGCGGG TCACGACGTG GTGGGCGTCG	300
ACGCGCTGCT GCGCGCGCG CACGGGCCAA ACCCGGTGCT GCCACCGGGC TGCCAGCGGG	360
TCGACGTGCG CGACGCCAGC GCGCTGGCCC CGTTGTTGGC CGGTGTCGAT CTGGTGTGTC	420
ACCAGGCCGC CATGGTGGGT GCCGGCGTCA ACGCCGCCGA CGCACCCGCC TATGGCGGCC	480
ACAACGATTT CGCCACCACG GTGCTGCTGG CGCAGATGTT CGCCGCCGGG GTCCGCCGTT	540
TGGTGCTGGC GTCGTCGATG GTGGTTTACG GGCAGGGGCG CTATGACTGT CCCCAGCATG	600
GACCGGTGCA CCCGCTGCCG CGGCGGCGAG CCGACCTGGA CAATGGGGTC TTCGAGCACC	660
GTTGCCCCGG GTGCGGCGAG CCAGTCATCT GGCAATTGGT CGACGAAGAT GCCCCGTTGC	720
GCCCCGCGCAG CCTGTACGCG GCAGCAAGAC CGCGCAGGAG CACTACGCGC TGGCGTGGTC	780
GGAAACGAAT GCGGTTCCG TGGTGGCGTT G	811

(2) INFORMATION FOR SEQ ID NO:200:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 966 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:200:

GTCCCGCGAT GTGGCCGAGC ATGACTTTTCG GCAACACCGG CGTAGTAGTC GAAGATATCG	60
GACTTTGTGG TCCCGGTGGC GGGATAGAGC ACCTGTCGGC GTTGGTCAGC GTCACCCGTT	120
GCTCGGACGC CGAACCCATG CTTTCAACGT AGCCTGTCGG TCACACAAGT CGCGAGCGTA	180
ACGTCACGGT CAAATATCGC GTGGAATTTC GCCGTGACGT TCCGCTCGCG GACAATCAAG	240
GCATACTCAC TTACATGCGA GCCATTTGGA CGGGTTCGAT CGCCTTCGGG CTGGTGAACG	300
TGCCGGTCAA GGTGTACAGC GCTACCGCAG ACCACGACAT CAGGTTCCAC CAGGTGCACG	360

CCAAGGACAA CGGACGCATC CGGTACAAGC GCGTCTGCGA GGCGTGTGGC GAGGTGGTCG	420
ACTACCGCGA TCTTGCCCGG GCCTACGAGT CCGGCGACGG CCAAATGGTG GCGATCACCG	480
ACGACGACAT CGCCAGCTTG CCTGAAGAAC GCAGCCGGGA GATCGAGGTG TTGGAGTTCTG	540
TCCCCGCCGC CGACGTGGAC CCGATGATGT TCGACCGCAG CTACTTTTTG GAGCCTGATT	600
CGAAGTCGTC GAAATCGTAT GTGCTGCTGG CTAAGACACT CGCCGAGACC GACCGGATGG	660
CGATCGTGGA TCGCCCCACC GGCCGTGAAT GCAGGAAAAA TAAGAGCCGC TATCCACAAT	720
TCGGCGTCGA GCTCGGTAC CACAAACGGT AGAACGATCG AGACATTCCC GAGCTGAAGT	780
GCGGCGCTAT AGAAGCCGCT CTGCGCGATT ATCAAACGCA AAATACGCTT ACTCATGCCA	840
TCGGCGCTGC TCACCCGATG CGACGTTTTT GCCACGCTCC ACCGCCTGCC GCGCGACCTC	900
AAGTGGGCAT GCATCCCACC CGTTCCCGGA AACCGGTTCC GCGGGGTCGG CTCATCGCTT	960
CATCCT	966

(2) INFORMATION FOR SEQ ID NO:201:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2367 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:201:

CCGCACCGCC GGCAATACCG CCAGCGCCAC CGTTACCGCC GTTGTGCGCCG TTGCCCCCGT	60
TGCGCGCCGT CCCGCCGGCC CCGCCGATGG AGTTCTCATC GCCAAAAGTA CTGGCGTTGC	120
CACEGGAGCC GCCGTTGCCG CCGTCACCGC CAGCCCCGCC GACTCCACCG GCCCCACCGA	180
CTCGGCCGCT GCCACCGTTG CCGCCGTTGC CGATCAACAT GCCGCTGGCG CCACCCTTGC	240
CACCCACGCC ACCGGCTCCG CCCACCCCGC CGACACCAAG CGAGCTGCCG CCGGAGCCAC	300
CATCACCACC TACGCCACCG ACCGCCGAGA CACCAGCGAC CGGGTCTTCG TGAAACGTCG	360
CGGTGCCACC ACCGCCGCCG TTACCGCCAA CCCACCGGC AACGCCGGCG CCGCCATCCC	420
CGCCGGCCCC GCGGTTGCCG CCGTTGCCGC CGTTGCCGAA CAACAACCCG CCGGCGCCGC	480
CGTTGCCGCC CGCGCCGCCG GTCCCGCCGG CGCCGCCGAC GCAAGGCCG CTGCCGCCCT	540
TGCCGCCATC ACCACCCTTG CCGCCGACCA CATCGGGTTC TGCCTCGGGG TCTGGGCTGT	600
CAAACCTCGC GATGCCAGCG TTGCCGCCG TTCCCCCGG CCCCCCGTG GCGCCGTCAC	660
CACCGATACC ACCCGCGCCA CCGGCGCCAC CGTTGCCGCC ATCACCGAAT AGCAACCCGC	720
CGGCGCCACC ATTGCCGCA GCTCCCCCTG CGCCACCGTC GGCGCCGGAG GCGGCACTGG	780

CAGCCCCGTT	ACCACCGAAA	CCGCCGCTAC	CACCGGTAGA	GGTGGCAGTG	GCGATGTGTA	840
CGAAAGCGCC	GCCTCCGGCG	CCGCCGCTAC	CACCCCCACT	GCCGGCGGGT	ACACCGTCGG	900
ACCCGTTGCC	ACCATCACCG	CCAAAGGCGC	TCGCAATGTC	GCCCTGCGCG	ACTCCGCCGT	960
CGCCGCCGTT	GCCGCCGCGG	CCACCGGCAG	CGGCGGTACC	GCCGTCACCA	CCGGCACC GC	1020
CGGTGGCCTT	GCCCCAGCCT	GCCGTCGCGG	TGGCACCGTC	GCCGCCGGTG	CCACCGGTGC	1080
GCGTGCCGGC	AGTGCCATGG	CCGCCCGTGC	CGCCGTCGCC	GCCGGTTTGA	TCACCGATGC	1140
CGGACACATC	TGCCGGGCTG	TCCCCGGTGC	TGGCCGCGGG	GCCGGGCGTG	GGATTGACCC	1200
CGTTTGCCCC	GGCGAGGCCG	GCGCCGCCGG	TACCACCGGC	GCCGCCATGG	CCGAACAGCC	1260
CGGCGTTGCC	GCCGTTACCG	CCCGCACCCC	CGATGCCTGC	GGCCACGCTG	GTGCCGCCGA	1320
CACCGCCGTT	GCCGCCGTTG	CCCCACAACC	ACCCCCCGTT	CCCACCGGCA	CCGCCGGCCG	1380
CGCCGGTACC	ACCGGCCCCG	CCGTTGCCGC	CGTTGCCGAT	CAACCCGGCC	GCGCCTCCGC	1440
TGCCGCCGGT	TTGACCGAAC	CCGCCAGCCG	CGCCGTTGCC	ACCGTTGCCA	AACAGCAACC	1500
CGCCGGCCGC	GCCAGGCTGC	CCGGGTGCCG	TCCCGTCGGC	GCCGTTTCCG	ATCAACGGGC	1560
GCCCCAAAAG	CGCCTCGGTG	GGCGCATTCA	CCGCACCCAG	CAGACTCCGC	TCAACAGCGG	1620
CTTCAGTGCT	GGCATAACCGA	CCCGCGGCCG	CAGTCAACGC	CTGCACAAAC	TGCTCGTGAA	1680
ACGCTGCCAC	CTGTACGCTG	AGCGCCTGAT	ACTGCCGAGC	ATGGGCCCCG	AACAACCCCG	1740
CAATCGCCGC	CGACACTTCA	TCGGCAGCCG	CAGCCACCAC	TTCCGTCGTC	GGGATCGCCG	1800
CGGCCGCATT	AGCCGCGCTC	ACCTGCGAAC	CAATAGTCGA	TAAATCCAAA	GCCGCAGTTG	1860
CCAGCAGCTG	CGGCGTCGCG	ATCACCAAGG	ACACCTCGCA	CCTCCGGATA	CCCCATATCG	1920
CCGCACCGTG	TCCCCAGCGG	CCACGTGACC	TTTGGTCGCT	GGCTGGCGGC	CCTGACTATG	1980
GCCGCGACGG	CCCTCGTTCT	GATTCGCCCC	GGCGCGCAGC	TTGTTGCGCG	AGTTGAAGAC	2040
GGGAGGACAG	GCCGAGCTTG	GTGTAGACGT	GGGTCAAGTG	GGAATGCACG	GTCCGCGGCG	2100
AGATGAATAG	GCGGACGCCG	ATCTCCTTGT	TGCTGAGTCC	CTCACCAGAC	AGTAGAGCCA	2160
CCTCAAGCTC	TGTCGGTGTC	AACGCGCCCC	AGCCACTTGT	CGGGCGTTTC	CGTGCACCGC	2220
GGCCTCGTTG	GGCGTAEGEG	ATCGCCTCAT	CGATCGATAA	CGCAGTTCCT	TCGGCCCAGG	2280
CATCGTCGAA	CTCGCTGTCA	CCCATGGATT	TTCGAAGGGT	GGCTAGCGAC	GAGTTACAGC	2340
CCGCCTGGTA	GATCCCGAAG	CGGACCG				2367

(2) INFORMATION FOR SEQ ID NO:202:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 376 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:202:

Gln Pro Ala Gly Ala Thr Ile Ala Ala Ser Ser Pro Cys Ala Thr Val
 1 5 10 15
 Gly Ala Gly Gly Gly Thr Gly Ser Pro Val Thr Thr Glu Thr Ala Ala
 20 25 30
 Thr Thr Gly Arg Gly Gly Ser Gly Asp Val Tyr Glu Ser Ala Ala Ser
 35 40 45
 Gly Ala Ala Ala Thr Thr Pro Thr Ala Gly Gly Tyr Thr Val Gly Pro
 50 55 60
 Val Ala Thr Ile Thr Ala Lys Gly Ala Arg Asn Val Ala Leu Arg Asp
 65 70 75 80
 Ser Ala Val Ala Ala Val Ala Ala Ala Ala Thr Gly Ser Gly Gly Thr
 85 90 95
 Ala Val Thr Thr Gly Thr Ala Gly Gly Leu Ala Arg Ala Cys Arg Arg
 100 105 110
 Gly Gly Thr Val Ala Ala Gly Ala Thr Gly Arg Arg Ala Gly Ser Ala
 115 120 125
 Met Ala Ala Arg Ala Ala Val Ala Ala Gly Leu Ile Thr Asp Ala Gly
 130 135 140
 His Ile Cys Arg Ala Val Pro Gly Ala Gly Arg Gly Ala Gly Arg Gly
 145 150 155 160
 Ile Asp Pro Val Cys Pro Gly Glu Ala Gly Ala Ala Gly Thr Thr Gly
 165 170 175
 Ala Ala Met Ala Glu Gln Pro Gly Val Ala Ala Val Thr Ala Arg Thr
 180 185 190
 Pro Asp Ala Cys Gly His Ala Gly Ala Ala Asp Thr Ala Val Ala Ala
 195 200 205
 Val Ala Pro Gln Pro Pro Pro Val Pro Thr Gly Thr Ala Gly Arg Ala
 210 215 220
 Gly Thr Thr Gly Pro Ala Val Ala Ala Val Ala Asp Gln Pro Gly Arg
 225 230 235 240
 Ala Ser Ala Ala Ala Gly Leu Thr Glu Pro Ala Ser Arg Ala Val Ala
 245 250 255
 Thr Val Ala Lys Gln Gln Pro Ala Gly Arg Ala Arg Leu Pro Gly Cys
 260 265 270
 Arg Pro Val Gly Ala Val Ser Asp Gln Arg Ala Pro Gln Lys Arg Leu
 275 280 285
 Gly Gly Arg Ile His Arg Thr Gln Gln Thr Pro Leu Asn Ser Gly Phe
 290 295 300

197

Ser Ala Gly Ile Pro Thr Arg Gly Arg Ser Gln Arg Leu His Lys Leu
 305 310 315 320

Leu Val Lys Arg Cys His Leu Tyr Ala Glu Arg Leu Ile Leu Pro Ser
 325 330 335

Met Gly Pro Glu Gln Pro Arg Asn Arg Arg Arg His Phe Ile Gly Ser
 340 345 350

Arg Ser His His Phe Arg Arg Arg Asp Arg Arg Gly Arg Ile Ser Arg
 355 360 365

Ala His Leu Arg Thr Asn Ser Arg
 370 375

(2) INFORMATION FOR SEQ ID NO:203:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2852 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:203:

GGCCAAAACG CCCCGGCGAT CGCGGCCACC GAGGCCGCCT ACGACCAGAT GTGGGCCCAG	60
GACGTGGCGG CGATGTTTGG CTACCATGCC GGGGCTTCGG CGGCCGTCTC GGCGTTGACA	120
CCGTTTCGGCC AGGCGCTGCC GACCGTGGCG GCGGGCGGTG CGCTGGTCAG CGCGGCCGCG	180
GCTCAGGTGA CCACGCGGGT CTTCCGCAAC CTGGGCTTGG CGAACGTCCG CGAGGGCAAC	240
GTCCGCAACG GTAATGTCCG GAACTTCAAT CTCGGCTCGG CCAACATCGG CAACGGCAAC	300
ATCGGCAGCG GCAACATCGG CAGCTCCAAC ATCGGGTTTG GCAACGTGGG TCCTGGGTTG	360
ACCGCAGCGC TGAACAACAT CGGTTTCGGC AACACCGGCA GCAACAACAT CGGGTTTGGC	420
AACACCGGCA GCAACAACAT CGGGTTCGGC AATACCGGAG ACGGCAACCG AGGTATCGGG	480
CTCACGGGTA GCGGTTTGTT GGGGTTTCGGC GGCCTGAACT CGGGCACCGG CAACATCGGT	540
CTGTTCAACT CGGGCACCGG AAACGTCGGG ATCGGCAACT CGGGTACCGG GAACTGGGGC	600
ATTGGCAACT CGGGCAACAG CTACAACACC GGTTTGGCA ACTCCGGCGA CGCCAACACG	660
GGCTTCTTCA ACTCCGGAAT AGCCAACACC GGCGTCGGCA ACGCCGGCAA CTACAACACC	720
GGTAGCTACA ACCCGGGCAA CAGCAATACC GGCGGCTTCA ACATGGGCCA GTACAACACG	780
GGCTACCTGA ACAGCGGCAA CTACAACACC GGCTTGGCAA ACTCCGGCAA TGTC AACACC	840
GGCGCCTTCA TTA CTGGCAA CTTCAACAAC GGCTTCTTGT GGCGCGGCGA CCACCAAGGC	900
CTGATTTTCG GGAGCCCCGG CTTCTTCAAC TCGACCAAGT CGCCGTCGTC GGGATTCTTC	960

AACAGCGGTG	CCGGTAGCGC	GTCCGGCTTC	CTGAACTCCG	GTGCCAACAA	TTCTGGCTTC	1020
TTCAACTCTT	CGTCGGGGGC	CATCGGTAAC	TCCGGCCTGG	CAAACGCGGG	CGTGCTGGTA	1080
TCGGGCGTGA	TCAACTCGGG	CAACACCGTA	TCGGGTTTGT	TCAACATGAG	CCTGGTGGCC	1140
ATCACAAACGC	CGGCCTTGAT	CTCGGGCTTC	TTCAACACCG	GAAGCAACAT	GTCGGGATTT	1200
TTCGGTGGCC	CACCGGTCTT	CAATCTCGGC	CTGGCAAACC	GGGGCGTCGT	GAACATTCTC	1260
GGCAACGCCA	ACATCGGCAA	TTACAACATT	CTCGGCAGCG	GAAACGTCGG	TGACTTCAAC	1320
ATCCTTGCCA	GCGGCAACCT	CGGCAGCCAA	AACATCTTGG	GCAGCGGCAA	CGTCGGCAGC	1380
TTCAATATCG	GCAGTGAAA	CATCGGAGTA	TTCAATGTCG	GTTCCGGAAG	CCTGGGAAAC	1440
TACAACATCG	GATCCGAAA	CCTCGGGATC	TACAACATCG	GTTTTGAAA	CGTCGGCGAC	1500
TACAACGTCG	GCTTCGGGAA	CGCGGGCGAC	TTCAACCAAG	GCTTTGCCAA	CACCGGCAAC	1560
AACAACATCG	GGTTCGCCAA	CACCGGCAAC	AACAACATCG	GCATCGGGCT	GTCCGGCGAC	1620
AACCAGCAGG	GCTTCAATAT	TGCTAGCGGC	TGGAATCGG	GCACCGGCAA	CAGCGGCCTG	1680
TTCAATTTCG	GCACCAATAA	CGTTGGCATC	TTCAACGCGG	GCACCGGAAA	CGTCGGCATC	1740
GCAAACCTCG	GCACCGGGAA	CTGGGGTATC	GGGAACCCGG	GTACCGACAA	TACCGGCATC	1800
CTCAATGCTG	GCAGCTACAA	CACGGGCATC	CTCAACGCCG	GCGACTTCAA	CACGGGCTTC	1860
TACAACACGG	GCAGCTACAA	CACCGGCGGC	TTCAACGTCG	GTAACACCAA	CACCGGCAAC	1920
TTCAACGTGG	GTGACACCAA	TACCGGCAGC	TATAACCCGG	GTGACACCAA	CACCGGCTTC	1980
TTCAATCCCG	GCAACGTCAA	TACCGGCGCT	TTCGACACGG	GCGACTTCAA	CAATGGCTTC	2040
TTGGTGGCGG	GCGATAACCA	GGGCCAGATT	GCCATCGATC	TCTCGGTCAC	CACTCCATTC	2100
ATCCCCATAA	ACGAGCAGAT	GGTCATTGAC	GTACACAACG	TAATGACCTT	CGGCGGCAAC	2160
ATGATCACGG	TCACCGAGGC	CTCGACCGTT	TTCCCCAAA	CCTTCTATCT	GAGCGGTTTG	2220
TTCTTCTTCG	GCCCCGTCAA	TCTCAGCGCA	TCCACGCTGA	CCGTTCCGAC	GATCACCCCTC	2280
ACCATCGGCG	GACCGACGGT	GACCGTCCCC	ATCAGCATTG	TCGGTGCTCT	GGAGAGCCGC	2340
ACGATTACCT	TCCTCAAGAT	CGATCCGGCG	CCGGGCATCG	GAAATTCGAC	CACCAACCCC	2400
TCGTCCGGCT	TCTTCAACTC	GGGCACCGGT	GGCACATCTG	GCTTCCAAA	CGTCGGCGGC	2460
GGCAGTTTCA	GCGTCTGGAA	CAGTGTTTGT	AGCAGCGCGA	TAGGGAATTC	GGGTTTCCAG	2520
AACCTCGGCT	CGCTGCAGTC	AGGCTGGGCG	AACCTGGGCA	ACTCCGTATC	GGGCTTTTTC	2580
AACACCAGTA	CGGTGAACCT	CTCCACGCCG	GCCAATGTCT	CGGGCCTGAA	CAACATCGGC	2640
ACCAACCTGT	CCGGCGTGTT	CCGCGGTCCG	ACCGGGACGA	TTTTCAACGC	GGGCCTTGCC	2700
AACCTGGGCC	AGTTGAACAT	CGGCAGCGCC	TCGTGCCGAA	TTCGGCACGA	GTTAGATACG	2760
GTTTCAACAA	TCATATCCGC	GTTTTGCGGC	AGTGCATCAG	ACGAATCGAA	CCCGGGAAGC	2820

GTAAGCGAAT AAACCGAATG GCGGCCTGTC AT

2852

(2) INFORMATION FOR SEQ ID NO:204:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 943 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:204:

Gly Gln Asn Ala Pro Ala Ile Ala Ala Thr Glu Ala Ala Tyr Asp Gln
 1 5 10 15
 Met Trp Ala Gln Asp Val Ala Ala Met Phe Gly Tyr His Ala Gly Ala
 20 25 30
 Ser Ala Ala Val Ser Ala Leu Thr Pro Phe Gly Gln Ala Leu Pro Thr
 35 40 45
 Val Ala Gly Gly Gly Ala Leu Val Ser Ala Ala Ala Ala Gln Val Thr
 50 55 60
 Thr Arg Val Phe Arg Asn Leu Gly Leu Ala Asn Val Arg Glu Gly Asn
 65 70 75 80
 Val Arg Asn Gly Asn Val Arg Asn Phe Asn Leu Gly Ser Ala Asn Ile
 85 90 95
 Gly Asn Gly Asn Ile Gly Ser Gly Asn Ile Gly Ser Ser Asn Ile Gly
 100 105 110
 Phe Gly Asn Val Gly Pro Gly Leu Thr Ala Ala Leu Asn Asn Ile Gly
 115 120 125
 Phe Gly Asn Thr Gly Ser Asn Asn Ile Gly Phe Gly Asn Thr Gly Ser
 130 135 140
 Asn Asn Ile Gly Phe Gly Asn Thr Gly Asp Gly Asn Arg Gly Ile Gly
 145 150 155 160
 Leu Thr Gly Ser Gly Leu Leu Gly Phe Gly Gly Leu Asn Ser Gly Thr
 165 170 175
 Gly Asn Ile Gly Leu Phe Asn Ser Gly Thr Gly Asn Val Gly Ile Gly
 180 185 190
 Asn Ser Gly Thr Gly Asn Trp Gly Ile Gly Asn Ser Gly Asn Ser Tyr
 195 200 205
 Asn Thr Gly Phe Gly Asn Ser Gly Asp Ala Asn Thr Gly Phe Phe Asn
 210 215 220
 Ser Gly Ile Ala Asn Thr Gly Val Gly Asn Ala Gly Asn Tyr Asn Thr
 225 230 235 240
 Gly Ser Tyr Asn Pro Gly Asn Ser Asn Thr Gly Gly Phe Asn Met Gly

200

245								250				255			
Gln	Tyr	Asn	Thr	Gly	Tyr	Leu	Asn	Ser	Gly	Asn	Tyr	Asn	Thr	Gly	Leu
			260					265					270		
Ala	Asn	Ser	Gly	Asn	Val	Asn	Thr	Gly	Ala	Phe	Ile	Thr	Gly	Asn	Phe
		275					280					285			
Asn	Asn	Gly	Phe	Leu	Trp	Arg	Gly	Asp	His	Gln	Gly	Leu	Ile	Phe	Gly
	290					295					300				
Ser	Pro	Gly	Phe	Phe	Asn	Ser	Thr	Ser	Ala	Pro	Ser	Ser	Gly	Phe	Phe
305					310				315						320
Asn	Ser	Gly	Ala	Gly	Ser	Ala	Ser	Gly	Phe	Leu	Asn	Ser	Gly	Ala	Asn
				325				330						335	
Asn	Ser	Gly	Phe	Phe	Asn	Ser	Ser	Ser	Gly	Ala	Ile	Gly	Asn	Ser	Gly
			340					345					350		
Leu	Ala	Asn	Ala	Gly	Val	Leu	Val	Ser	Gly	Val	Ile	Asn	Ser	Gly	Asn
		355					360					365			
Thr	Val	Ser	Gly	Leu	Phe	Asn	Met	Ser	Leu	Val	Ala	Ile	Thr	Thr	Pro
	370					375					380				
Ala	Leu	Ile	Ser	Gly	Phe	Phe	Asn	Thr	Gly	Ser	Asn	Met	Ser	Gly	Phe
385					390					395					400
Phe	Gly	Gly	Pro	Pro	Val	Phe	Asn	Leu	Gly	Leu	Ala	Asn	Arg	Gly	Val
				405				410						415	
Val	Asn	Ile	Leu	Gly	Asn	Ala	Asn	Ile	Gly	Asn	Tyr	Asn	Ile	Leu	Gly
			420				425						430		
Ser	Gly	Asn	Val	Gly	Asp	Phe	Asn	Ile	Leu	Gly	Ser	Gly	Asn	Leu	Gly
		435					440					445			
Ser	Gln	Asn	Ile	Leu	Gly	Ser	Gly	Asn	Val	Gly	Ser	Phe	Asn	Ile	Gly
	450					455				460					
Ser	Gly	Asn	Ile	Gly	Val	Phe	Asn	Val	Gly	Ser	Gly	Ser	Leu	Gly	Asn
465					470				475						480
Tyr	Asn	Ile	Gly	Ser	Gly	Asn	Leu	Gly	Ile	Tyr	Asn	Ile	Gly	Phe	Gly
				485				490						495	
Asn	Val	Gly	Asp	Tyr	Asn	Val	Gly	Phe	Gly	Asn	Ala	Gly	Asp	Phe	Asn
			500				505						510		
Gln	Gly	Phe	Ala	Asn	Thr	Gly	Asn	Asn	Asn	Ile	Gly	Phe	Ala	Asn	Thr
		515					520					525			
Gly	Asn	Asn	Asn	Ile	Gly	Ile	Gly	Leu	Ser	Gly	Asp	Asn	Gln	Gln	Gly
	530					535					540				
Phe	Asn	Ile	Ala	Ser	Gly	Trp	Asn	Ser	Gly	Thr	Gly	Asn	Ser	Gly	Leu
545					550				555						560
Phe	Asn	Ser	Gly	Thr	Asn	Asn	Val	Gly	Ile	Phe	Asn	Ala	Gly	Thr	Gly
				565				570						575	

201

Asn Val Gly Ile Ala Asn Ser Gly Thr Gly Asn Trp Gly Ile Gly Asn
 580 585 590
 Pro Gly Thr Asp Asn Thr Gly Ile Leu Asn Ala Gly Ser Tyr Asn Thr
 595 600 605
 Gly Ile Leu Asn Ala Gly Asp Phe Asn Thr Gly Phe Tyr Asn Thr Gly
 610 615 620
 Ser Tyr Asn Thr Gly Gly Phe Asn Val Gly Asn Thr Asn Thr Gly Asn
 625 630 635 640
 Phe Asn Val Gly Asp Thr Asn Thr Gly Ser Tyr Asn Pro Gly Asp Thr
 645 650 655
 Asn Thr Gly Phe Phe Asn Pro Gly Asn Val Asn Thr Gly Ala Phe Asp
 660 665 670
 Thr Gly Asp Phe Asn Asn Gly Phe Leu Val Ala Gly Asp Asn Gln Gly
 675 680 685
 Gln Ile Ala Ile Asp Leu Ser Val Thr Thr Pro Phe Ile Pro Ile Asn
 690 695 700
 Glu Gln Met Val Ile Asp Val His Asn Val Met Thr Phe Gly Gly Asn
 705 710 715 720
 Met Ile Thr Val Thr Glu Ala Ser Thr Val Phe Pro Gln Thr Phe Tyr
 725 730 735
 Leu Ser Gly Leu Phe Phe Phe Gly Pro Val Asn Leu Ser Ala Ser Thr
 740 745 750
 Leu Thr Val Pro Thr Ile Thr Leu Thr Ile Gly Gly Pro Thr Val Thr
 755 760 765
 Val Pro Ile Ser Ile Val Gly Ala Leu Glu Ser Arg Thr Ile Thr Phe
 770 775 780
 Leu Lys Ile Asp Pro Ala Pro Gly Ile Gly Asn Ser Thr Thr Asn Pro
 785 790 795 800
 Ser Ser Gly Phe Phe Asn Ser Gly Thr Gly Gly Thr Ser Gly Phe Gln
 805 810 815
 Asn Val Gly Gly Gly Ser Ser Gly Val Trp Asn Ser Gly Leu Ser Ser
 820 825 830
 Ala Ile Gly Asn Ser Gly Phe Gln Asn Leu Gly Ser Leu Gln Ser Gly
 835 840 845
 Trp Ala Asn Leu Gly Asn Ser Val Ser Gly Phe Phe Asn Thr Ser Thr
 850 855 860
 Val Asn Leu Ser Thr Pro Ala Asn Val Ser Gly Leu Asn Asn Ile Gly
 865 870 875 880
 Thr Asn Leu Ser Gly Val Phe Arg Gly Pro Thr Gly Thr Ile Phe Asn
 885 890 895
 Ala Gly Leu Ala Asn Leu Gly Gln Leu Asn Ile Gly Ser Ala Ser Cys
 900 905 910

Arg Ile Arg His Glu Leu Asp Thr Val Ser Thr Ile Ile Ser Ala Phe
915 920 925

Cys Gly Ser Ala Ser Asp Glu Ser Asn Pro Gly Ser Val Ser Glu
930 935 940

(2) INFORMATION FOR SEQ ID NO:205:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 53 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:205:

GGATCCATAT GGGCCATCAT CATCATCATC ACGTGATCGA CATCATCGGG ACC

53

(2) INFORMATION FOR SEQ ID NO:206:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:206:

CCTGAATTCA GGCCTCGGTT GCGCCGGCCT CATCTTGAAC GA

42

(2) INFORMATION FOR SEQ ID NO:207:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:207:

GGATCCTGCA GGCTCGAAAC CACCGAGCGG T

31

(2) INFORMATION FOR SEQ ID NO:208:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:208:
CTCTGAATTC AGCGCTGGAA ATCGTCGCGA T 31
- (2) INFORMATION FOR SEQ ID NO:209:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:209:
GGATCCAGCG CTGAGATGAA GACCGATGCC GCT 33
- (2) INFORMATION FOR SEQ ID NO:210:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:210:
GGATATCTGC AGAATTCAGG TTAAAGCCC ATTTGCGA 38
- (2) INFORMATION FOR SEQ ID NO:211:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:211:
CCGCATGCGA GCCACGTGCC CACAACGGCC 30
- (2) INFORMATION FOR SEQ ID NO:212:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:212:

CTTCATGGAA TTCTCAGGCC GGTAAGGTCC GCTGCGG

37

(2) INFORMATION FOR SEQ ID NO:213:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7676 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:213:

TGGCGAATGG GACGCGCCCT GTAGCGGCGC ATTAAGCGCG GCGGGTGTGG TGGTTACGCG	60
CAGCGTGACC GCTACACTTG CCAGCGCCCT AGCGCCCGCT CCTTTCGCTT TCTTCCCTTC	120
CTTTCTCGCC ACGTTCGCCG GCTTTCCCCG TCAAGCTCTA AATCGGGGGC TCCCTTTAGG	180
GTTCCGATTT AGTGCTTTAC GGCACCTCGA CCCCAGAAAA CTTGATTAGG GTGATGGTTC	240
ACGTAGTGGG CCATCGCCCT GATAGACGGT TTTTCGCCCT TTGACGTTGG AGTCCACGTT	300
CTTTAATAGT GGACTCTTGT TCCAACTGG AACAACACTC AACCCTATCT CGGTCTATTC	360
TTTTGATTTA TAAGGGATTT TGCCGATTTT GGCCTATTGG TTAAAAAATG AGCTGATTTA	420
ACAAAAATTT AACGCGAATT TTAACAAAAT ATTAACGTTT ACAATTTTCA GTGGCACTTT	480
TCGGGGAAAT GTGCGCGGAA CCCCTATTTG TTTATTTTTT TAAATACATT CAAATATGTA	540
TCCGCTCATG AATTAATTCT TAGAAAACT CATCGAGCAT CAAATGAAAC TGCAATTTAT	600
TCATATCAGG ATTATCAATA CCATATTTTT GAAAAAGCCG TTTCTGTAAT GAAGGAGAAA	660
ACTCACCGAG GCAGTTCCAT AGGATGGCAA GATCCTGGTA TCGGTCTGCG ATTCCGACTC	720
GTCCAACATC AATACAACCT ATTAATTTCC CCTCGTCAAA AATAAGGTTA TCAAGTGAGA	780
AATCACCATG AGTGACGACT GAATCCGGTG AGAATGGCAA AAGTTTATGC ATTTCTTTCC	840
AGACTTGTTT AACAGGCCAG CCATTACGCT CGTCATCAAA ATCACTCGCA TCAACCAAAC	900
CGTTATTCAT TCGTGATTGC GCCTGAGCGA GACGAAATAC GCGATCGCTG TAAAAGGAC	960
AATTACAAAC AGGAATCGAA TGCAACCGGC GCAGGAACAC TGCCAGCGCA TCAACAATAT	1020
TTTACCTGA ATCAGGATAT TCTTCTAATA CCTGGAATGC TGTTTTCCCG GGGATCGCAG	1080
TGGTGAGTAA CCATGCATCA TCAGGAGTAC GGATAAAATG CTTGATGGTC GGAAGAGGCA	1140

TAAATTCCGT	CAGCCAGTTT	AGTCTGACCA	TCTCATCTGT	AACATCATTG	GCAACGCTAC	1200
CTTTGCCATG	TTTCAGAAAC	AACTCTGGCG	CATCGGGCTT	CCCATACAAT	CGATAGATTG	1260
TCGCACCTGA	TTGCCCGACA	TTATCGCGAG	CCCATTTATA	CCCATATAAA	TCAGCATCCA	1320
TGTTGGAATT	TAATCGCGGC	CTAGAGCAAG	ACGTTTCCCG	TTGAATATGG	CTCATAACAC	1380
CCCTTGATT	ACTGTTTATG	TAAGCAGACA	GTTTTATTGT	TCATGACCAA	AATCCCTTAA	1440
CGTGAGTTTT	CGTTCCACTG	AGCGTCAGAC	CCCGTAGAAA	AGATCAAAGG	ATCTTCTTGA	1500
GATCCTTTTT	TTCTGCGCGT	AATCTGCTGC	TTGCAAACAA	AAAAACCACC	GCTACCAGCG	1560
GTGGTTTGTT	TGCCGGATCA	AGAGCTACCA	ACTCTTTTTC	CGAAGGTAAC	TGGCTTCAGC	1620
AGAGCGCAGA	TACCAAATAC	TGTCCTTCTA	GTGTAGCCGT	AGTTAGGCCA	CCACTTCAAG	1680
AACTCTGTAG	CACCGCCTAC	ATACCTCGCT	CTGCTAATCC	TGTTACCAGT	GGCTGCTGCC	1740
AGTGGCGATA	AGTCGTGTCT	TACCGGGTTG	GACTCAAGAC	GATAGTTACC	GGATAAGGCG	1800
CAGCGGTCGG	GCTGAACGGG	GGGTTCGTGC	ACACAGCCCA	GCTTGGAGCG	AACGACCTAC	1860
ACCGAACTGA	GATACCTACA	GCGTGAGCTA	TGAGAAAGCG	CCACGCTTCC	CGAAGGGAGA	1920
AAGGCGGACA	GGTATCCGGT	AAGCGGCAGG	GTCGGAACAG	GAGAGCGCAC	GAGGGAGCTT	1980
CCAGGGGGAA	ACGCCTGGTA	TCTTTATAGT	CCTGTCGGGT	TCGCCACCT	CTGACTTGAG	2040
CGTCGATTTT	TGTGATGCTC	GTCAGGGGGG	CGGAGCCTAT	GGAAAAACGC	CAGCAACGCG	2100
GCCTTTTTAC	GGTTCTGGC	CTTTTGCTGG	CCTTTTGCTC	ACATGTTCTT	TCCTGCGTTA	2160
TCCCCTGATT	CTGTGGATAA	CCGTATTACC	GCCTTTGAGT	GAGCTGATAC	CGCTCGCCGC	2220
AGCCGAACGA	CCGAGCGCAG	CGAGTCAGTG	AGCGAGGAAG	CGGAAGAGCG	CCTGATGCGG	2280
TATTTTCTCC	TTACGCATCT	GTGCGGTATT	TCACACCGCA	TATATGGTGC	ACTCTCAGTA	2340
CAATCTGCTC	TGATGCCGCA	TAGTTAAGCC	AGTATACACT	CCGCTATCGC	TACGTGACTG	2400
GGTCATGGCT	GCGCCCCGAC	ACCCGCCAAC	ACCCGCTGAC	GCGCCCTGAC	GGGCTTGCT	2460
GCTCCCGGCA	TCCGCTTACA	GACAAGCTGT	GACCGTCTCC	GGGAGCTGCA	TGTGTCAGAG	2520
GTTTTACCG	TCATCACCGA	AACGCGCGAG	GCAGCTGCGG	TAAAGCTCAT	CAGCGTGGTC	2580
GTGAAGCGAT	TCACAGATGT	CTGCCTGTTC	ATCCGCGTCC	AGCTCGTTGA	GTTTCTCCAG	2640
AAGCGTTAAT	GTCTGGCTTC	TGATAAAGCG	GGCCATGTTA	AGGGCGGTTT	TTTCTGTTT	2700
GGTCACTGAT	GCCTCCGTGT	AAGGGGGATT	TCTGTTTCATG	GGGGTAATGA	TACCGATGAA	2760
ACGAGAGAGG	ATGCTCACGA	TACGGGTAC	TGATGATGAA	CATGCCCGGT	TACTGGAACG	2820
TTGTGAGGGT	AAACAACCTGG	CGGTATGGAT	GCGGCGGGAC	CAGAGAAAAA	TCACTCAGGG	2880
TCAATGCCAG	CGCTTCGTTA	ATACAGATGT	AGGTGTTCCA	CAGGGTAGCC	AGCAGCATCC	2940
TGCGATGCAG	ATCCGGAACA	TAATGGTGCA	GGGCGCTGAC	TTCCGCGTTT	CCAGACTTTA	3000

CGAAACACGG AAACCGAAGA CCATTCATGT TGTTGCTCAG GTCGCAGACG TTTTGCAGCA 3060
GCAGTCGCTT CACGTTGCT CGCGTATCGG TGATTCATTC TGCTAACCAG TAAGGCAACC 3120
CCGCCAGCCT AGCCGGGTCC TCAACGACAG GAGCACGATC ATGCGCACCC GTGGGGCCGC 3180
CATGCCGGCG ATAATGGCCT GCTTCTCGCC GAAACGTTTG GTGGCGGGAC CAGTGACGAA 3240
GGCTTGAGCG AGGGCGTGCA AGATTCCGAA TACCGCAAGC GACAGGCCGA TCATCGTCGC 3300
GCTCCAGCGA AAGCGGTCCT CGCCGAAAAT GACCCAGAGC GCTGCCGGCA CCTGTCCTAC 3360
GAGTTGCATG ATAAAGAAGA CAGTCATAAG TCGGCGCAGC ATAGTCATGC CCCGCGCCCA 3420
CCGGAAGGAG CTGACTGGGT TGAAGGCTCT CAAGGGCATC GGTCGAGATC CCGGTGCCTA 3480
ATGAGTGAGC TAACCTTACAT TAATTGCGTT GCGCTCACTG CCCGCTTTCC AGTCGGGAAA 3540
CCTGTCGTGC CAGCTGCATT AATGAATCGG CCAACGCGCG GGGAGAGGCG GTTTGCGTAT 3600
TGGGCGCCAG GGTGGTTTTT CTTTTACCA GTGAGACGGG CAACAGCTGA TTGCCCTTCA 3660
CCGCCTGGCC CTGAGAGAGT TGCAGCAAGC GGTCCACGCT GGTTTGCCCC AGCAGGCGAA 3720
AATCCTGTTT GATGGTGGTT AACGGCGGGA TATAACATGA GCTGTCTTCG GTATCGTCGT 3780
ATCCCACTAC CGAGATATCC GCACCAACGC GCAGCCCGGA CTCGGTAATG GCGCGCATTG 3840
CGCCCAGCGC CATCTGATCG TTGGCAACCA GCATCGCAGT GGGAACGATG CCCTCATTCA 3900
GCATTTGCAT GGTTTGTTGA AAACCGGACA TGGCACTCCA GTCGCCTTCC CGTTCCGCTA 3960
TCGGCTGAAT TTGATTGCGA GTGAGATATT TATGCCAGCC AGCCAGACGC AGACGCGCCG 4020
AGACAGAACT TAATGGGCCC GCTAACAGCG CGATTTGCTG GTGACCCAAT GCGACCAGAT 4080
GCTCCACGCC CAGTCGCGTA CCGTCTTCAT GGGAGAAAAT AATACTGTTG ATGGGTGTCT 4140
GGTCAGAGAC ATCAAGAAAT AACGCCGGA CATTAGTGCA GGCAGCTTCC ACAGCAATGG 4200
CATCCTGGTC ATCCAGCGGA TAGTTAATGA TCAGCCCACT GACGCGTTGC GCGAGAAGAT 4260
TGTGCACCGC CGCTTTACAG GCTTCGACGC CGCTTCGTTT TACCATCGAC ACCACCACGC 4320
TGGCACCCAG TTGATCGGCG CGAGATTTAA TCGCCGCGAC AATTTGCGAC GGCGCGTGCA 4380
GGGCCAGACT GGAGGTGGCA ACGCCAATCA GCAACGACTG TTTGCCCGCC AGTTGTTGTG 4440
CCACGCGGTT GGGAAATGTAA TTCAGTCCG CCATCGCCGC TTCCACTTTT TCCGCGTTTT 4500
TCGAGAAAC GTGGCTGGCC TGTTTACCA CGCGGGAAAC GGTCTGATAA GAGACACCGG 4560
CATACTCTGC GACATCGTAT AACGTTACTG GTTTCACATT CACCACCCTG AATTGACTCT 4620
CTTCCGGGCG CTATCATGCC ATACCGCGAA AGGTTTTGCG CCATTGATG GTGTCCGGGA 4680
TCTCGACGCT CTCCCTTATG CGACTCCTGC ATTAGGAAGC AGCCAGTAG TAGGTTGAGG 4740
CCGTTGAGCA CCGCCCGCGC AAGGAATGGT GCATGCAAGG AGATGGCGCC CAACAGTCCC 4800
CCGGCCACGG GGCCTGCCAC CATACCCACG CCGAAACAAG CGCTCATGAG CCCGAAGTGG 4860

CGAGCCCGAT	CTTCCCCATC	GGTGATGTCG	GCGATATAGG	CGCCAGCAAC	CGCACCTGTG	4920
GCGCCGGTGA	TGCCGGCCAC	GATGCGTCCG	GCGTAGAGGA	TCGAGATCTC	GATCCCGCGA	4980
AATTAATACG	ACTCACTATA	GGGGAATTGT	GAGCGGATAA	CAATTCCCCT	CTAGAAATAA	5040
TTTTGTTTAA	CTTTAAGAAG	GAGATATACA	TATGGGCCAT	CATCATCATC	ATCACGTGAT	5100
CGACATCATC	GGGACCAGCC	CCACATCCTG	GGAACAGGCG	GCGGCGGAGG	CGGTCCAGCG	5160
GGCGCGGGAT	AGCGTCGATG	ACATCCGCGT	CGCTCGGGTC	ATTGAGCAGG	ACATGGCCGT	5220
GGACAGCGCC	GGCAAGATCA	CCTACCGCAT	CAAGCTCGAA	GTGTCGTTCA	AGATGAGGCC	5280
GGCGCAACCG	AGGGGCTCGA	AACCACCGAG	CGGTTCGCCT	GAAACGGGCG	CCGGCGCCGG	5340
TACTGTCGCG	ACTACCCCCG	CGTCGTCGCC	GGTGACGTTG	GCGGAGACCG	GTAGCACGCT	5400
GCTCTACCCG	CTGTTCAACC	TGTGGGGTCC	GGCCTTTCAC	GAGAGGTATC	CGAACGTCAC	5460
GATCACCGCT	CAGGGCACCG	GTTCTGGTGC	CGGGATCGCG	CAGGCCGCCG	CCGGGACGGT	5520
CAACATTGGG	GCCTCCGACG	CCTATCTGTC	GGAAGGTGAT	ATGGCCGCGC	ACAAGGGGCT	5580
GATGAACATC	GCGCTAGCCA	TCTCCGCTCA	GCAGGTCAAC	TACAACCTGC	CCGGAGTGAG	5640
CGAGCACCTC	AAGCTGAACG	GAAAAGTCCT	GGCGGCCATG	TACCAGGGCA	CCATCAAAAC	5700
CTGGGACGAC	CCGCAGATCG	CTGCGCTCAA	CCCCGGCGTG	AACCTGCCCC	GCACCGCGGT	5760
AGTTCCGCTG	CACCGCTCCG	ACGGGTCCGG	TGACACCTTC	TTGTTACCCC	AGTACCTGTC	5820
CAAGCAAGAT	CCCGAGGGCT	GGGGCAAGTC	GCCCGGCTTC	GGCACCACCG	TCGACTTCCC	5880
GGCGGTGCCG	GGTGCGCTGG	GTGAGAACGG	CAACGGCGGC	ATGGTGACCG	GTTGCGCCGA	5940
GACACCGGGC	TGCGTGGCCT	ATATCGGCAT	CAGCTTCCTC	GACCAGGCCA	GTCAACGGGG	6000
ACTCGGCGAG	GCCCAACTAG	GCAATAGCTC	TGGCAATTTT	TTGTTGCCCG	ACGCGCAAAG	6060
CATTACAGGC	GCGGCGGCTG	GCTTCGCATC	GAAAACCCCG	GCGAACCAGG	CGATTTTCGAT	6120
GATCGACGGG	CCCGCCCCGG	ACGGCTACCC	GATCATCAAC	TACGAGTACG	CCATCGTCAA	6180
CAACCGGCAA	AAGGACGCCG	CCACCGCGCA	GACCTTGCAG	GCATTTCTGC	ACTGGGCGAT	6240
CACCGACGGC	AACAAGGCCT	CGTTCCTCGA	CCAGGTTTCAT	TTCCAGCCGC	TGCCGCCCCG	6300
GGTGGTGAAG	TTGTCTGACG	CGTTGATCGC	GACGATTTC	AGCGCTGAGA	TGAAGACCGA	6360
TGCCGCTACC	CTCGCGCAGG	AGGCAGGTAA	TTTCGAGCGG	ATCTCCGGCG	ACCTGAAAAC	6420
CCAGATCGAC	CAGGTGGAGT	CGACGGCAGG	TTCGTTGCAG	GGCCAGTGGC	GCGGCGCGGC	6480
GGGGACGGCC	GCCCAGGCCG	CGGTGGTGCG	CTTCCAAGAA	GCAGCCAATA	AGCAGAAGCA	6540
GGAACCTGAC	GAGATCTCGA	CGAATATTCG	TCAGGCCGGC	GTCCAATACT	CGAGGGCCGA	6600
CGAGGAGCAG	CAGCAGGCGC	TGTCCTCGCA	AATGGGCTTT	GTGCCACAA	CGGCCGCCTC	6660
GCCGCCGTCG	ACCGCTGCAG	CGCCACCCGC	ACCGGCGACA	CCTGTTGCCG	CCCCACCACC	6720

```

GGCCGCCGCC AACACGCCGA ATGCCCAGCC GGGCGATCCC AACGCAGCAC CTCCGCCGGC 6780
CGACCCGAAC GCACCGCCGC CACCTGTCAT TGCCCCAAAC GCACCCCAAC CTGTCCGGAT 6840
CGACAACCCG GTTGGAGGAT TCAGCTTCGC GCTGCCTGCT GGCTGGGTGG AGTCTGACGC 6900
CGCCCCTTC GACTACGGTT CAGCACTCCT CAGCAAAACC ACCGGGGACC CGCCATTTCC 6960
CGGACAGCCG CCGCCGGTGG CCAATGACAC CCGTATCGTG CTCGGCCGGC TAGACCAAAA 7020
GCTTTACGCC AGCGCCGAAG CCACCGACTC CAAGGCCGCG GCCCGGTTGG GCTCGGACAT 7080
GGGTGAGTTC TATATGCCCT ACCCGGGCAC CCGGATCAAC CAGGAAACCG TCTCGCTTGA 7140
CGCCAACGGG GTGTCTGGAA GCGCGTCGTA TTACGAAGTC AAGTTCAGCG ATCCGAGTAA 7200
GCCGAACGGC CAGATCTGGA CGGGCGTAAT CGGCTCGCCC GCGGCGAACG CACCGGACGC 7260
CGGGCCCCCT CAGCGCTGGT TTGTGGTATG GCTCGGGACC GCCAACAACC CGGTGGACAA 7320
GGGCGCGGCC AAGGCGCTGG CCGAATCGAT CCGGCCTTTG GTCGCCCCGC CGCCGGCGCC 7380
GGCACCGGCT CCTGCAGAGC CCGCTCCGGC GCCGGCGCCG GCCGGGGAAG TCGTCCTAC 7440
CCCGACGACA CCGACACCGC AGCGGACCTT ACCGGCCTGA GAATTCTGCA GATATCCATC 7500
ACACTGGCGG CCGCTCGAGC ACCACCACCA CCACCACTGA GATCCGGCTG CTAACAAAGC 7560
CCGAAAGGAA GCTGAGTTGG CTGCTGCCAC CGCTGAGCAA TAACTAGCAT AACCCTTGG 7620
GGCCTCTAAA CGGGTCTTGA GGGGTTTTTT GCTGAAAGGA GGAATATAT CCGGAT 7676

```

(2) INFORMATION FOR SEQ ID NO:214:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 802 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:214:

```

Met Gly His His His His His Val Ile Asp Ile Ile Gly Thr Ser
1           5           10           15
Pro Thr Ser Trp Glu Gln Ala Ala Ala Glu Ala Val Gln Arg Ala Arg
20           25           30
Asp Ser Val Asp Asp Ile Arg Val Ala Arg Val Ile Glu Gln Asp Met
35           40           45
Ala Val Asp Ser Ala Gly Lys Ile Thr Tyr Arg Ile Lys Leu Glu Val
50           55           60
Ser Phe Lys Met Arg Pro Ala Gln Pro Arg Gly Ser Lys Pro Pro Ser
65           70           75           80

```

209

Gly Ser Pro Glu Thr Gly Ala Gly Ala Gly Thr Val Ala Thr Thr Pro
 85 90 95
 Ala Ser Ser Pro Val Thr Leu Ala Glu Thr Gly Ser Thr Leu Leu Tyr
 100 105 110
 Pro Leu Phe Asn Leu Trp Gly Pro Ala Phe His Glu Arg Tyr Pro Asn
 115 120 125
 Val Thr Ile Thr Ala Gln Gly Thr Gly Ser Gly Ala Gly Ile Ala Gln
 130 135 140
 Ala Ala Ala Gly Thr Val Asn Ile Gly Ala Ser Asp Ala Tyr Leu Ser
 145 150 155 160
 Glu Gly Asp Met Ala Ala His Lys Gly Leu Met Asn Ile Ala Leu Ala
 165 170 175
 Ile Ser Ala Gln Gln Val Asn Tyr Asn Leu Pro Gly Val Ser Glu His
 180 185 190
 Leu Lys Leu Asn Gly Lys Val Leu Ala Ala Met Tyr Gln Gly Thr Ile
 195 200 205
 Lys Thr Trp Asp Asp Pro Gln Ile Ala Ala Leu Asn Pro Gly Val Asn
 210 215 220
 Leu Pro Gly Thr Ala Val Val Pro Leu His Arg Ser Asp Gly Ser Gly
 225 230 235 240
 Asp Thr Phe Leu Phe Thr Gln Tyr Leu Ser Lys Gln Asp Pro Glu Gly
 245 250 255
 Trp Gly Lys Ser Pro Gly Phe Gly Thr Thr Val Asp Phe Pro Ala Val
 260 265 270
 Pro Gly Ala Leu Gly Glu Asn Gly Asn Gly Gly Met Val Thr Gly Cys
 275 280 285
 Ala Glu Thr Pro Gly Cys Val Ala Tyr Ile Gly Ile Ser Phe Leu Asp
 290 295 300
 Gln Ala Ser Gln Arg Gly Leu Gly Glu Ala Gln Leu Gly Asn Ser Ser
 305 310 315 320
 Gly Asn Phe Leu Leu Pro Asp Ala Gln Ser Ile Gln Ala Ala Ala Ala
 325 330 335
 Gly Phe Ala Ser Lys Thr Pro Ala Asn Gln Ala Ile Ser Met Ile Asp
 340 345 350
 Gly Pro Ala Pro Asp Gly Tyr Pro Ile Ile Asn Tyr Glu Tyr Ala Ile
 355 360 365
 Val Asn Asn Arg Gln Lys Asp Ala Ala Thr Ala Gln Thr Leu Gln Ala
 370 375 380
 Phe Leu His Trp Ala Ile Thr Asp Gly Asn Lys Ala Ser Phe Leu Asp
 385 390 395 400
 Gln Val His Phe Gln Pro Leu Pro Pro Ala Val Val Lys Leu Ser Asp
 405 410 415

Ala Leu Ile Ala Thr Ile Ser Ser Ala Glu Met Lys Thr Asp Ala Ala
 420 425 430
 Thr Leu Ala Gln Glu Ala Gly Asn Phe Glu Arg Ile Ser Gly Asp Leu
 435 440 445
 Lys Thr Gln Ile Asp Gln Val Glu Ser Thr Ala Gly Ser Leu Gln Gly
 450 455 460
 Gln Trp Arg Gly Ala Ala Gly Thr Ala Ala Gln Ala Ala Val Val Arg
 465 470 475 480
 Phe Gln Glu Ala Ala Asn Lys Gln Lys Gln Glu Leu Asp Glu Ile Ser
 485 490 495
 Thr Asn Ile Arg Gln Ala Gly Val Gln Tyr Ser Arg Ala Asp Glu Glu
 500 505 510
 Gln Gln Gln Ala Leu Ser Ser Gln Met Gly Phe Val Pro Thr Thr Ala
 515 520 525
 Ala Ser Pro Pro Ser Thr Ala Ala Ala Pro Pro Ala Pro Ala Thr Pro
 530 535 540
 Val Ala Pro Pro Pro Pro Ala Ala Ala Asn Thr Pro Asn Ala Gln Pro
 545 550 555 560
 Gly Asp Pro Asn Ala Ala Pro Pro Pro Ala Asp Pro Asn Ala Pro Pro
 565 570 575
 Pro Pro Val Ile Ala Pro Asn Ala Pro Gln Pro Val Arg Ile Asp Asn
 580 585 590
 Pro Val Gly Gly Phe Ser Phe Ala Leu Pro Ala Gly Trp Val Glu Ser
 595 600 605
 Asp Ala Ala His Phe Asp Tyr Gly Ser Ala Leu Leu Ser Lys Thr Thr
 610 615 620
 Gly Asp Pro Pro Phe Pro Gly Gln Pro Pro Pro Val Ala Asn Asp Thr
 625 630 635 640
 Arg Ile Val Leu Gly Arg Leu Asp Gln Lys Leu Tyr Ala Ser Ala Glu
 645 650 655
 Ala Thr Asp Ser Lys Ala Ala Ala Arg Leu Gly Ser Asp Met Gly Glu
 660 665 670
 Phe Tyr Met Pro Tyr Pro Gly Thr Arg Ile Asn Gln Glu Thr Val Ser
 675 680 685
 Leu Asp Ala Asn Gly Val Ser Gly Ser Ala Ser Tyr Tyr Glu Val Lys
 690 695 700
 Phe Ser Asp Pro Ser Lys Pro Asn Gly Gln Ile Trp Thr Gly Val Ile
 705 710 715 720
 Gly Ser Pro Ala Ala Asn Ala Pro Asp Ala Gly Pro Pro Gln Arg Trp
 725 730 735
 Phe Val Val Trp Leu Gly Thr Ala Asn Asn Pro Val Asp Lys Gly Ala

211

740

745

750

Ala Lys Ala Leu Ala Glu Ser Ile Arg Pro Leu Val Ala Pro Pro Pro
755 760 765

Ala Pro Ala Pro Ala Pro Ala Glu Pro Ala Pro Ala Pro Ala Pro Ala
770 775 780

Gly Glu Val Ala Pro Thr Pro Thr Thr Pro Thr Pro Gln Arg Thr Leu
785 790 795 800

Pro Ala

CLAIMS

1. A polypeptide comprising an immunogenic portion of a soluble *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen has an N-terminal sequence selected from the group consisting of:

- (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Cys-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu; (SEQ ID No. 120)
- (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser; (SEQ ID No. 121)
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg; (SEQ ID No. 122)
- (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro; (SEQ ID No. 123)
- (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val; (SEQ ID No. 124)
- (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro; (SEQ ID No. 125)
- (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Thr-Ala-Ala-Ser-Pro-Pro-Ser; (SEQ ID No. 126)
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly; (SEQ ID No. 127)
- (i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Leu-Thr-Ser-Leu-Leu-Asn-Ser-Leu-Ala-Asp-Pro-Asn-Val-Ser-Phe-Ala-Asn; (SEQ ID No. 128) and
- (j) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly; (SEQ ID No. 136)

wherein Xaa may be any amino acid.

2. A polypeptide comprising an immunogenic portion of an *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative

substitutions and/or modifications, wherein said antigen has an N-terminal sequence selected from the group consisting of:

- (a) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID No. 129) and
- (b) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID No. 137), wherein Xaa may be any amino acid.

3. A polypeptide comprising an immunogenic portion of a soluble *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID Nos.: 1, 2, 4-10, 13-25, 52, 99 and 101, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID Nos.: 1, 2, 4-10, 13-25, 52, 99 and 101 or a complement thereof under moderately stringent conditions.

4. A polypeptide comprising an immunogenic portion of a *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID Nos.: 26-51, 138, 139, 163-183 and 201, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID Nos.: 26-51, 138, 139, 163-183 and 201 or a complement thereof under moderately stringent conditions.

5. A DNA molecule comprising a nucleotide sequence encoding a polypeptide according to any one of claims 1-4.

6. An expression vector comprising a DNA molecule according to claim 5.

7. A host cell transformed with an expression vector according to claim 6.

8. The host cell of claim 7 wherein the host cell is selected from the group consisting of *E. coli*, yeast and mammalian cells.
9. A pharmaceutical composition comprising one or more polypeptides according to any one of claims 1-4 and a physiologically acceptable carrier.
10. A pharmaceutical composition comprising one or more DNA molecules according to claim 5 and a physiologically acceptable carrier.
11. A pharmaceutical composition comprising one or more DNA sequences recited in SEQ ID Nos.: 3, 11, 12, 140 and 141; and a physiologically acceptable carrier.
12. A vaccine comprising one or more polypeptides according to any one of claims 1-4 and a non-specific immune response enhancer.
13. A vaccine comprising:
a polypeptide having an N-terminal sequence selected from the group consisting of sequences recited in SEQ ID NO: 134 and 135; and
a non-specific immune response enhancer.
14. A vaccine comprising:
one or more polypeptides encoded by a DNA sequence selected from the group consisting of SEQ ID Nos.: 3, 11, 12, 140 and 141, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID Nos.: 3, 11, 12, 140 and 141; and
a non-specific immune response enhancer.
15. The vaccine of claims 12-14 wherein the non-specific immune response enhancer is an adjuvant.

16. A vaccine comprising one or more DNA molecules according to claim 5 and a non-specific immune response enhancer.

17. A vaccine comprising one or more DNA sequences recited in SEQ ID Nos.: 3, 11, 12, 140 and 141; and a non-specific immune response enhancer.

18. The vaccine of claims 16 or 17 wherein the non-specific immune response enhancer is an adjuvant.

19. A pharmaceutical composition according to any one of claims 9-11, for use in the manufacture of a medicament for inducing protective immunity in a patient.

20. A vaccine according to any one of claims 12-18, for use in the manufacture of a medicament for inducing protective immunity in a patient.

21. A fusion protein comprising two or more polypeptides according to any one of claims 1-4.

22. A fusion protein comprising one or more polypeptides according to any one of claims 1-4 and ESAT-6.

23. A fusion protein comprising one or more polypeptides according to any one of claims 1-4 and the *M. tuberculosis* antigen 38 kD (SEQ ID NO:155).

24. A pharmaceutical composition comprising a fusion protein according to any one of claims 21-23 and a physiologically acceptable carrier.



25. A vaccine comprising a fusion protein according to any one of claims 21-23 and a non-specific immune response enhancer.

26. The vaccine of claim 25 wherein the non-specific immune response enhancer is an adjuvant.



27. A pharmaceutical composition according to claim 24, for use in the manufacture of a medicament for inducing protective immunity in a patient.
28. A vaccine according to claims 25 or 26, for use in the manufacture of a medicament for inducing protective immunity in a patient.
29. A method for detecting tuberculosis in a patient, comprising:
- (a) contacting dermal cells of a patient with one or more polypeptides according to any one of claims 1-4; and
 - (b) detecting an immune response on the patient's skin and therefrom detecting tuberculosis in the patient.
30. A method for detecting tuberculosis in a patient, comprising:
- (a) contacting dermal cells of a patient with a polypeptide having an N-terminal sequence selected from the group consisting of sequences recited in SEQ ID NO: 134 and 135; and
 - (b) detecting an immune response on the patient's skin and therefrom detecting tuberculosis in the patient.
31. A method for detecting tuberculosis in a patient, comprising:
- (a) contacting dermal cells of a patient with one or more polypeptides encoded by a DNA sequence selected from the group consisting of SEQ ID Nos.: 3, 11, 12, 140, 141, 156-160, 189-193, 199, 200 and 203, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID Nos.: 3, 11, 12, 140, 141, 156-160, 189-193, 199, 200 and 203; and
 - (b) detecting an immune response on the patient's skin and therefrom detecting tuberculosis in the patient.
32. The method of any one of claims 29-31 wherein the immune response is induration.

33. A diagnostic kit comprising:
- (a) a polypeptide according to any one of claims 1-4; and
 - (b) apparatus sufficient to contact said polypeptide with the dermal cells of a patient.
34. A diagnostic kit comprising:
- (a) a polypeptide having an N-terminal sequence selected from the group consisting of sequences recited in SEQ ID NO: 134 and 135; and
 - (b) apparatus sufficient to contact said polypeptide with the dermal cells of a patient.
35. A diagnostic kit comprising:
- (a) a polypeptide encoded by a DNA sequence selected from the group consisting of SEQ ID Nos.: 3, 11, 12, 140, 141, 156-160, 189-193, 199, 200 and 203, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID Nos.: 3, 11, 12, 140, 141, 156-160, 189-193, 199, 200 and 203; and
 - (b) apparatus sufficient to contact said polypeptide with the dermal cells of a patient.
36. A diagnostic kit comprising:
- (a) a fusion protein according to any one of claims 21-23; and
 - (b) apparatus sufficient to contact said fusion protein with the dermal cells of a patient.
37. A fusion protein according to claim 23 comprising an amino acid sequence selected from the group consisting of sequences recited in SEQ ID NO: 153 and 209.



1/11

*Fig. 1A-1**Fig. 1A-2*

2/11

*Fig. 1B-1**Fig. 1B-2*

3/11

*Fig. 2A**Fig. 2B*

4/11

M 1 2 3 4 5
68-
43-
29-
18-
14-

Fig. 3B

M 1 2 3 4 5
97-
68-
43-
29-
18-
14-

Fig. 3D

M 1 2 3 4 5
97-
68-
43-
29-
18-
14-

Fig. 3C

M 1 2 3 4 5
97-
68-
43-
29-
18-
14-

Fig. 3E

5/11









Fig. 4A



6/11



*Fig. 4B*

*Fig. 5A**Fig. 5B*



*Fig. 6A**Fig. 6B*

9/11

*Fig. 7A**Fig. 7B*

*Fig. 8A**Fig. 8B*

11/11

*Fig. 9A**Fig. 9B*



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/31, C07K 14/35, A61K 39/04, 48/00, 49/00, C12N 15/62, C07K 19/00, G01N 33/50, 33/60, 33/569, C12N 1/19, 1/20, 1/21, 5/10 // (C12N 1/21, C12R 1:19)	A3	(11) International Publication Number: WO 98/16646 (43) International Publication Date: 23 April 1998 (23.04.98)
(21) International Application Number: PCT/US97/18293 (22) International Filing Date: 7 October 1997 (07.10.97) (30) Priority Data: 08/730,510 11 October 1996 (11.10.96) US 08/818,112 13 March 1997 (13.03.97) US (71) Applicant: CORIXA CORPORATION [US/US]; 1124 Columbia Street, Seattle, WA 98104 (US). (72) Inventors: REED, Steven, G.; 2843 - 122nd Place N.E., Bellevue, WA 98005 (US). SKEIKY, Yasir, A., W.; 8327 - 25th Avenue N.W., Seattle, WA 98107 (US). DILLON, Davin, C.; 21607 N.E. 24th Street, Redmond, WA 98053 (US). CAMPOS-NETO, Antonio; 9308 Midship Court N.E., Bainbridge Island, WA 98021 (US). HOUGHTON, Raymond; 2636 - 242nd Place S.E., Bothell, WA 98021 (US). VEDVICK, Thomas, S.; 124 South 300th Place, Federal Way, WA 98003 (US). TWARDZIK, Daniel, R.; 10195 South Beach Drive, Bainbridge Island, WA 98110 (US). LODES, Michael, J.; 9223 - 36th Avenue S.W., Seattle, WA 98126 (US).		(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US). (81) Designated States: AL, AM, AT, AU, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 8 October 1998 (08.10.98)
(54) Title: COMPOUNDS AND METHODS FOR IMMUNOTHERAPY AND DIAGNOSIS OF TUBERCULOSIS (57) Abstract <p>Compounds and methods for inducing protective immunity against tuberculosis are disclosed. The compounds provided include polypeptides that contain at least one immunogenic portion of one or more <i>M. tuberculosis</i> proteins and DNA molecules encoding such polypeptides. Such compounds may be formulated into vaccines and/or pharmaceutical compositions for immunization against <i>M. tuberculosis</i> infection, or may be used for the diagnosis of tuberculosis.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/18293

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6	C12N15/31	C07K14/35	A61K39/04	A61K48/00	A61K49/00
	C12N15/62	C07K19/00	G01N33/50	G01N33/60	G01N33/569
	C12N1/19	C12N1/20	C12N1/21	C12N5/10	

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 95 01441 A (STATENS SERUMSINSTITUT ;ANDERSEN PETER (DK); ANDERSEN AASE BENGAAR) 12 January 1995</p> <p>see abstract see page 9, line 3 - page 20, line 31 see page 25, line 1-25 see page 32, line 6 - page 35, line 6 see page 43, line 9 - page 50, line 24 see claims</p> <p style="text-align: center;">--- -/--</p>	<p>1,3, 5-10,12, 15,16, 18-29, 32,33, 36,37</p>



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

5 March 1998

Date of mailing of the international search report

16 -07- 1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, T.x. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Macchia, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/18293

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 //(C12N1/21,C12R1:19)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 95 01440 A (STATENS SERUMINSTITUT ;HASLOEV KAARE (DK); ANDERSEN AASE BENGARD) 12 January 1995</p> <p>see abstract see page 1, line 1-30 see page 4, line 8-31 see page 6, line 22 - page 7, line 21 see page 11, line 18 - page 12, line 31 see page 17, line 8-24 see page 24, line 18-25; figure 1 see page 30, line 26 - page 32, line 22 see page 36 - page 43; examples 2-5 see claims</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	<p>1,3, 5-10,12, 15,16, 18-29, 32,33, 36,37</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

5 March 1998

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Macchia, G

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 97/18293

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ANDERSEN P. ET AL.: "Identification of immunodominant antigens during infection with Mycobacterium tuberculosis" SCANDINAVIAN JOURNAL OF IMMUNOLOGY, vol. 36, 1992, pages 823-831, XP002057524 ---	1,3, ~ 5-10,12, 15,16, 18-29, 32,33, 36,37
A	WO 95 31216 A (MEDICAL RES COUNCIL ;LOWRIE DOUGLAS BRUCE (GB)) 23 November 1995 see abstract see claims ---	10,16, 18-20
A	WO 96 15241 A (MERCK & CO INC ;INNOGENETICS NV (BE); LIU MARGARET A (US); MONTGOM) 23 May 1996 see abstract see claims ---	10,16, 18-20
A	WO 96 23885 A (PASTEUR INSTITUT ;LAQUEYRERIE ANNE (FR); MARCHAL GILLES (FR); PESCE) 8 August 1996 ---	
A	WO 92 21758 A (PASTEUR INSTITUT) 10 December 1992 ---	
A	AUSUBEL ET AL.: "Current Protocols in Molecular Biology" 1993 , WILEY & SONS , US, NEW YORK XP002026411 cited in the application see page 10.19.1 - page 10.19.12 ---	
A	YOUNG ET AL.: "Screening of a recombinant mycobacterial DNA library with polyclonal antiserum and molecular weight analysis of expressed antigens" INFECTION AND IMMUNITY, vol. 55, no. 6, June 1987, pages 1421-1425, XP002026410 ---	
A	WO 94 00493 A (KAPOOR ARCHANA ;MUNSHI ANIL (US)) 6 January 1994 ---	
A	FR 2 265 402 A (MITSUI PHARMACEUTICALS) 24 October 1975 ---	
A	FR 2 244 539 A (MITSUI PHARMACEUTICALS) 18 April 1975 ---	
A	ROMAIN ET AL.: "Preparation of Tuberculin antigen L" ANN. INST. PASTEUR/MICROBIOL., vol. 136B, 1985, pages 235-248, XP002026409 ---	
	--- -/--	

INTERNATIONAL SEARCH REPORT

Intern: al Application No
PCT/US 97/18293

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 97 09428 A (CORIXA CORP) 13 March 1997 see abstract see page 158 - page 163; claims ---	1,3, 5-10,12, 15,16, 18-22, 24-29, 32,33,36
P,X	WO 97 09429 A (CORIXA CORP) 13 March 1997 see abstract see page 173-181; claims -----	1,3,5-8, 21,22

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/ 18293

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 29 and 32
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See annex

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

See annex subject 1 -

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1, 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

A polypeptide comprising an immunogenic portion of a soluble M. tuberculosis antigen or a variant, having an N-terminal aminoacid sequence as in Seq.ID:120. A DNA molecule encoding said polypeptide as in Seq.ID:101. An expression vector comprising said DNA molecule, a host transformed with said expression vector. A pharmaceutical composition or vaccine comprising said polypeptide or said DNA molecule. Fusion protein comprising said polypeptide and pharmaceutical composition or vaccine thereof. A diagnostic kit comprising said polypeptide or fusion protein.

2. Claims: 1, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:121.

3. Claims: 1, 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:122 and 25.

4. Claims: 1, 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:123 and 24.

5. Claims: 1, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:124.

6. Claims: 1, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:125.

7. Claims: 1, 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:126 and 52.

8. Claims: 1, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:127.

9. Claims: 1, 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:128 and 99.

10. Claims: 1, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:136.

11. Claims: 2, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:129.

12. Claims: 2, 5-10, 12, 15, 16, 18-29, 31-33,
35-37 all partially.

Same as invention 1 but for Seq.ID:137 and 203.

13. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:1.

14. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:2.

15. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:4 and 17.

16. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:5.

17. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:6.

18. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:7.

19. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:8.

20. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially

Same as invention 1 but for Seq.ID:9.

21. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:10 and 13.

22. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:14.

23. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:15 and 153.

24. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:16.

25. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:18.

26. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:19.

27. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:20.

28. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:21.

29. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:22.

30. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:23.

31. Claims: 3, 5-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:99.

32. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:26.

33. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:27.

34. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:28.

35. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:29.

36. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:30.

37. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:31.

38. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:32.

39. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:33.

40. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:34.

41. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:35.

42. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:36.

43. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:37.

44. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:38.

45. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:39.

46. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:40.

47. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:41.

48. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:42.

49. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:43, 44 and 183.

50. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:45.

51. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:46 and 153.

52. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:47.

53. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:48.

54. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:49.

55. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:50.

56. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:51.

57. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:138.

58. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:139.

59. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:163.

60. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:165.

61. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:166.

62. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:167.

63. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:168.

64. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:169 and 170.

65. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:171 and 172.

66. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:173 and 174.

67. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:175 and 176.

68. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:177 and 178.

69. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:179 and 180.

70. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

Same as invention 1 but for Seq.ID:181 and 182.

71. Claims: 4-10, 12, 15, 16, 18-29, 32, 33, 36,
37 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 1 but for Seq.ID:201.

72. Claims: 11, 14, 15, 17-20, 31, 32, 35 all partially.

Pharmaceutical composition or vaccine comprising a DNA sequence as in Seq.ID:3. Vaccine comprising a polypeptide encoded by said Seq.ID:3, complement or hybridizing sequences thereof. Use of said polypeptide in diagnostic.

73. Claims: 11, 14, 15, 17-20, 31, 32, 35 all partially.

Same as invention 72 but for Seq.ID:11.

74. Claims: 11, 14, 15, 17-20, 31, 32, 35 all partially.

Same as invention 72 but for Seq.ID:12.

75. Claims: 11, 14, 15, 17-20, 31, 32, 35 all partially.

Same as invention 72 but for Seq.ID:140.

76. Claims: 11, 14, 15, 17-20, 31, 32, 35 all partially.

Same as invention 72 but for Seq.ID:141.

77. Claims: 13, 15, 20, 30, 32, 34 all partially.

Vaccine comprising a polypeptide having an N-terminal sequence as in Seq.ID:134. Use of said polypeptide in diagnostic.

78. Claims: 13, 15, 20, 30, 32, 34 all partially.

Same as invention 77 but for Seq.ID:135.

79. Claims: 31, 32, 35 all partially.

Use in diagnostic of a polypeptide encoded by a DNA sequence as in Seq.ID:156.

80. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:157.

81. Claims: 31, 32, 35 all partially.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Same as invention 79 but for Seq.ID:158.

82. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:159 and 160.

83. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:189.

84. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:190.

85. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:191.

86. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:192.

87. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:193.

88. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:199 and 200.

89. Claims: 31, 32, 35 all partially.

Same as invention 79 but for Seq.ID:203.

Polypeptides comprising an immunogenic portion of a soluble M. tuberculosis antigen are well documented in the prior art.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/18293

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9501441 A	12-01-95	AU 682879 B	23-10-97
		AU 7068894 A	24-01-95
		CA 2165949 A	12-01-95
		EP 0706571 A	17-04-96
		NZ 267984 A	22-09-97

WO 9501440 A	12-01-95	AU 685133 B	15-01-98
		AU 7068694 A	24-01-95
		EP 0749486 A	27-12-96

WO 9531216 A	23-11-95	AU 2529095 A	05-12-95
		EP 0804230 A	05-11-97
		JP 10500128 T	06-01-98

WO 9615241 A	23-05-96	US 5736524 A	07-04-98
		AU 4110296 A	06-06-96
		CZ 9701451 A	15-10-97
		EP 0792358 A	03-09-97
		FI 972034 A	11-07-97
		HU 77028 A	02-03-98
		NO 972196 A	11-07-97
		PL 320091 A	15-09-97
		SK 59797 A	14-01-98
		ZA 9509608 A	29-05-96

WO 9623885 A	08-08-96	US 5714593 A	03-02-98
		AU 4667596 A	21-08-96
		CA 2210928 A	08-08-96
		EP 0807178 A	19-11-97

WO 9221758 A	10-12-92	FR 2677365 A	11-12-92
		CA 2110389 A	10-12-92
		EP 0589943 A	06-04-94
		JP 6508513 T	29-09-94

WO 9400493 A	06-01-94	US 5330754 A	19-07-94
		AU 689075 B	26-03-98
		AU 4651193 A	24-01-94
		EP 0649435 A	26-04-95
		JP 7508649 T	28-09-95

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/18293

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9400493 A		US 5559011 A	24-09-96
FR 2265402 A	24-10-75	NONE	
FR 2244539 A	18-04-75	NONE	
WO 9709428 A	13-03-97	AU 7158696 A	27-03-97
WO 9709429 A	13-03-97	AU 7158796 A	27-03-97
		EP 0850305 A	01-07-98